

Fasdol Suspension 250mg/5ml (Strawberry)**Fasdol Suspension 250mg/5ml (Orange)**

Paracetamol 250mg /5ml

What is in this leaflet

1. What Fasdol suspension is used for
2. How Fasdol suspension works
3. Before you take Fasdol suspension
4. How to use Fasdol suspension
5. While you are using Fasdol suspension
6. Side effects
7. Storage and Disposal
8. Product Description
9. Manufacturer & Product Registration Holder
10. Date of revision

1. What with Fasdol suspension is used for

Fasdol suspension is used for the relief of fever, headache and symptoms of cold and flu, toothache, discomfort of teething and fever after vaccination.

2. How Fasdol suspension works

Fasdol suspension contains Paracetamol which is a centrally acting analgesic and antipyretic with minimal anti-inflammatory properties.

3. Before you use Fasdol suspension*-When you must not use it*

- If you have ever had an allergic reaction to paracetamol or any of the other ingredients in the product.
- If you are taking other medicines containing paracetamol.

-Before you start to use it

Check with your doctor

- If you suffer from liver or kidney problems, including alcoholic liver disease
- If you suffer from mild arthritis and need to take painkillers every day
- If you have a weak kidney or liver.

-Taking another medicines

Please talk to your doctor or pharmacist, if you are taking these medicines:

- if you are taking blood thinning drugs (anticoagulants e.g. warfarin) and you need to take a pain reliever on a daily basis
- If you have recently taken any other medicines, including medicines obtained without prescription.

-Pregnancy and breast-feeding

Use in pregnancy:

- Considered to be the analgesic of choice in pregnant patients.
- Although it crosses placenta, paracetamol is considered to be safe in normal therapeutic doses for short-term use as a minor analgesic/antipyretic in pregnancy.

Use in lactation:

- Excreted in breast milk.
- Maternal ingestion of paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

-Driving and using machines

Fasdol suspension is unlikely to impair a patient's ability to drive or use machinery

4. How to use Fasdol suspension*How much to use it*

Children

1-3 years: 3 -5ml

4-6 years: 5-7 ml

7-9 years: 7-10 ml

10-12 years: 10-15 ml

Shake well before use.

Do not give more than 4 times daily

-When to use it

Fasdol suspension should be taken after meal.

-How long to use it

The duration of treatment with Fasdol suspension is determined by your doctor.

-If you forget to use it

Take the next dose when needed. Do not take a double dose to make up for a forgotten dose.

-If you use overdose

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

5. While you are using Fasdol suspension*-Thing you must do*

Keep this leaflet. You may need to read it again. If you have any further questions, ask your doctor or pharmacist

-Things you must not do

Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

-Things to be careful of

Do not take more than the recommended dose. If your symptoms continue or your headache becomes persistent, meet your doctor.

6. Side Effect

Adverse effects of paracetamol are rare and usually mild, although haematological reactions have been reported. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Check with your doctor or pharmacist for further information. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.gov.my (Consumer → Reporting Side Effects to Medicines [ConSERF] or Vaccines [AEFI])

7. Storage and Disposal*Storage*

Store at below 30°C in dry place away from heat, moisture and light. Store it in the original package. Keep this medicine in a safe place out of the sight and reach of children.

Disposal

Medicine should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Fasdol Suspension 250mg/5ml (Strawberry)**Fasdol Suspension 250mg/5ml (Orange)**Paracetamol 250mg /5ml

8. Product Description-What it looks like

Fasdol Suspension strawberry: A pink rose coloured and strawberry flavour suspension.

Fasdol Suspension orange: Orange coloured and orange flavour suspension.

-Ingredients

Active ingredient: paracetamol

Other ingredients:

Strawberry: Methyl paraben, sodium saccharin, refined sugar, sorbitol solution, xanthan gum, strawberry flavour, colour pink rose and purified water.

Orange: Methyl paraben, sodium saccharin, refined sugar, sorbitol solution, xanthan gum, banana flavour, orange flavour, colour orange and purified water.

-Registration Number:

Fasdol Suspension 250mg/5ml (Strawberry)
MAL XCZ

Fasdol Suspension 250mg/5ml (Orange)
MAL XCZ

9. Manufacturer & Product**Registration Holder**-Product registration holder:

Advance Pharma Sdn Bhd 539885-W
28, Jalan BP6/6, Bandar Bukit Puchong,
47120 Puchong, Selangor.

-Manufacturer:

AV Manufacturing Sdn. Bhd. (667760-v)
Lot 10621, Jalan Permata 2,
Arab Malaysian Industrial Park
71800 Nilai, Negeri Sembilan.

10. Date of revision

31 Jan 2020

Serial number

NPRA (R2) 19/226