

POLYLAB T-EUSOL (SOLUTION A & B)

Chlorinated Lime (2.5% w/v)

What is in this leaflet

1. What T-Eusol (Solution A & B) is used for
2. How T-Eusol works
3. Before using T-Eusol
4. How to use T-Eusol
5. While you are using T-Eusol
6. Side Effects
7. Storage and Disposal of T-Eusol
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What T-Eusol is used for

For cleansing of wound and ulcer.

How T-Eusol works

Inhibiting bacterial and fungal growth.

Before using T-Eusol

When you must not use it

You are allergic to ingredients present in T-Eusol Solution A and B.

Before you start to use it

Please consult your doctor or pharmacist before using it.

Taking other medicines

Please inform your doctor or pharmacist if you are taking any other medications.

How to use T-Eusol

How much to use

Mix equal volumes of T-Eusol (Solution A) and T-Eusol (Solution B) before use.

When to use

Follow the directions on the label carefully or as instructed by your doctor or pharmacist.

How long to use

Please refer to your doctor or pharmacist.

If you forget to use it

Apply as soon as you remember. If it is too near the next dose timing, apply the next dose at the normal time. Do not apply extra dose to compensate for the missed dose.

If you use too much (overdose)

Seek medical attention as soon as possible.

While you are using T-Eusol

Things you must do

Consult your doctor or pharmacist if symptoms persist. Doctor or pharmacist will assess your situation and advise on the need to continue application of T-Eusol.

Things you must not do

- Do not ingest internally. Ingestion causes irritation and damage of mucous membrane along with pain and vomiting.
- Do not use T-Eusol to treat other complaints unless advised by your doctor.

Things to be careful of

- The antimicrobial activity of chlorine is diminished by presence of organic matter.
- Inhalation of fumes can cause breathing problems (respiratory tract irritation and pulmonary oedema).
- Avoid contact with eyes.

Side Effects

May give rise to swelling in a certain area (localised oedema), burning sensation in wounds and irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of T-Eusol

Storage

- Store in a dry place below 30°C.
- Protect from light.
- Keep container tightly closed.

Disposal

T-Eusol should not be disposed via wastewater or household waste. Ask advice from pharmacist on the correct disposal of the product once no longer required. This will help to protect the environment.

Product Description

What it looks like

Packed in two different bottles;

Solution A: Contains 2.5% chlorinated lime in an aqueous buffer solution.

Solution B: Contains 1.2% glacial acetic acid.

Contains not less than 0.25% w/v available chlorine when freshly prepared by mixing equal volumes of solution A and B.

Active ingredient:

- Chlorinated lime

Inactive ingredients:

- Sodium acetate
- Glacial acetic acid
- Deionised water

MAL Number

MAL07081672XCS

Manufacturer

Sanchuan Medical Sdn. Bhd.
No.72, Lengkok Rishah 2,
Taman Rishah,
Kawasan Perindustrian Silibin,
30100 Ipoh, Perak.

Product Registration Holder

Polylab Biotech Sdn. Bhd.
6A, Jalan Kaskas 3,
Taman Cheras,
56100 Kuala Lumpur,
W. P. Kuala Lumpur.

Date of Revision

08/02/2021

Serial Number

NPRA(R4/1)060519/00117