

AJALORIN ORODISPERSIBLE TABLET

Desloratadine 2.5mg & 5mg

What is in this leaflet

1. What Ajalorin Orodispersible Tablet is used for?
2. How Ajalorin Orodispersible Tablet works?
3. Before you use Ajalorin Orodispersible Tablet?
4. How to use Ajalorin Orodispersible Tablet?
5. While you are using Ajalorin Orodispersible Tablet
6. Side Effect
7. Storage and Disposal of Ajalorin Orodispersible Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Ajalorin Orodispersible Tablet is used for:

Ajalorin Orodispersible Tablet is used for the relief of symptoms relating to allergic rhinitis and urticarial.

How Ajalorin Orodispersible Tablets works:

Ajalorin Orodispersible Tablet contains the active ingredient desloratadine which known as *antihistamines*. During an allergic reaction, the body produces a chemical called *histamine*, which causes allergy symptoms. Desloratadine works by blocking the action of histamine in the body and relieve the symptoms.

Before you use Ajalorin Orodispersible Tablet:

-When you must not use it:

- If you are allergic to the active substance

deslaratadine, loratadine or any of the other ingredients of this medicine.

- If you are pregnant or are breast-feeding.
- If you are under 12 years old of age.
- If you have case of severe renal insufficiency.
- This product contains aspartame, it may be harmful to people with phenylketonuria.

-Before you start to use it:

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

-Taking other medicines

There is no clinical relevant interactions were observed in trials with desloratadine.

How to use Ajalorin Orodispersible Tablet

-How much to use

Ajalorin 2.5mg Orodispersible Tablet;

- Children (6-11years old); 1 tablet per day.
- Adults/adolescent (12years old and above); 2 tablets per day.

Ajalorin 5mg Orodispersible Tablet;

- Adults/adolescent (12 years old and above); 1 tablet per day.

-When to use it

Always take this medicines exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

You can take Ajalorin Orodispersible Tablet with or without food.

-How long to use it

The duration you need to take Ajalorin Orodispersible Tablet depends on your symptoms and doctor advice.

-If you forget to use it

If you forget to take your tablet you should take it as soon as you remember on the same day. If you do not remember on the same day, take your normal dose on the next day.

Do not take a double dose to make up for a forgotten dose. If you are unsure about what to do ask your doctor or pharmacist.

-If you use too much (overdose)

If you take more than the prescribed, get medical help immediately.

While you are using Ajalorin Orodispersible Tablet

-Things you must do

It is important that you complete the course of treatment even if you begin to feel better after a few days. If you stop taking Ajalorin Orodispersible Tablet too soon your allergic may not be completely cured and the symptoms may return or your condition may get worse

-Things you must not do

- Do not give this medicine to children under the age of 12 because efficacy and safety have not been established for this age group.

-Things to be careful of

- Hypersensitivity to desloratadine, loratadine or to any of the excipients.
- Pregnancy and lactation.
- If you are below 12 years old.

Consumer Medication Information Leaflet

AJALORIN ORODISPERSIBLE TABLET

Desloratadine 2.5mg & 5mg

(RiMUP)

- If you have case of severe renal insufficiency.
- This product contains aspartame, it may be harmful to people with phenylketonuria.

Side Effect

- Common; fatigue, dry mouth and headache.
- Rarely; hallucination, dizziness, palpitations, abdominal pain, nausea, vomiting, diarrhoea and hypersensitivity reactions. Please consult your doctor for further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public Reporting Medicinal Problems/ Side Effects/AEFI/Vaccine Safety).

Storage and Disposal of Ajalorin Orodispersible Tablet

-Storage

Store below 30°C in the original package to protect from light

-Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use.

Product Description

-What it looks like

Ajalorin 2.5mg Orodispersible Tablet

Red-brick color, round, flat tablets with bevel edges and embossed with "2.5"

Ajalorin 2.5mg Orodispersible Tablet

Red-brick color, round, flat tablets with bevel edges and embossed with "5"

-Ingredients

Active Ingredient
Desloratadine

Inactive Ingredient
Polacrillin potassium, Citric Acid Monohydrate, Red Iron Oxide E172, Magnesium Stearate, Croscarmellose Sodium, Tutti Frutti Flavor, Aspartame, Microcrystalline Cellulose, Mannitol, Purified water.

-MAL number:

Ajalorin 2.5mg Orodispersible tablet: MAL19016047ACZ
Ajalorin 5mg Orodispersible tablet: MAL19016044ACZ

Manufacturer:

Genepharm S.A
18km Marathon Avenue
15351 Pallini Attikis
Greece

Product Registration

Holder:

AJ Research & Pharma Sdn.Bhd.
Unit 27.1, Level 27,
Menara Felda, Platinum Park, No. 11, Perisaran KLCC, 50088 Kuala Lumpur Malaysia.

Date of revision:

April 2023

Serial number

NPRA (R1/1) 11042023/074