

AJENAFIL CHEAWABLE TABLET

Sildenafil Citrate 25mg, 50mg and 100mg

What is in this leaflet

1. What Ajenafil Chewable Tablet is used for
2. How Ajenafil Chewable Tablet it works
3. Before you use Ajenafil Chewable Tablet
4. How to use Ajenafil Chewable Tablet
5. While you are using Ajenafil Chewable Tablet
6. Side Effect
7. Storage and Disposal Ajenafil Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Ajenafil Chewable Tablet is used for:

Treatment for adult men with erectile dysfunction, sometimes known as impotence. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity.

How Ajenafil Chewable Tablet works:

Ajenafil contains the active substance sildenafil which belongs to a group of medicines called phosphodiesterase type 5 (PDE5) inhibitors. It works by helping to relax the blood vessels in your penis, allowing blood to flow into your penis when you get sexually excited. Ajenafil will only help you to get an erection if you are sexually stimulated.

Before you use Ajenafil Chewable Tablet:

When you must not use it:

- Hypersensitivity to the active substance or to any of the excipients.
- Sexual activity is inadvisable.
- Have loss of vision in one eye because of Non-Arteritic

Anterior Ischaemic Optic Neuropathy (NAION).

- Severe hepatic impairment, hypotension (blood pressure <90/50 mmHg), recent history of stroke or myocardial infarction and known hereditary degenerative retinal disorders such as *retinitis pigmentosa*.

Before you start to use it:

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking other medicines

Tell your healthcare provider if you take any of the following:

- CYP3A4 inhibitor such as Ketoconazole, erythromycin and cimetidine.
- Medicines called HIV protease inhibitors, such as ritonavir, saquinavir
- Antacid such as magnesium hydroxide/aluminium hydroxide.
- CYP2C9 inhibitors such as tolbutamide, warfarin, phenytoin and CYP2D6 inhibitors such as thiazide.
- Rifampicin and barbiturates.
- Nicorandil and other nitrates.
- Alpha-blocker therapy

How to use Ajenafil Chewable Tablet

How much to use

Use in adults:

The recommended dose is 50 mg. The tablets should be chewed before swallowed.

Based on efficacy and toleration, the dose may be increased to 100 mg or decreased to 25 mg. The maximum recommended

dose is 100 mg. The maximum recommended dosing frequency is once per day. If Ajenafil is taken with food, the onset of activity may be delayed compared to the fasted state.

Use in the elderly:

Dosage adjustments are not required in elderly patients.

When to use it

Taken as needed approximately one hour before sexual activity.

How long to use it

Take as advised by your doctor.

If you forget to use it

You can take it one hour before sexual activity.

If you use too much (overdose)

Doses above 100 mg do not increase the efficacy. Contact your doctor if you take more tablets than you should.

While you are using Ajenafil Chewable Tablet

Things you must do

Take 1 hour before the sexual activity.

Things you must not do

- You should not use Ajenafil with any other oral or local treatments for erectile dysfunction.
- You should not take Ajenafil Chewable Tablet if you do not have erectile dysfunction.
- Ajenafil Chewable Tablet contains Aspartame. You should not take if you are a phenylketonuria patients.
- Ajenafil Chewable Tablet contains lactose. If you have hereditary problems of

galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Things to be careful of

- Your cardiovascular status.
- If you have deformation of penis.
- If you have sickle cell anaemia, multiple myeloma or leukaemia.

Side Effect

Headache, flushing, dyspepsia, visual disorders, nasal congestion, dizziness, nausea and visual color distortion.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Ajenafil Chewable Tablet

Storage

Store below 30°C in the original package to protect from light

Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use.

Product Description

What it looks like

25 mg: White, triangular, with a side of 7.3 mm, biconvex, embossed with “25” on one side.

50 mg: White, triangular, with a side of 8.8 mm, biconvex, embossed with “50” on one side.

100 mg: White, triangular, with a side of 11.8 mm,

biconvex, embossed with “100” on one side.

Ingredients

Active Ingredient
Sildenafil Citrate

Inactive Ingredient
Polacrillin potassium, purified water, magnesium stearate, silica colloidal anhydrous, aspartame, croscarmellose sodium, peppermint flavor, lactose monohydrate, povidone K-30, hydrochloride acid/potassium hydroxide.

MAL no:

Ajenafil 25mg Chewable
Tablet: MAL18046144ACZ
Ajenafil 50mg Chewable
Tablet: MAL18046145ACZ
Ajenafil 100mg Chewable
Tablet: MAL18046146ACZ

Manufacturer:

Genepharm S.A
18 Km Marathon Avenue -
15351 Pallini
Greece

Product Registration

Holder:

AJ Research & Pharma
Sdn.Bhd.
Unit 27.1, Level 27, Menara
Felda, Platinum Park, No.
11, Persiaran KLCC,
50088 Kuala Lumpur
Malaysia.

Date of revision:

December 2022

Serial Number:

NPRA (R1/1) 09022023/040