

COXA 500 TABLET

Consumer Medication Information Leaflet (RiMUP)

Paracetamol 500mg

What is in this leaflet

1. What Coxa 500 Tablet is used for
2. How Coxa 500 Tablet works
3. Before you use Coxa 500 Tablet
4. How to use Coxa 500 Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Coxa 500 Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Coxa 500 Tablet is used for

Coxa 500 tablets is used for the relief of fever and reduce pain.

How Coxa 500 Tablet works

Coxa Tablet contains Paracetamol which has analgesic (pain relief) and antipyretic (anti-fever) properties.

Before you use Coxa 500 Tablet

- When you must not use it

Do not take Coxa 500 Tablet if:

- You have ever had an allergic reaction to Paracetamol or to any of the other ingredients
- You are taking other medicines containing paracetamol.

If you are not sure whether you should take this medicine, ask your doctor or pharmacist for advice before taking any medicine.

- Before you start use it

Tell your doctor or pharmacist if

- You have liver or kidney disease.
- You are pregnant or breastfeeding. Ill effects in human have not been reported when taking paracetamol during pregnancy or breastfeeding.
- You are taking other medicine that affect the liver

WARNING

This preparation contains
PARACETAMOL.

Do not take any other paracetamol containing medicines at the same time.

- Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking if you are taking cholestyramine, metoclopramide, domperidone, warafin, chloramphenicol, non-steroidal anti-inflammatory drug (NSAIDs), alcohol, hepatic enzyme inducers, hepatotoxic medications.

How to use Coxa 500 Tablet

- How much to use

- Adults and children over 12 years: 1 - 2 tablets three to four times daily as required. Not more than 8 tablets in 24 hours unless instructed by a doctor.
- Children under 12 years: Not recommended.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Follow the direction on the label carefully and as instructed by your doctor or pharmacist

- How long to use it

As instructed by your doctor or pharmacist

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose.

Do not take a double dose to make up for the missed dose.

If you are not sure, consult your doctor or pharmacist on what you should do when you forget to use this medicine.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause skin to turn pale, nausea or vomiting, stomach cramps or pain, loss of appetite and liver damage.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Coxa 500 Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Coxa 500 Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Avoid drinking alcohol while taking this medication.

Side effects

Like all medicines, Coxan Tablet can cause side effects, although not everybody gets them. Side effects of paracetamol are usually mild, though allergic and hematological reactions had been reported. Stop taking the medicine and tell your doctor immediately if you experience allergic reactions including skin rashes, swelling of the deep layers of skin, Steven Johnson Syndrome/Toxic Epidermal.

Allergy Alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers→

Reporting Side Effects to Medicines
(ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Coxa 500 Tablet

- Storage

Store below 30°C. Protect from light.
Keep out of reach of children.

- Disposal

Medicines should not be disposed of via
wastewater or household waste. Ask
your pharmacist how to dispose of
medicines no longer required. These
measures will help to protect the
environment.

Product Description

- What it looks like

Coxa Tablet is a white round, uniform
and scored on one side. The score line is
not intended to divide the tablet into
equal doses.

- Ingredients

- Active ingredient
Paracetamol 500mg

- Inactive ingredients
Pregelatinized starch, stearic acid,
Povidone, and purified water.

- MAL number:

MAL22036020XCZ

Manufacturer

Medinfar Manufacturing, S.A.
Parque Industrial Armando Martins
Tavares
Rua Outeiro da Armada, 5
Condeixa-a-Nova
3150-194 Sebal
Portugal

Product Registration Holder

Alpharme Plc Sdn Bhd.
Lot 6-01, 6th Floor, Choo Plaza,
41 Lorong Abu Siti, 10400
Georgetown, Pulau Pinang,
Malaysia.

Date of revision

21/02/2024
MYX066RE (01)

Serial Number

NPRA (R2) 21/081