

ULTRACET® TABLET

Tramadol hydrochloride / acetaminophen (37.5mg/325mg)

What is in this leaflet

1. What ULTRACET® is used for
2. How ULTRACET® works
3. Before you use ULTRACET®
4. How to use ULTRACET®
5. While you are using it
6. Side effects
7. Storage and disposal of ULTRACET®
8. Product description
9. Manufacturer and product registration holder
10. Date of revision

What ULTRACET® is used for

ULTRACET® is used to treat moderate pain to severe pain.

How ULTRACET® works

ULTRACET® tablets have a combination of two pain relievers – tramadol hydrochloride (an opioid analgesic) and acetaminophen (same as paracetamol).

Before you use ULTRACET®

- When you must not use it

- You are less than 12 years old;
- You have slow or shallow breathing, or other breathing problems;
- You are pregnant;
- You are breastfeeding.

You must not take ULTRACET® if you have previously had an allergic reaction (hypersensitivity) to tramadol hydrochloride or acetaminophen, or to any other opioid pain relievers. Please tell your doctor if this applies to you. Hypersensitivity can be recognized for instance by skin rash, itching, shortness of breath or swollen face. If any of these occur, contact your doctor right away.

Do not give ULTRACET® to a child under 18 years of age that has just an operation to remove their tonsils or adenoids.

Do not take ULTRACET® if you are taking:

- Sleeping pills e.g. hypnotics;
- Medicines that affect your mood or emotions e.g. narcotics and psychotropic drugs;
- Pain relievers that affect your central nervous system e.g. centrally acting analgesics and opioids;

- Antidepressants such as Monoamine Oxidase Inhibitors (MAO inhibitors) currently or within the last 14 days.

Pregnancy and lactation

If you are, or think you may be, pregnant, you should inform your doctor, who will decide if you can take ULTRACET®. Opioid medicines can slow the breathing of a newborn child. Prolonged use of ULTRACET® during pregnancy can cause withdrawal symptoms in your new-born baby that could be life-threatening if not recognized and treated. This risk is increased in the last trimester of pregnancy.

Breastfeeding

You are advised not to take ULTRACET® if you are breastfeeding, because ULTRACET® may end up in your milk.

- Before you start use it

BEFORE you use ULTRACET®, be sure to tell your doctor or pharmacist if you:

- suffer from serious liver or kidney disease,
- have diabetes,
- are over 65 years of age,
- have abdominal problems,
- have had a previous head injury,
- have chronic alcoholism,
- have sepsis (when bacteria and their toxins circulate in the blood leading to organ damage)
- suffer from malnutrition
- are pregnant or plan to become pregnant,
- are breast-feeding,

if you are taking any other medications including natural health products, prescription drugs, salicylates, other pain and fever relief medications or nonsteroidal anti-inflammatory drugs (NSAIDs).

This will help your doctor to decide whether you should use ULTRACET® and what extra care should be taken during its use.

- Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines. This includes

medicines that you buy without a prescription or herbal medicines. In particular, tell your doctor or pharmacist if you are taking any of the following medicines as ULTRACET® may affect how well they work:

- Monoamine Oxidase Inhibitors (MAOIs), used to treat mental conditions such as depression e.g. phenelzine, tranylcypromine. You should not take ULTRACET® with MAOIs or within 14 days of having last taken them;
- Selective Serotonin Reuptake Inhibitors (SSRI's), used to treat depression or Parkinson's Disease;
- Drugs used to treat depression e.g. mirtazapine; tricyclic antidepressants; trazadone;
- Triptans, used to treat migraines;
- 5-HT₃ receptor antagonists, used to treat nausea and vomiting;
- Medicines that slow down your central nervous system such as sedatives, sleeping pills, drugs used for surgery (anesthetics), tranquilizers (benzodiazepines), drugs used to treat certain mental conditions (psychotropic drugs), some strong painkillers (opioids), or alcohol;
- Antibiotics to treat infections, e.g. erythromycin, rifampin, linezolid, flucloxacillin;
- Anti-fungals used to treat fungal infections, e.g. ketoconazole;
- Anti-retrovirals used to treat HIV/AIDs, e.g. ritonavir;
- Phenytoin, used to treat seizures;
- Carbamazepine, used to treat epilepsy and some types of pain;
- Quinidine, used to treat heart conditions (antiarrhythmics);
- Warfarin like compounds, used to stop blood clotting (blood thinners);
- Drugs that slow down or reduce the conversion (metabolism) of this drug to its active form (CYP2D6 inhibitors); e.g. fluoxetine, paroxetine, amitriptyline, and bupropion;
- Cimetidine, used to treat heart burn and peptic ulcers.

Taking ULTRACET® with a benzodiazepine (medicine used as

ULTRACET® TABLET

Tramadol hydrochloride / acetaminophen (37.5mg/325mg)

sedatives or treat anxiety) can depress your central nervous system. Inform your doctor if you are currently taking any benzodiazepine.

Taking ULTRACET® at regular doses for a prolonged period or together with flucloxacillin can cause serious metabolic acidosis (a blood and fluid abnormality). Symptoms of metabolic acidosis include: serious breathing difficulties with deep rapid breathing, drowsiness, feeling sick (nausea) and being sick (vomiting).

Seek medical attention immediately if you or the person taking this medication experience(s) symptoms of unusual dizziness or lightheadedness, extreme sleepiness, slowed or difficulty in breathing, or unresponsiveness.

How to use ULTRACET®

- How to use it

ULTRACET® comes in the form of tablet to be taken by mouth. You can take ULTRACET® at any time of the day with or without food. Do not take more than the recommended dose of ULTRACET®. The lowest effective dose should be used for the shortest period of time.

- How much to use

Adults (including elderly) and Children over 16 years:

Take 1 to 2 tablets every 4 to 6 hours as needed for pain relief.

Do not take more than 8 tablets in a 24-hour period.

Do not take ULTRACET® if you have severe liver problems.

Tell your doctor if you have a kidney disorder (a creatinine clearance of less than 30 mL/min). Do not take more than 2 tablets every 12 hours.

If in doubt, ask your doctor or pharmacist for advice.

- When to use it

Always take ULTRACET® exactly as your doctor has told you. You should

check with your doctor if you are not sure.

- How long to use it

Continue taking ULTRACET® for as long as your doctor recommends.

- If you forget to use it

If you miss a dose, take it as soon as you remember. But, if it is almost time for the next dose, do not take the missed dose. Instead, take the next

scheduled dose. Do not try to make up for the missed dose by taking a double dose next time.

- If you use too much (overdose)

Tell your doctor immediately if you have taken too much ULTRACET®, especially if you have any of the following symptoms:

Breathing difficulties with slow or shallow breathing and/or fits (seizures), irritation and discomfort in the stomach and gut, loss of appetite, nausea, vomiting, feeling unwell, unusually pale colour and sweating. Cases of abnormal electrical conduction in the heart have been reported.

Your doctor will take the necessary measures.

Accidental swallowing of ULTRACET® tablets, especially by children, can result in breathing difficulties, with slow or shallow breathing, and/or fits (seizures). Deaths have been reported.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking ULTRACET®.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not take more than your recommended dose.

Discontinuation:

Stopping the ULTRACET® tablets too quickly may cause some side effects, so your doctor may ask you to gradually stop taking your medication over several days. To get the full benefit from your treatment, it is important to take ULTRACET® tablets as prescribed.

- Things to be careful of

Tramadol is not to be used during breastfeeding. Small amounts of tramadol is excreted into breast milk. On a single dose, it is usually not necessary to interrupt breastfeeding. If you have taken ULTRACET® when you are breastfeeding, seek immediate medical attention if you notice your baby has any changes in their breathing (such as weak, difficult or fast breathing).

Serotonin syndrome:

ULTRACET® may cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. If you have some or all of these symptoms: feeling confused, feeling restless, sweating, shaking, shivering, hallucinations, sudden jerks in your muscles or a fast heartbeat, seek medical attention immediately.

Adrenal insufficiency:

Long-term use of ULTRACET® may cause adrenal insufficiency, a potentially life-threatening condition that may present with non-specific symptoms and signs such as nausea, vomiting, decreased appetite, fatigue, weakness, dizziness, and low blood pressure. Seek medical attention if you experience a constellation of these symptoms.

Infertility:

Long-term use of ULTRACET® may cause reduced fertility. It is known whether these effects on fertility are reversible.

ULTRACET[®] TABLET

Tramadol hydrochloride / acetaminophen (37.5mg/325mg)

Opioids can be abused and misused, and you are at risk for opioids addiction, even if you take your dose as prescribed.

Tell your doctor if you:

- Have an allergic reaction (characterized by significant skin rash, itching, hives, difficulty breathing, and/or swollen face) while taking codeine or other opioids.
- Taking other medicines, which contain tramadol or acetaminophen.
- Taking any medicines, because taking some medicines together with ULTRACET[®] may increase your risk of fits (seizures).
- Have asthma, any history of serious breathing problems or if you experience breathing difficulties, with slow or shallow breathing.
- ULTRACET[®] can cause low levels of oxygen in the blood and a problem called sleep apnea (stopping breathing from time to time whilst sleeping). Tell your doctor if you have a history of sleep apnea or if anyone notices you stop breathing from time to time whilst sleeping.
- ULTRACET[®] is not for use in children under 12 years old as it may result in breathing difficulties, with slow or shallow breathing, that could lead to death.
- Taking any drugs that slow down your central nervous system (for example, drugs that make you sleepy, reduce anxiety or decrease awareness). Taking these kinds of drugs with ULTRACET[®] can cause severe drowsiness, decreased awareness and breathing difficulties, with slow or shallow breathing.
- Have recently suffered from a head injury or severe headaches associated with vomiting.
- Have a history of substance dependence or abuse of drugs, alcohol or any opioid painkillers.
- Have a history of mental health problems.
- Want to stop taking ULTRACET[®]. You may become dependent on ULTRACET[®] and may need to reduce the dose of ULTRACET[®] slowly to avoid withdrawal symptoms.
- Stop taking ULTRACET[®] and tell your doctor immediately if you experience skin rash, especially:
 - A red rash covered with small pus-filled bumps that can spread over the body, sometimes with a fever (acute generalized exanthematous pustulosis).
 - A life-threatening rash with blisters and peeling skin over much of the body (toxic epidermal necrolysis).
 - A severe rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome).
- Some patients taking this medicine may experience low levels of sodium in the blood (hyponatremia). Symptoms of low blood sodium may include nausea and vomiting, headaches, feeling confused, feeling very tired, feeling restless, feeling irritable, muscle weakness, spasms or cramps and seizures. Elderly patients and patients taking other medicines that lower sodium in the blood are most at risk for this side effect. If you get any of these symptoms while taking ULTRACET[®], tell your doctor immediately.
- Tell your doctor if you have problems with or have had surgery on your gall bladder or bile duct so you can be monitored for potential development of acute pancreatitis (inflamed pancreas).
- Opioids can cause a high level of the hormone 'prolactin' and low levels of sex hormones in the blood. The effects of these hormone changes may include:
 - In females – unexpected production of breast milk; loss of menstrual periods
 - In males – enlarged breasts; difficulty getting and keeping an erection (impotence)
 - In females or males – decreased sex drive; infertility.

Tell your doctor if you notice any of these symptoms. You may need blood tests, and your doctor may tell you to stop using ULTRACET[®].

Warning

This preparation contains
PARACETAMOL.

Do not take any other
paracetamol containing
medicines at the same time.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Alcohol

You should not take ULTRACET[®] tablets with any alcohol-containing beverages. Also, you should tell your doctor if you drink alcohol regularly, or have a history of alcoholism.

Driving and using machines

ULTRACET[®] may make you feel tired, dizzy or drowsy. If this happens, do not drive or use any machines.

Side effects

Like all medicines, ULTRACET[®] can cause side effects, although not everybody gets them. Tell your doctor if you notice any of the following side effects while using ULTRACET[®]:

Very Common

- Nausea

Common

- Loss of appetite, difficulty sleeping, dizziness, headache, feeling sleepy, vomiting, constipation, diarrhea, indigestion, dry mouth, stomach discomfort, itching, increased sweating, feeling tired, hot flushes, stomach pain, wind, rash

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug

ULTRACET® TABLET

Tramadol hydrochloride / acetaminophen (37.5mg/325mg)

Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting side effects to medicines (ConSERF) or vaccines (AEFI)].

starch, polyethylene glycol, polysorbate 80, powdered cellulose, pregelatinized starch, sodium starch glycolate, and titanium dioxide.

Storage and disposal of ULTRACET®

- Storage

ULTRACET® tablets should be stored at room temperature (15°C to 30°C).

Do not use ULTRACET® tablets after the expiry date, even if it has been stored properly.

Keep this and all medicines in a safe place away from children.

Store this medicine securely, where other people cannot access it. It may harm people who may take this medicine by accident, or intentionally when it has not been prescribed for them.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

- MAL number:

MAL20051462AZ

Manufacturer

Janssen-Cilag S.p.A.
Via C. Janssen (loc. Borgo S. Michele)
04100 Latina (LT), Italy

Product registration holder

Johnson & Johnson Sdn Bhd (3718-D)
Level 8, The Pinnacle,
Persiaran Lagoon, Bandar Sunway,
46150, Petaling Jaya, Selangor, Malaysia

Date of revision

06/08/2025 (MY PI based on CCDS v02May25)

Serial number

NPRA (R1/6) 15112024/207

Product description

- What it looks like

A light-yellow tablet with an off-white core, with debossing as follows: 'J-C' on one side and 'T/P' on the other side.

ULTRACET® tablets are packaged in blisters of standard PVC with aluminium foil backing, polypropylene with aluminium foil backing and polypropylene with polypropylene backing. Each blister contains ten tablets. Pack sizes available are 2, 10 and multiples of 10 up to 60 tablets per pack (however not all pack sizes listed may be marketed).

- Ingredients

- Active ingredients in ULTRACET® are 37.5mg tramadol hydrochloride and 325 mg acetaminophen.

- Inactive ingredients
Carnauba wax, hypromellose, iron oxide, magnesium stearate, maize