

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

What is in this leaflet

1. What Stelara[®] is used for
2. How Stelara[®] works
3. Before you use Stelara[®]
4. How to use Stelara[®]
5. While you are using it
6. Side effects
7. Storage and Disposal of Stelara[®]
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Stelara[®] is used for

Stelara is a prescription medicine that is approved for the treatment of:

- Adults with moderate to severe plaque psoriasis that is chronic (doesn't go away).
- Adults with active psoriatic arthritis, an inflammatory disease of the joints that is usually accompanied by psoriasis.
- Adults with moderately to severely active Crohn's disease or ulcerative colitis.

How Stelara[®] works

Stelara blocks the action of two proteins in your body called interleukin 12 (IL-12) and interleukin 23 (IL-23). IL-12 and IL-23 are made by your body's immune system. In people with psoriasis, psoriatic arthritis or Crohn's disease, IL-12 and IL-23 can cause their immune system to attack normal healthy parts of their body. Stelara can block the IL-12 and IL-23 from causing the immune system to attack the skin, nails, joints or the digestive tract.

Before you use Stelara[®]

- *When you must not use it*

You should not receive Stelara if you:

- are allergic to ustekinumab or any of the other ingredients of this medicine. See the end of this labeling for a complete list of ingredients in Stelara.
- have an active infection which your doctor thinks is important.

- *Before you start to use it*

Talk to your doctor or pharmacist before using Stelara. Your doctor will check how

well you are before each treatment. Make sure you tell your doctor about any illness you have before each treatment. Also tell your doctor if you have recently been near anyone who might have tuberculosis. Your doctor will examine you and do a test for tuberculosis, before you have Stelara. If your doctor thinks you are at risk of tuberculosis, you may be given medicines to treat it.

Before you use Stelara tell your doctor if you:

- ever had an allergic reaction to Stelara. Ask your doctor if you are not sure.
- have ever had any type of cancer – this is because immunosuppressants like Stelara weaken part of the immune system. This may increase the risk of cancer.
- have been treated for psoriasis with other biologic medicines (a medicine produced from a biological source and usually given by injection) – the risk of cancer may be higher.
- have or have had a recent infection.
- have any new or changing lesions within psoriasis areas or on normal skin.
- If you have ever had an allergic reaction to latex or Stelara injection – the container of this medicinal product contains latex rubber, which may cause severe allergic reactions in people who are sensitive to latex. See 'Side effects' for the signs of an allergic reaction.
- are having any other treatment for psoriasis and/or psoriatic arthritis – such as another immunosuppressant or phototherapy (when your body is treated with a type of ultraviolet (UV) light). These treatments may also weaken part of the immune system. Using these therapies together with Stelara has not been studied. However it is possible it may increase the chance of diseases related to a weaker immune system.
- are having or have ever had injections to treat allergies – it is not known if

Stelara may affect these.

- are 65 years of age or over – you may be more likely to get infections.

Some patients have experienced lupus-like reactions including skin lupus or lupus-like syndrome during treatment with ustekinumab. Talk to your doctor right away if you experience a red, raised, scaly rash sometimes with a darker border, in areas of the skin that are exposed to the sun or with joint pains.

Heart attack and strokes

Heart attack and strokes have been observed in a study in patients with psoriasis treated with Stelara. Your doctor will regularly check your risk factors for heart disease and stroke in order to ensure that they are appropriately treated. Seek medical attention right away if you develop chest pain, weakness or abnormal sensation on one side of your body, facial droop, or speech or visual abnormalities.

Children and adolescents

The Stelara pre-filled pen is not recommended for use in children and adolescents under 18 years of age with psoriasis or Crohn's disease because it has not been studied in this age group. The pre-filled syringe or vial should be used instead for children 6 years of age and older and adolescents with psoriasis. The solution for infusion, pre-filled vial or syringe should be used instead for children weighing at least 40 kg with Crohn's disease.

Stelara is not recommended for use in children and adolescents under 18 years of age with psoriatic arthritis, or ulcerative colitis or children with Crohn's disease who weigh less than 40 kg because it has not been studied in this age group.

Pregnancy and lactation

- If you are pregnant, think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine.
- A higher risk of birth defects has not been seen in babies exposed to Stelara

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

in the womb. However, there is limited experience with Stelara in pregnant women. It is therefore preferable to avoid the use of Stelara in pregnancy.

- If you are a woman of childbearing potential, you are advised to avoid becoming pregnant and must use adequate contraception while using Stelara and for at least 15 weeks after the last Stelara treatment.
- Stelara can pass across the placenta to the unborn baby. If you received Stelara during your pregnancy, your baby may have a higher risk for getting an infection.
- It is important that you tell your baby's doctors and other health care professionals if you received Stelara during your pregnancy before the baby receives any vaccine. Live vaccines such as the BCG vaccine (used to prevent tuberculosis) are not recommended for your baby in the first twelve months after birth if you received Stelara during the pregnancy unless your baby's doctor recommends otherwise.
- Ustekinumab may pass into breast milk in very small amounts. Talk to your doctor if you are breast feeding or are planning to breast feed. You and your doctor should decide if you should breast-feed or use Stelara - do not do both.

- Taking other medicines

Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of your medicines and show them to your doctor and pharmacist when you get a new medicine.

Tell your doctor or pharmacist if you have recently had or are going to have a vaccination. Some types of vaccines (live vaccines) should not be given while using Stelara.

If you received Stelara while pregnant,

tell your baby's doctor about your Stelara treatment before the baby receives any vaccine, including live vaccines, such as the BCG vaccine (used to prevent tuberculosis). Live vaccines are not recommended for your baby in the first twelve months after birth if you received Stelara during the pregnancy unless your baby's doctor recommends otherwise.

How to use Stelara[®]

- For treatment of psoriasis and psoriatic arthritis, Stelara is given by injection under the skin.
- For treatment of Crohn's disease or ulcerative colitis, the first dose of Stelara is given by an intravenous infusion, which means that the medicine will be given to you through a needle placed in a vein. This dose is given by your healthcare provider. After that, Stelara is given by injection under the skin.
- Stelara is intended for use under the guidance and supervision of your doctor. If your doctor determines that it is appropriate, you may be able to administer Stelara under the skin to yourself, after proper training in injection technique using the right type of syringe and the amount (volume) to be injected (see the "Instructions for Injecting Stelara under the skin yourself").
- Your doctor will determine the right dose of Stelara for you, the amount for each injection and how often you should receive it. Make sure to discuss with your doctor when you will receive injections and to come in for all your scheduled follow-up appointments.

Instructions for Injecting Stelara under the skin yourself

Your first Stelara may be administered by your healthcare provider. However, your healthcare provider may decide that it is right for you or your caregiver to learn how to inject Stelara under the skin (subcutaneously) yourself. **Before you**

self-inject Stelara, you must be trained by a healthcare professional. If you have not been trained, please contact your healthcare provider to schedule a training session. Call your healthcare provider if you have any questions about giving yourself an injection.

Stelara should not be used:

- after the expiry date on the label;
- if the seal is broken;
- if the liquid is discolored, cloudy or you can see other particulate matter floating in it;
- if you know, or think that it may have been exposed to extreme temperatures (such as accidentally frozen or heated).


Always keep medicine out of the reach of children.

INSTRUCTIONS FOR INJECTING STELARA USING A PRE-FILLED SYRINGE

To reduce the risk of accidental needle sticks to users, each pre-filled syringe is equipped with a needle guard that is automatically activated to cover the needle after complete delivery of the syringe content.

Do not shake Stelara at any time. Prolonged vigorous shaking may damage the product. If the product has been shaken vigorously, don't use it.

STEP 1: PREPARING FOR PRE-FILLED SYRINGE USE



Take the Syringe out of the Refrigerator

Check the package to make sure that it is the right dose, **either 45 mg or 90 mg as prescribed by your healthcare provider.**

Check Expiration Date

Open the box and remove the pre-filled syringe. Check the expiration date on the

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

pre-filled syringe and the label of the box. If the expiration date has passed, don't use it.

Assemble Additional Supplies

Assemble the additional supplies you will need for your injection. These include an antiseptic wipe, a cotton ball or gauze, and a sharps container for syringe disposal.

Check Solution in Syringe

Hold the pre-filled syringe with the covered needle pointing upward. Make sure the syringe is not damaged. Look at the solution or liquid in the syringe to make sure that it is clear to slightly opalescent and colorless to slightly yellow. **DO NOT** use if it is frozen, discolored, cloudy or contains large particles and contact your healthcare provider for assistance.


DO NOT remove the needle cover from the pre-filled syringe.

DO NOT pull back on the plunger head at any time.

STEP 2: CHOOSING AND PREPARING THE INJECTION SITE

Choose the injection site*

Good sites are the top of the thigh and around the tummy (abdomen) but about 2 inches away from the belly button (navel). Avoid, if possible, skin involved with psoriasis. If your caregiver is giving you the injection, they may use the upper arms as well.



*Areas in gray are recommended injection sites.


Prepare the Injection Site

Thoroughly wash your hands with soap and warm water. Wipe the injection site with an antiseptic wipe. **DO NOT** touch this area again before giving the injection.

STEP 3: INJECTING THE MEDICATION

Remove the Needle Cover


When you are ready to inject, pick up the pre-filled syringe, hold the body of the syringe with one hand and pull the needle cover straight off. Throw the needle cover into the trash. You may notice a small air bubble in the pre-filled syringe. You do not need to remove the air bubble. You may also see a drop of liquid at the end of the needle – this is normal. Do not touch the needle or allow it to touch any surface.



Note: The needle cover should **NOT** be removed until you are ready to inject the dose. Do not use syringe if it is dropped without the needle cover in place. If you drop the syringe without the needle cover in place, please contact your healthcare provider for assistance.

Inject the Medication

Gently pinch the cleaned skin between your thumb and index finger.




Don't squeeze it.

Push the syringe needle into the pinched skin.

Push the plunger with your thumb as far as it will go to inject all of the liquid. Push it slowly and evenly, keeping the skin pinched.

When the plunger meets the end of the syringe barrel, and all of the medication has been injected, release the pinched skin and gently remove the needle. Following

complete injection, the needle guard will automatically extend over the needle and lock as you take your hand off the plunger.



STEP 4: AFTER THE INJECTION

Dispose of the Empty Syringe

Immediately dispose of the empty syringe into the sharps container. For your safety and health and for the safety of others, needles and syringes must **NEVER** be re-used. Dispose of sharps container according to your local regulations.




Use a Cotton Ball or Gauze

There may be a small amount of blood or liquid at the injection site, which is normal. You can press a cotton ball or gauze over the injection site and hold for 10 seconds. Do not rub the injection site. You may cover the injection site with a small adhesive bandage, if necessary.

INSTRUCTIONS FOR INJECTING STELARA USING A ONE-PRESS PRE-FILLED PEN

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen



Important

Stelara comes in a single-dose One-press pre-filled pen containing one 45 mg dose or one 90 mg dose.

During injection, push handle all the way down until purple body is not visible to inject the full dose. DO NOT LIFT ONE-PRESS PRE-FILLED PEN during injection! If you do, the One-Press pre-filled pen will lock and you will not get the full dose.

If your healthcare provider (doctor, nurse, or pharmacist) decides that you or a caregiver may be able to give your injections of Stelara at home, you should receive training on the right way to prepare and inject Stelara using the One-Press One-Press pre-filled pen. **Do not try to inject yourself until you have been trained by your healthcare provider.**

Each One-Press pre-filled pen can only be used one time. Throw it away (see Step 3) after use even if there is medicine left in it.

Do not reuse your One-Press pre-filled pen.

If you cannot give yourself the injection:

- ask your doctor or nurse to help you, or
- ask someone who has been trained by a doctor or nurse to give your injections.

To reduce the risk of accidental needle sticks, each One-Press pre-filled pen has a needle guard that automatically covers the needle and locks after you have given your injection and the injector is lifted. Do not lift the One-Press pre-filled pen during the injection until the injection is complete.


The needle cover on the syringe in the STELARA pre-filled syringe / inside the bottom cap of the One-Press pre-filled pen contains latex. **Do not handle the needle cover if you are allergic to latex.**

One-Press/pre-filled pen injector parts



Gather the following supplies.
Provided in the carton:
• One-Press/pre-filled pen injector
Not provided in the carton:
• Alcohol swabs
• Cotton balls or gauze pads
• Adhesive bandages
• Sharps container (See Step 3)

STEP 1: PREPARING TO INJECT STELARA



Gather carton(s)


If refrigerated, remove your One-Press pre-filled pen carton(s) from the refrigerator and place on a flat surface.

Leave at room temperature for at least 30 minutes before use.

Do not warm any other way.

If your dose is 45 mg, you will receive one 45 mg One-Press pre-filled pen. If your dose is 90 mg, you will receive one 90 mg or two 45 mg One-Press pre-filled pens. If you receive two 45 mg One-Press pre-filled pens, follow Steps 1-3 for both injections.

Choose a different injection site for the second injection.



Check the expiration date ('EXP') and seals on the carton(s)

Do not use your One-Press pre-filled pen if the seals on the carton are broken or if the expiration date has passed.

Do not use your One-Press pre-filled pen if it has been kept at room temperature for longer than 30 days or if it has been stored above 30°C. Call your healthcare provider or pharmacist for a new One-Press pre-filled pen.



STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

Choose injection site

Select from the following areas for your injection:


- Front of thighs
- Lower stomach area (lower abdomen), except for a 2-inch area right around your navel (belly-button)

If someone else is giving you the injection, they may also use:

- Back of upper arms

Do not inject into skin that is tender, bruised, red or hard.

Use a different injection site for each injection.



Wash hands

Wash your hands well with soap and warm water.

Clean injection site

Wipe your chosen injection site with an alcohol swab and allow it to dry.

Do not touch, fan, or blow on the injection site after you have cleaned it.




Inspect liquid in window

Choose a well lit, clean, flat work surface. Take your One-Press pre-filled pen out of the carton and check for damage.

Check the liquid in the viewing window. It should be **clear and colorless** to **slightly yellow** and may contain **tiny white or clear particles** and **one or more air bubbles**. This is normal.

Do not inject if the liquid is frozen, cloudy, discolored, or has large particles. Call your healthcare provider or pharmacist for a new One-Press pre-filled pen.

STEP 2: INJECTING STELARA




Pull off bottom cap

Keep hands away from the needle guard after the cap is removed. It is normal to see a few drops of liquid.

Inject STELARA within 5 minutes of removing the cap.

Do not put the cap back on. This could damage the needle.

Do not use a One-Press pre-filled pen if it is dropped after removing the cap. Call your healthcare provider or pharmacist for a new One-Press pre-filled pen.



Place straight on skin. Push handle all the way down until purple body is not visible.


DO NOT LIFT ONE-PRESS during injection!

If you do, the needle guard will lock, showing a yellow band, and you will not get the full dose.

You may hear a click when the injection begins. Keep pushing.

If you feel resistance, keep pushing. This is normal.

The medication injects as you push. Do this at a speed that is comfortable for you.




Confirm your injection is complete

Your injection is complete when:

- **The purple body is not visible.**
- You cannot press the handle down anymore.
- You may hear a click.

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen



Lift straight up

The yellow band indicates that the needle guard is locked into place.

STEP 3: AFTER YOUR INJECTION




Dispose of your One-Press pre-filled pen

Put your used pen in a sharps disposal container right away after use.

Do not throw away (dispose of) your One-Press pre-filled pens in your household trash.

Do not recycle your used sharps disposal container.

For more information, see “Disposal”.



Check injection site

There may be a small amount of blood or liquid at the injection site. This is normal. Press a cotton ball or gauze pad to the injection site until any bleeding stops.

Do not rub the injection site.

If needed, cover the injection site with a bandage.

If you receive two 45 mg One-Press pre-filled pens for a 90 mg dose, repeat Steps 1-3 with your second One-Press pre-filled pen. **Choose a different injection site for the second injection.**

This information sheet summarizes the most important information about Stelara. You can ask your healthcare provider for information about Stelara that is written for healthcare professionals.

- How much to use

Your doctor will determine the right dose of Stelara for you, the amount for each injection and how often you should receive it. Make sure to discuss with your doctor when you will receive injections and to come in for all your scheduled follow-up appointments.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Stelara for as long as your doctor recommends.

- If you forget to use it

Make the next injection as soon as you remember. Do not double up the injections.

If you have missed more than one dose, or are not sure what to do, check with your doctor or pharmacist.

If you have trouble remembering when to use your medicine, ask your pharmacist for some hints.

- If you use too much (overdose)

Immediately telephone your doctor, or go to Accident and Emergency at your nearest hospital.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

Always follow your doctor's instructions carefully.

- Tell your doctor about your medical conditions before each treatment (see *Before you use Stelara*).
- Tell your doctor if you become pregnant while using Stelara.
- If you are about to start taking a new medicine, tell your doctor and pharmacist that you are using Stelara.

- Things you must not do

- You should not receive a live vaccine while taking Stelara.
- If you used Stelara while pregnant, tell your baby's doctor about your Stelara use before the baby receives any vaccine, including live vaccines, such as the BCG vaccine (used to prevent tuberculosis), rotavirus vaccine, or any other live vaccines.
- Medicines are sometimes prescribed for purposes that are not mentioned in patient information sheets. Do not use Stelara for a condition for which it was not prescribed.
- Do not give this medicine to anyone else, even if their symptoms seem similar to yours.

- Things to be careful of

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

Driving and using machines

Stelara has no or negligible influence on the ability to drive and use machines.

Side effects

Like all medicines, Stelara can cause side effects, although not everybody gets them.

Serious side effects

Some patients may have serious side effects that may need urgent treatment.

Allergic reactions – these may need urgent treatment. Tell your doctor or get emergency medical help straight away if you notice any of the following signs.

- Serious allergic reactions ('anaphylaxis') are rare in people taking Stelara (may affect up to 1 in 1,000 people). Signs include:
 - difficulty breathing or swallowing
 - low blood pressure, which can cause dizziness or light headedness
 - swelling of the face, lips, mouth or throat.
- Common signs of an allergic reaction include skin rash and hives (these may affect up to 1 in 100 people).

In rare cases, allergic lung reactions and lung inflammation have been reported in patients who receive ustekinumab. Tell your doctor right away if you develop symptoms such as cough, shortness of breath, and fever.

If you have a serious allergic reaction, your doctor may decide that you should not use Stelara again.

Infections – these may need urgent treatment. Tell your doctor straight away if you notice any of the following signs.

- Infections of the nose or throat and common cold are common (may affect up to 1 in 10 people)
- Infections of the chest are uncommon (may affect up to 1 in 100 people)
- Inflammation of tissue under the skin ('cellulitis') is uncommon (may affect up to 1 in 100 people)

- Shingles (a type of painful rash with blisters) are uncommon (may affect up to 1 in 100 people)

Stelara may make you less able to fight infections. Some infections could become serious and may include infections caused by viruses, fungi, bacteria (including tuberculosis), or parasites, including infections that mainly occur in people with a weakened immune system (opportunistic infections). Opportunistic infections of the brain (encephalitis, meningitis), lungs, and eye have been reported in patients receiving treatment with ustekinumab.

You must look out for signs of infection while you are using Stelara. These include:

- fever, flu like symptoms, night sweats, weight loss
- feeling tired or short of breath; cough which will not go away
- warm, red and painful skin, or a painful skin rash with blisters
- burning when passing water
- diarrhoea
- visual disturbance or vision loss
- headache, neck stiffness, light sensitivity, nausea or confusion.

Tell your doctor straight away if you notice any of these signs of infection. These may be signs of infections such as chest infections, or skin infections or shingles or opportunistic infections that could have serious complications. Tell your doctor if you have any kind of infection that will not go away or keeps coming back. Your doctor may decide that you should not use Stelara until the infection goes away. Also tell your doctor if you have any open cuts or sores as they might get infected.

Shedding of skin – increase in redness and shedding of skin over a larger area of the body may be symptoms of erythrodermic psoriasis or exfoliative dermatitis, which are serious skin conditions. You should tell your doctor straight away if you notice any of these signs.

Other side effects

Common side effects (may affect up to 1 in 10 people):

- Diarrhea
- Nausea
- Vomiting
- Feeling tired
- Feeling dizzy
- Headache
- Itching ('pruritus')
- Back, muscle or joint pain
- Sore throat
- Redness and pain where the injection is given
- Sinus infection

Uncommon side effects (may affect up to 1 in 100 people):

- Tooth infections
- Vaginal yeast infection
- Depression
- Blocked or stuffy nose
- Bleeding, bruising, hardness, swelling and itching where the injection is given
- Feeling weak
- Drooping eyelid and sagging muscles on one side of the face ('facial palsy' or 'Bell's palsy'), which is usually temporary
- A change in psoriasis with redness and new tiny, yellow or white skin blisters, sometimes accompanied by fever (pustular psoriasis)
- Peeling of the skin (skin exfoliation)
- Acne

Rare side effects (may affect up to 1 in 1,000 people):

- Redness and shedding of skin over a larger area of the body, which may be itchy or painful (exfoliative dermatitis). Similar symptoms sometimes develop as a natural change in the type of psoriasis symptoms (erythrodermic psoriasis)
- Inflammation of small blood vessels, which can lead to a skin rash with small red or purple bumps, fever or

STELARA[®] SOLUTION FOR INJECTION

Ustekinumab (45mg/0.5ml, 90mg/1ml) in Prefilled Syringe and One-Press Solution for Injection in Pre-filled Pen

joint pain (vasculitis)

Very rare side effects (may affect up to 1 in 10,000 people):

- Blistering of the skin that may be red, itchy, and painful (Bullous pemphigoid).
- Skin lupus or lupus-like syndrome (red, raised scaly rash on areas of the skin exposed to the sun possibly with joint pains).

These are not all of the side effects with Stelara. Tell your doctor about any side effect that bothers you or does not go away. Ask your doctor or pharmacist for more information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.moh.gov.my [Consumers → Reporting side effects to medicines (ConSERF) or vaccines (AEFI)].

Storage and Disposal of Stelara[®]

- Storage

If you are using Stelara at home, it is important to store the product in your refrigerator although not in the freezer compartment. Stelara should not be frozen. Keep the product in the original carton to protect from light until the time of use.

If needed, individual Stelara pre-filled syringes and One-Press pre-filled pens may also be stored at room temperature up to 30°C for a maximum single period of up to 30 days in the original carton with protection from light. Record the date when the pre-filled syringe or One-Press pre-filled pen is first removed from the refrigerator and the new expiry date on the carton in the space provided. The new expiry date must not exceed the original expiry date printed on the carton. Once a syringe or One-Press pre-filled pen has been stored at room temperature, it should not be returned to the refrigerator. Discard the syringe or One-Press pre-filled pen if

not used within 30 days at room temperature storage.

- Disposal

Used syringes should be placed in a puncture-resistant container, like a sharps container. Dispose of your sharps container according to your local regulations. Empty vials, antiseptic wipes, and other supplies can be placed in your regular trash. The syringe, needle and vial must never be re-used. Discard any unused portion of Stelara in accordance with local requirements.

Product Description

- What it looks like

Stelara is a colorless to light yellow product and may contain a few small translucent or white particles of protein. This appearance is not unusual for solutions containing protein.

- Ingredients

- Active ingredients
The active ingredient is ustekinumab.

- Inactive ingredients

The inactive ingredients include: L-histidine, L-histidine monohydrochloride monohydrate polysorbate 80, sucrose and water for injection. No preservatives are present.

- MAL number

Stelara 45mg/0.5mL Solution for Injection in Prefilled Syringe: MAL20112139AZ

Stelara 90mg/1.0mL Solution for Injection in Prefilled Syringe: MAL20112140AZ

Stelara One-Press 45mg/0.5mL Solution for Injection in Pre-filled Pen: MALXXXXXXXX

Stelara One-Press 90mg/1.0mL Solution for Injection in Pre-filled Pen: MALXXXXXXXX

Manufacturer

Cilag AG,
Hochstrasse 201, 8200, Schaffhausen,
Switzerland.

Product Registration Holder

Johnson & Johnson Sdn Bhd (3718-D)
Level 8, The Pinnacle,
Persiaran Lagoon, Bandar Sunway,
46150, Petaling Jaya, Selangor, Malaysia

Date of Revision

09/03/2026 (MY PI based on CCDS
v04Mar2024 & EU SmPC v24Mar2025)

Serial Number

NPRA(R3/01) 23092024/0320