

VALOSIN[®] FG 50

MEDICATED PREMIX: CONTAINS 42.5 g TYLVALOSIN (AS TARTRATE) PER kg

Active substance: 42.5 g (as Tylvalosin tartrate)

INDICATIONS FOR USE:

For the control and treatment of Enzootic Pneumonia, Swine Dysentery and ileitis (Porcine Proliferative Enteropathy).

NET WEIGHT: 5 kg

DOSAGE AND ADMINISTRATION: For oral administration in feed only. To obtain therapeutic levels of Tylvalosin, mix **VALOSIN[®] FG 50** with feed at the following rates:

Indication	Dose of active ingredient	Duration of Treatment	Quantity of VALOSIN [®] FG 50 per ton of feed
Treatment and prevention of swine enzootic pneumonia	2.125 mg / kg bw / day	7 days	1 kg / tonne *
Treatment of PPE (Ileitis)	4.25 mg / kg bw / day	10 days	2 kg / tonne *
Treatment and prevention of swine dysentery	4.25 mg / kg bw / day	10 days	2 kg / tonne *

* **Important:** these inclusion rates assumes a pig eats the equivalent of 5% of their body weight per day.

VALOSIN[®] FG 50 It is essential to mix well.

SIGNS AND SYMPTOMS OF OVERDOSE AND TREATMENT: No signs of intolerance have been observed in growing pigs at up to 10 times the recommended dosage.

WARNINGS AND PRECAUTIONS: Operators should avoid contact with the skin and eyes and should use dust masks when preparing medicated feed.

Jauhi dari kanak-kanak. Keep out the reach of children. For animal treatment only.

Usage During Pregnancy and lactation: The safety of the use of **VALOSIN[®] FG 50** during pregnancy and lactation has not been established. Use in accordance with the risk/benefit assessment by the responsible veterinarian.

No Known Contra-Indications or Incompatibilities: Undesirable side effects-adverse reations-none known.

Withdrawal period: Meat and offal: 2 days.

Storage: Do not store above 25°C. Keep the container tightly closed. Store in the original container.

Disposal of Containers: Dispose of in accordance with the laws of the Local Authorities.

Shelf life after incorporation into meal or pelleted feed: 1 month.

Pharmacodynamics Pharmacokinetics: Tylvalosin tartrate is a macrolide antibiotic that has antibacterial activity against Gram positive, some Gram negative organisms and mycoplasma.

It acts by inhibiting protein synthesis in the bacterial cell. Tylvalosin tartrate is rapidly absorbed after oral administration of **VALOSIN[®] FG 50**.

After administration of the recommended dose lung concentrations of 0.06-0.066 pg/ml were found at 2 and 12 hours post-treatment. The parent compound is widely distributed in the tissues with the highest concentrations found in the lungs, bile, intestinal mucosa, spleen, kidney and liver.

Reg. No.: MAL 10100040HA For animal treatment only - to be supplied only on veterinary prescription.

MANUFACTURER: Cod Beck Blenders Limited, Cod Beck Estate, Dalton Lane, Dalton, Thirsk, North Yorkshire, YO7 3HR, United Kingdom.

REGISTRATION HOLDER: ECO Animal Health Malaysia Sdn Bhd. Menara Maxis, 36th Floor, Kuala Lumpur City Centre, Kuala Lumpur, Malaysia 5088.



ECO ANIMAL HEALTH LIMITED.

78 Coombe Road,
New Malden, Surrey,
KT3 4QS, U.K.

Tel: + 44 (0)20 8447 8899
Fax: + 44 (0)20 8447 9292

Email: enquiries@ecoanimalhealth.com

Date of revision: 06/2019

5 kg



Batch:

Manufacture Date:

Expiry Date:

CONTROLLED MEDICINE

**NON VARNISH
AREA**

V24-1

DIE SIZE: 209.55 x 296.33mm

OUTER MAGENTA KEYLINE SHOWS CUTTING DIE ONLY & DOES NOT PRINT

THIS COLOUR PROOF IS NOT ACCURATE FOR COLOUR