

PACKAGE INSERT

1 NAME OF THE MEDICINAL PRODUCT

Salofalk® 500mg
gastro-resistant tablets

Mesalazine

2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each gastro-resistant tablet contains 500 mg mesalazine.

Excipients with known effect: sodium carbonate and croscarmellose sodium

For a full list of excipients, see section 6.1.

3 PHARMACEUTICAL FORM

Gastro-resistant tablets

Appearance: butter-yellow to ochre enteric-coated oblong formed tablets, lustreless with smooth surface, no cracking perceptible.

4 CLINICAL PARTICULARS

4.1 Therapeutic indications

- For the treatment of acute episodes and the maintenance of remission of ulcerative colitis.
- For the treatment of acute episodes of Crohn's disease.

4.2 Posology and method of administration

Posology

Adults and elderly
Depending upon the clinical requirements in individual cases, the following daily doses are recommended:

	Crohn's disease	Ulcerative colitis	
	Acute episode	Acute episode	Prevention of recurrence/long-term treatment
Mesalazine (active substance)	1.5 g - 4.5 g	1.5 g - 3.0 g	1.5 g
Salofalk 500mg tablets	1 tablet three times per day to 3 tablets three times per day	1 tablet three times per day to 2 tablets three times per day	1 tablet three times per day

Children
There is only limited documentation for an effect in children (age 6-18 years).



Children 6 years of age and older

Active disease: To be determined individually, starting with 30-50 mg/kg body weight/day in divided doses. Maximum dose: 75 mg/kg body weight/day. The total dose should not exceed the maximum adult dose.

Maintenance treatment (ulcerative colitis): To be determined individually, starting with 15-30 mg/kg body weight/day in divided doses. The total dose should not exceed the recommended adult dose.

It is generally recommended that half the adult dose may be given to children up to a body weight of 40 kg; and the normal adult dose to those above 40 kg.

Duration of treatment

The treatment of acute episodes of ulcerative colitis usually lasts 8 weeks. The duration of use is determined by the physician.

For maintenance of remission in ulcerative colitis, the dose can usually be reduced to 1.5 g mesalazine/day (adults and adolescents with a body weight over 40 kg) and 0.75 g mesalazine/day (children/adolescents).

Method of administration

Salofalk 500mg tablets should be taken in the morning, at midday and in the evening, 1 hour before meals. They should be swallowed whole, not chewed, and taken with plenty of fluid.

Treatment with Salofalk 500mg tablets should be administered regularly and consistently, both in the acute inflammatory stage and during maintenance therapy in order to achieve the desired therapeutic effect.

4.3 Contraindications

- Salofalk 500mg tablets are contraindicated in patients with
- known hypersensitivity to the active substance, salicylates or to any of the excipients listed in section 6.1
 - severe impairment of hepatic or renal function.

4.4 Special warnings and precautions for use

Blood tests (differential blood count; liver function parameters such as ALT or AST; serum creatinine) and urinary status (dip sticks) should be determined prior to and during treatment, at the discretion of the treating physician. As a guideline, follow-up tests are recommended 14 days after commencement of treatment, then a further two to three tests at intervals of 4 weeks.

If the findings are normal, follow-up tests should be carried out every 3 months. If additional symptoms occur, these tests should be performed immediately.

Caution is recommended in patients with impaired hepatic function.

Mesalazine should not be used in patients with impaired renal function. Mesalazine-induced renal toxicity should be considered, if renal function deteriorates during treatment. If this is the case, Salofalk 500mg tablets should be discontinued immediately.

Cases of nephrolithiasis have been reported with the use of mesalazine including stones with a 100% mesalazine content. It is recommended to ensure adequate fluid intake during treatment.

Mesalazine may produce red-brown urine discoloration after contact with sodium hypochlorite bleach (e.g. in toilets cleaned with sodium hypochlorite contained in certain bleaches).

Serious blood dyscrasias have been reported very rarely with mesalazine. Hematological investigations should be performed if patients suffer from unexplained haemorrhages,

bruises, purpura, anaemia, fever or pharyngolaryngeal pain. Salofalk 500mg tablets should be discontinued in case of suspected or confirmed blood dyscrasia.

Cardiac hypersensitivity reactions (myocarditis, and pericarditis) induced by mesalazine have been rarely reported. Salofalk 500mg tablets should then be discontinued immediately.

Patients with pulmonary disease, in particular asthma, should be very carefully monitored during a course of treatment with mesalazine.

Severe cutaneous adverse reactions

Severe cutaneous adverse reactions (SCARs), including drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported in association with mesalazine treatment. Mesalazine should be discontinued, at the first appearance of signs and symptoms of severe skin reactions, such as skin rash, mucosal lesions, or any other sign of hypersensitivity.

Idiopathic intracranial hypertension

Idiopathic intracranial hypertension (pseudotumor cerebri) has been reported in patients receiving mesalazine. Patients should be warned for signs and symptoms of idiopathic intracranial hypertension, including severe or recurrent headache, visual disturbances or tinnitus. If idiopathic intracranial hypertension occurs, discontinuation of mesalazine should be considered.

Patients with a history of adverse drug reactions to preparations containing sulphasalazine should be kept under close medical surveillance on commencement of a course of treatment with mesalazine. Should Salofalk 500mg tablets cause acute intolerance reactions such as abdominal cramps, acute abdominal pain, fever, severe headache and rash, therapy should be discontinued immediately.

Photosensitivity

More severe reactions are reported in patients with pre-existing skin conditions such as atopic dermatitis and atopic eczema.

Note

In rare cases, in patients who have undergone bowel resection/bowel surgery in the ileocecal region with removal of the ileocecal valve, it has been observed that Salofalk 500 mg tablets were excreted undissolved in the stool, due to an excessively rapid intestinal passage.

This medicinal product contains 49 mg sodium per tablet, equivalent to 2.5 % of the WHO recommended maximum daily intake of 2 g sodium for an adult. The maximum daily dose of this product is equivalent to 22 % of the WHO recommended maximum daily intake for sodium. Salofalk 500 mg tablets are considered high in sodium. This should be particularly taken into account for those on a low salt diet.

4.5 Interaction with other medicinal products and other forms of interaction

Specific interaction studies have not been performed.

In patients who are concomitantly treated with azathioprine, 6-mercaptopurine or thioguanine, a possible increase in the myelosuppressive effects of azathioprine, 6-mercaptopurine or thioguanine should be taken into account.

There is weak evidence that mesalazine might decrease the anticoagulant effect of warfarin.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no adequate data on the use of mesalazine in pregnant women. However, data on a limited number of exposed pregnancies indicate no adverse effect of mesala-

zine on pregnancy or on the health of the fetus/newborn child. To date no other relevant epidemiologic data are available. In one single case after long-term use of a high dose mesalazine (2-4 g/day, orally) during pregnancy, renal failure in a neonate was reported.

Animal studies on oral mesalazine do not indicate direct or indirect harmful effects with respect to pregnancy, embryonic/ fetal development, parturition or postnatal development. Salofalk 500mg tablets should only be used during pregnancy if the potential benefit outweighs the possible risk.

Breastfeeding

N-acetyl-5-aminosalicylic acid and to a lesser degree mesalazine are excreted in breast milk. Only limited experience during lactation in women is available to date. Hypersensitivity reactions such as diarrhoea in the infant cannot be excluded. Therefore, Salofalk 500mg tablets should only be used during breast-feeding if the potential benefit outweighs the possible risk. If the infant develops diarrhoea, the breast-feeding should be discontinued.

4.7 Effects on ability to drive and use machines

Mesalazine has no or negligible effects on the ability to drive and use machines.

4.8 Undesirable effects

The following undesirable effects have been observed after administration of mesalazine (see table at the end of this package insert)

4.9 Overdose

There are rare data on overdosage (e.g. intended suicide with high oral doses of mesalazine), which do not indicate renal or hepatic toxicity. There is no specific antidote and treatment is symptomatic and supportive.

5 PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Intestinal anti-inflammatory agents, Aminosalicilyc acid and similar agents
ATC code: A07EC02

Mechanism of action

The mechanism of the anti-inflammatory action is unknown. The results of in-vitro studies indicate that inhibition of lipoxxygenase may play a role. Effects on prostaglandin concentrations in the intestinal mucosa have also been demonstrated. Mesalazine (5-Aminosalicilyc acid / 5-ASA) may also function as a radical scavenger of reactive oxygen compounds.

Pharmacodynamic effects

Mesalazine, orally administered, acts predominantly locally at the gut mucosa and in the submucous tissue from the luminal side of the intestine. It is important, therefore, that mesalazine is available at the regions of inflammation. Systemic bioavailability / plasma concentrations of mesalazine therefore are of no relevance for therapeutic efficacy, but rather a factor for safety. In order to fulfil these criteria, Salofalk 500mg tablets are coated with Eudragit L; they are thus gastro-resistant and release of mesalazine is pH-dependent.

5.2 Pharmacokinetic properties

General considerations of mesalazine

Absorption

Mesalazine absorption is highest in proximal gut regions and lowest in distal gut areas.

Biotransformation

Mesalazine is metabolised both pre-systemically by the intestinal mucosa and the liver to the pharmacologically inactive N-acetyl-5-aminosalicylic acid (N-Ac-5-ASA). The acetylation seems to be independent of the acetylator phenotype of the patient. Some acetylation also occurs through the action of colonic bacteria. Protein binding of mesalazine and N-Ac-5-ASA is 43% and 78%, respectively.

Elimination

Mesalazine and its metabolite N-Ac-5-ASA are eliminated via the faeces (major part), renally (varies between 20 and 50%, dependent on kind of application, pharmaceutical preparation and route of mesalazine release, respectively), and biliary (minor part). Renal excretion predominantly occurs as N-Ac-5-ASA. About 1% of total orally administered mesalazine dose is excreted into the breast milk mainly as N-Ac-5-ASA.

Salofalk 500mg tablets specific

Distribution

A combined pharmacoscintigraphic/pharmacokinetic study showed that Salofalk 500mg tablets reach the ileocecal region after approximately 3-4 hours in fasting subjects and reach the ascending colon within approximately 4–5 hours. The total transit time in the colon is approximately 17 hours.

Absorption

Release of mesalazine from Salofalk 500mg tablets, begins after a lag-phase of approximately 3–4 hours. Peak plasma concentrations are reached after approximately 5 hours (ileocecal region) and, at 3 x 500 mg mesalazine/ day under steady-state conditions, are 3.0 ± 1.6 µg/ml for mesalazine and 3.4 ± 1.6 µg/ml for the metabolite, N-Ac-5-ASA.

Elimination

The total renal elimination rate for mesalazine and N-Ac-5-ASA over 24 hours during multiple intake (3 x 1 Salofalk 500mg tablets, for 2 days; 1 tablet on the third day=examination day) was approximately 60%. The non-metabolised mesalazine fraction after oral administration was approximately 10%.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, genotoxicity, carcinogenicity (rat) or toxicity to reproduction.

Kidney toxicity (renal papillary necrosis and epithelial damage in the proximal tubule (pars convoluta) or the whole nephron) has been seen in repeat-dose toxicity studies with high oral doses of mesalazine. The clinical relevance of this finding is unknown.

6 PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Basic butylated methacrylate copolymer (Ph. Eur.) (rel. molar mass: approx. 150000 (e.g. Eudragit E)), Calcium stearate (Ph. Eur.) [herbal], Croscarmellose sodium, Iron oxide yellow (E 172), Glycine, Silica, colloidal anhydrous, Hypromellose, Macrogol 6000, Methacrylic acid methyl methacrylate copolymer (1:1) (Ph. Eur.) (rel. molar mass: approx. 135000 (e.g. Eudragit L)), Cellulose, microcrystalline, Sodium carbonate, Povidone K25, Talc, Titanium dioxide (E 171)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Store below 30 °C.

6.5 Nature and contents of container

Container
Blister: PVC/PE/PVdC (orange-transparent) /aluminium blister foil

Package sizes

Blister packs with 100 (10x10's) Salofalk 500mg gastro-resistant tablets.

6.6 Special precautions for disposal and other handling

No special requirements.

7 MANUFACTURER

Losan Pharma GmbH
Otto-Hahn-Straße 13
79395 Neuenburg
Germany

For

DR. FALK PHARMA GmbH



Leinenweberstr. 5
79108 Freiburg im Breisgau
Germany

8 PRODUCT REGISTRATION HOLDER

DCH Auriga (Malaysia) Sdn Bhd (719-V),
Lot 6, Persiaran Perusahaan, Seksyen 23,
40300 Shah Alam,
Selangor Darul Ehsan, Malaysia.

9 PRODUCT REGISTRATION NUMBER

MAL19987336ACSZ

10 DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 24.09.1998
Date of last renewal: 17.10.2025

11 DATE OF REVISION OF THE TEXT

July 2025

12 GENERAL CLASSIFICATION FOR SUPPLY

Medicinal product subject to medical prescription

Undesirable effects

<i>System Organ Class</i>	<i>Frequency according to MedDRA convention</i>				
	<i>Common</i> (≥ 1/100 to < 1/10)	<i>Uncommon</i> (≥ 1/1,000 to < 1/100)	<i>Rare</i> (≥ 1/10,000 to < 1/1,000)	<i>Very rare</i> (< 1/ 10,000)	<i>Not known (cannot be estimated from the available data)</i>
Blood and lymphatic system disorders				Altered blood counts (aplastic anaemia, agranulocytosis, pancytopenia, neutropenia, leukopenia, thrombocytopenia)	
Immune system disorders				Hypersensitivity reactions such as allergic exanthema, drug fever, lupus erythematosus syndrome, pancolitis	
Nervous system disorders	Headache		Dizziness	Peripheral neuropathy	Idiopathic intracranial hypertension (see section 4.4)
Cardiac disorders			Myocarditis, pericarditis		
Respiratory, thoracic and mediastinal disorders				Allergic and fibrotic lung reactions (including dyspnoea, cough, bronchospasm, alveolitis, pulmonary eosinophilia, lung infiltration, pneumonitis)	
Gastrointestinal disorders		Abdominal pain, diarrhoea, dyspepsia, flatulence, nausea, vomiting, acute pancreatitis			
Hepatobiliary disorders			Cholestatic hepatitis	Hepatitis	
Skin and subcutaneous tissue disorders	Rash, pruritus		Photosensitivity	Alopecia	Drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN)
Musculoskeletal and connective tissue disorders			Arthralgia	Myalgia	
Renal and urinary disorders				Impairment of renal function including acute and chronic interstitial nephritis and renal insufficiency	Nephrolithiasis*
Reproductive system and breast disorders				Oligospermia (reversible)	
General disorders			Asthenia, fatigue		
Investigations		Changes in liver function parameters (increase in transaminases and parameters of cholestasis), changes in pancreatic enzymes (lipase and amylase increased), eosinophil count increased			

* See section 4.4 for further information.

Severe cutaneous adverse reactions (SCARs), including drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), have been reported in association with mesalazine treatment (see section 4.4).

1251130

MY/07.25