

CAP SEEN SHEI MINYAK ANGIN ZHUI FONG YEW

Methyl Salicylate 57%w/v, Menthol 8%w/v

What is in this leaflet

1. What Zhui Fong Yew is used for
2. How Zhui Fong Yew works
3. Before you use Zhui Fong Yew
4. How to use Zhui Fong Yew
5. While you are using it
6. Side effects
7. Storage and Disposal of Zhui Fong Yew
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Zhui Fong Yew is used for

It is used to relieve muscle and joint pain related to rheumatic conditions.

How Zhui Fong Yew works

It contains menthol and methyl salicylate. Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action. Methyl salicylate is used topically as a counter-irritant. Upon application, it is absorbed through the skin and is applied for the relief of pain in rheumatic conditions and painful muscle or joints.

Before you use Zhui Fong Yew

- When you must not use it

Do not use it if you are allergic to any ingredients of this Zhui Fong Yew.

For external use only. Not to be taken.

- Before you start to use it

Please follow the advice of the doctor or pharmacist.

- Taking other medicines

Please inform the doctor or pharmacist if taking other medicines.

For patients taking warfarin, excessive application on the skin for muscle or joint pains may increase the chances of bleeding.

How to use Zhui Fong Yew

- How much to use

Rub on site of pain 3 to 4 times daily.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the

instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

Please follow the advice of the doctor or pharmacist.

- If you use too much (overdose)

If you have used too much Zhui Fong Yew, contact your doctor, an emergency department or a poisoning center immediately.

While you are using Zhui Fong Yew

- Things you must do

Inform your doctor or pharmacist of all the medications you are taking.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying to a larger skin area. Discontinue use if irritation develops.

- Things you must not do

Avoid contact with the eyes and mucous membrane. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads.

Do not take any new medicines without consulting your doctor.

Do not give Zhui Fong Yew to anyone else, even if they have the same symptoms or condition as you.

Side effects

Skin irritation, redness or swelling, burning or stinging sensation. Adverse reaction is rare. Symptoms of dizziness and nausea may occur.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78018464/ 8470, or visiting the website nptra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI).

Storage and Disposal of Zhui Fong Yew

- Storage

Store below 30°C. Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Brown-coloured oil with characteristic odour.

- Ingredients

- Active ingredient
Menthol, methyl salicylate

- Inactive ingredients
Chlorophyll, Red 17, liquid paraffin

- MAL number:

MALXXXXXXXXXX

Manufacturer

Wellmetech Pharmaceutical Industries Sdn Bhd
Lot 8058, Kota Samarahan Industrial Park,
94300 Kota Samarahan, Sarawak,
Malaysia.

Product Registration Holder

Kian Feng Trading,
No. 81, Lalan Rokam 13, Pekan Razaki,
31350 Ipoh, Perak, Malaysia.

Date of revision

9/3/2018