

TAFLOTAN[®]-S OPHTHALMIC SOLUTION 0.0015% (single-dose container)

Tafluprost (0.0015%)

What is in this leaflet

1. What Taflotan-S is used for
2. How Taflotan-S works
3. Before you use Taflotan-S
4. How to use Taflotan-S
5. While you are using it
6. Side effects
7. Storage and Disposal of Taflotan-S
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Taflotan-S is used for

It is usually used to treat primary open-angle glaucoma or ocular hypertension. (a buildup of fluid inside the eye).

How Taflotan-S works

This medicine promotes aqueous humour (intraocular fluid) outflow in order to lower pressure in the eye.

Before you use Taflotan-S*- When you must not use it*

Be sure to tell your doctor and pharmacist,

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines.
- If you have history of hypersensitivity to any of the ingredients of this product.

- Before you start use it

Be sure to tell your doctor and pharmacist,

- If you are patient with aphakic/pseudophakic eyes (absence of the lens of the eye due to surgical removal), bronchial asthma/history of bronchial asthma, or intraocular inflammation (iritis, uveitis).
- If you are pregnant or breastfeeding.

- Taking other medicines

Be sure to tell your doctor and pharmacist,

- If you are taking any other medicinal products. (Some

medicines may interact to enhance or diminish medicinal effects.

Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

How to use Taflotan-S

Before applying, wash your hands thoroughly with soap and running water. Next, twist and remove the tip of the container and discard 1 to 2 drops. Then, pull down your lower eyelid so that the dropper tip does not touch eye or eyelashes and apply the medicine. After applying, keep your eyes closed for 1 to 5 minutes pressing the inner corner of the eye lightly. Wipe off any excess liquid with a wet gauze or tissue immediately, or wash your face with eyes closed. After applying, make sure to dispose the dropper. Once the cap is open, remaining solution must not be stored for later applying. If you need to apply other eye drops at the same time, wait more than 5 minutes.

- How much to use

In general, apply 1 drop at a time, once a day. Strictly follow the instructions. Do not use more than once daily because more frequent administration may lessen the intraocular pressure-lowering effect (reduce the pressure of buildup of fluid inside the eye).

- When to use it

Once a day.

- How long to use it

Do not stop applying this medicine unless your doctor instructs you to do so.

- If you forget to use it

If you miss a dose, apply 1 drop as soon as possible in case of the same day. If you notice on the next day, skip the missed dose and continue your regular dosing schedule. You should not apply twice a day or 2 drops at one time.

- If you use too much (overdose)

If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.

While you are using it*- Things you must do*

- Take your medication exactly as directed by your doctor.
- Consult your doctor if you have any questions regarding the medication
- Tell your doctor if you notice or are worried by any of the side effects listed or notice any other effects not listed.

- Things you must not do

Do not give your medicine to anyone else, even if they have the same condition as you. It may not be safe for another person to use Taflotan-S.

Do not stop using your Taflotan-S or change the dosage without checking with your doctor.

- Things to be careful of

This medicine may cause blurred vision temporarily after applying. Do not operate machinery or drive a car until these symptoms recover.

Side effects

The most commonly reported adverse reactions include bloodshot eyes, changes in eyelashes (increased length, thickness and number of lashes), itchy/stinging eyes, eye pain, eyelid pigmentation (darkening around the eyes), corneal epithelial disorder, and hypertrichosis (excessive hair growth) in eyelid, etc. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

- changes in the color of the iris [iris pigmentation]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

TAFLOTAN[®]-S OPHTHALMIC SOLUTION 0.0015% (single-dose container)

Tafluprost (0.0015%)

You may report any side effects or adverse drug reactions directly to The National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.moh.gov.my (Consumers → Reporting → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Product Registration Holder

Santen Pharma Malaysia Sdn. Bhd.
Unit #23A-10, Q Sentral,
No. 2A, Jalan Stesen Sentral 2,
Kuala Lumpur Sentral,
50470 Kuala Lumpur, Malaysia.

Date of revision

22/06/2026

Storage and Disposal of Taflotan-S

- Storage

Keep out of reach of children.

Store in a refrigerator (2-8°C) and protect from light.

After opening the aluminum foil pouch

Stored below 25°C and protect from light. Used within one month. Use the product in the pouch which has been opened first.

- Disposal

Discard the remainder. Do not store them. Ask your pharmacy or medical institution how to discard.

Product Description

- What it looks like

Clear, colourless, sterile aqueous ophthalmic solution.

- Ingredients

- Active ingredient

Each mL contains 15µg of tafluprost.
Each container (0.3mL) contains 4.5µg of tafluprost.

- Inactive ingredients

Polysorbate 80, concentrated glycerin, disodium edetate hydrate, sodium dihydrogen phosphate dihydrate, sodium hydroxide, hydrochloric acid and purified water.

- MAL number(s):

MAL17085062ARZ

Manufacturer

Santen Pharmaceutical Co., Ltd.
Noto plant: 2-14, Shikinami,
Hodatsushimizu-cho,
Hakui-gun, Ishikawa, Japan.