

pharmaniaga®



Germacid Cream

Fusidic acid 2%

Ointment

Sodium fusidate 2%

DESCRIPTION

Pharmaniaga Germacid Cream

A smooth, white cream.

Pharmaniaga Germacid Ointment

A smooth, white ointment.

COMPOSITIONS

Pharmaniaga Germacid Cream

Each gram contains Fusidic acid 2% w/w.

Pharmaniaga Germacid Ointment

Each gram contains Sodium fusidate 2% w/w.

Also contains Chlorocresol 0.1% w/w as preservative.

PHARMACODYNAMICS

Pharmacotherapeutic group: Other antibiotics for topical use, ATC code: D06AX01

Fusidic acid is a strong antibacterial agent. Fusidic acid and its salts exhibit fat and water solubility properties with strong surface activity, and show uncommon ability to penetrate intact skin.

Fusidic acid or sodium fusidate is very active against *staphylococci*. Concentration of 0.03-0.12 mcg fusidic acid per ml inhibit nearly all strains of *Staphylococcus aureus*. Topical application of fusidic acid is also effective against *Streptococci*, *Corynebacterial*, *Neisseria* and certain *Clostridium* Strain.

PHARMACOKINETICS

Fusidic acid is well absorbed from the gastrointestinal tract. It is widely distributed into tissues and body fluids, including bone, pus and synovial fluid; it penetrates cerebral abscesses but does not enter cerebrospinal fluid in applicable amounts. About 95% or more of fusidate in circulation is bound to plasma protein.

Fusidic acid is excreted in bile, almost entirely as metabolites, some of which have weak antimicrobial activity. About 2% appears unchanged in the faeces. Little is excreted in the urine or removed by haemodialysis.

INDICATIONS

Pharmaniaga Germacid Cream

Pharmaniaga Germacid Cream is indicated for treatment of skin infections caused by staphylococci, streptococci, *Propionibacterium* acnes, *Corynebacterium minutissimum*, and other organisms sensitive to fusidic acid. The most important indications being: Impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia, erythrasma.

Pharmaniaga Germacid Cream is an invisible, non-staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

Pharmaniaga Germacid Ointment

For treatment of skin lesions primary or secondarily infected with *Staphylococci*,

Streptococci, *Corynebacterium minutissimum* and other sodium fusidate sensitive organisms. The most important indications being: impetigo, infected wounds, folliculitis, boils, sycosis barbae, carbuncles, hidradenitis, paronychia and erythrasma.

CONTRAINDICATIONS

Known hypersensitivity to fusidic acid or sodium fusidate infection caused by non-susceptible organisms. In particular *Pseudomonas aeruginosa*.

Patients with hypersensitivity to any of the ingredients.

PRECAUTIONS

Avoid contact with eyes.

Bacterial resistance is reported to have occurred when this product is applied topically. As in all topical antibiotics, extended and recurrent applications may increase the risk of contact sensitization and development of antibiotic resistance.

Fusidic cream contains cetyl alcohol. This excipient may cause local skin reactions (e.g. contact dermatitis).

Instruct patients not to smoke or go near naked flames as there is risk of severe burn. Fabric (e.g. clothing, bedding, dressings, etc) that has been in contact with this product burns more easily and is a serious fire hazard. Washing clothing and bedding may reduce product build-up but not totally remove it.

DRUG INTERACTIONS

No interaction studies have been performed. Interactions with systemically administered medicinal products are considered minimal as the systemic absorption of topical fusidic acid or sodium fusidate is negligible.

155mm

PREGNANCY AND LACTATION

Pregnancy

There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effects.

Breast-feeding

Safety in nursing mothers has not been established.

Germacid Cream can be used during breast-feeding but is recommended to avoid applying Germacid Cream on the breast.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Fusidic acid or sodium fusidate administered topically has no or negligible influence on the ability to drive and use machines.

ADVERSE REACTIONS

Immune system disorders

Rare: Hypersensitivity

Eye disorders

Rare: Conjunctivitis

Skin and subcutaneous tissue disorders

Uncommon: Dermatitis (including dermatitis contact, eczema), rash*, pruritus and erythema.

*Various types of rash reactions such as erythematous, pustular, vesicular, maculo-papular and papular have been reported. Rash generalised has also occurred.

Rare: Angioedema, urticaria and blister

General disorders and administration site conditions

Uncommon: Application site pain

(including skin burning sensation) and application site irritation.

DOSAGE AND ADMINISTRATION

Pharmaniaga Germacid Cream is applied to the affected area 2-3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

Pharmaniaga Germacid Ointment is applied to affected area 2-3 times daily, for a period of 7 days. It can be used with or without a covering dressing.

ROUTE OF ADMINISTRATION

Topical.

OVERDOSAGE

Overdose is unlikely to occur.

STORAGE CONDITIONS

Store below 30°C.

Protect from light.

Keep container tightly closed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

Tubes of 5g and 15g.

Date of revision: 01 March 2024

PRODUCT REGISTRATION HOLDER:

Pharmaniaga Manufacturing Berhad (198001006232)

No. 11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650 Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

MANUFACTURER:

Idaman Pharma Manufacturing Sdn Bhd (200401023395)

Lot 24 & 25, Jalan Perusahaan Lapan, Bakar Arang Industrial Estate, 08000 Sungai Petani, Kedah, Malaysia.

Lt-401.00

148mm