

ACTIMOL TABLET 650MG

Paracetamol 650 mg

What is in this leaflet

1. What Actimol Tablet 650 mg is used for
2. How Actimol Tablet 650 mg works
3. Before you use Actimol Tablet 650 mg
4. How to use Actimol Tablet 650 mg
5. While you are using it
6. Side effects
7. Storage and disposal of Actimol Tablet 650 mg
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What Actimol Tablet 650 mg is used for

Actimol Tablet 650 mg contains 650 mg of paracetamol. It is used for relief of headache, backache, period pain, and aches due to cold and flu, pain related to mild arthritis and fever.

How Actimol Tablet 650 mg works

Actimol Tablet 650 mg has analgesic (pain relief) and antipyretic (anti-fever) properties.

Before you use Actimol Tablet 650 mg

-When you must not use it

Do not take Actimol Tablet 650 mg if you are allergic to paracetamol or acetaminophen which is another name for paracetamol.

If you are not sure whether you should use this medicine, talk to your doctor or pharmacist first.

-Before you start to use it

Tell your doctor or pharmacist if you have allergies to paracetamol/ acetaminophen, any other medicines, foods, preservatives or dyes.

Your doctor or pharmacist needs to know before you take Actimol Tablet 650 mg if you have any medical condition.

Discuss with your doctor or pharmacist first if you are pregnant, plan to become pregnant or are breastfeeding

Taking other medicines

This preparation contains Paracetamol/acetaminophen.

Do not take any other Paracetamol/ acetaminophen containing medicines at the same time.

The effects of some medicines may be affected when used with Actimol Tablet 650 mg at the same time.

Tell your doctor or pharmacist what other prescription and non-prescription medicines that you are taking.

Avoid drinking alcohol while taking Actimol Tablet 650 mg. and kidney impairment, please consult your doctor for dose adjustments.

How to use Actimol Tablet 650 mg

Swallow with water or other fluids. Always take the medicine exactly as your doctor or pharmacist has told you. Read the direction on the label carefully. If you have any concerns about how to take this medicine, talk to your doctor or pharmacist.

-How much to use

Your pharmacist can advise you how many doses are needed each day and for how long you may need to take it. This depends on your condition and whether or not you are taking any other medicines.

Adults: 1 to 1½ tablets, three or four times a day (Maximum: 6 tablets a day).

Children should not be given paracetamol for more than 3 days without consulting a doctor.

Should not be taken more than four times a day and with an interval not less than four hours between each dose.

-When to use it

Follow the direction label carefully and as instructed by your doctor or pharmacist.

-How long to use it

Follow the direction on the label carefully and as instructed by your doctor or pharmacist.

-If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed. This may increase the chance of you getting unwanted side effects.

-If you use too much (overdose)

Immediately go to the emergency department at your nearest hospital if you or

ACTIMOL TABLET 650MG

Paracetamol 650mg

anyone else has taken more than the recommended amount of this medication even if there are no signs of discomfort or poisoning because of the risk of liver failure.

While you are using Actimol Tablet 650 mg

-Things you must do

- Take the medication exactly as prescribed by your doctor or as instructed by your pharmacist and as on the label of your medicine. Talk to your doctor or pharmacist if need any clarification or your symptoms do not improve.
- When seeking other medical treatment, inform all the doctors, dentists and pharmacists that you are taking Actimol Tablet 650 mg.
- Tell your doctor if you experienced persistent headaches.
- Tell your doctor or pharmacist immediately if you become pregnant while taking this medication.

-Things you must not do

Do not give your medicine to anyone else, even if their symptoms seem similar to yours.

Do not take your medicine to treat any other complaints unless your doctor or pharmacist tells you to.

-Things to be careful of

If you are pregnant or are breast-feeding, you should not take Actimol Tablet 650 mg without discussing it with your doctor or pharmacist first.

Side Effects

All medicines may have some unwanted side effects. Unwanted side effects do not always occur in every person. Tell your doctor or pharmacist as soon as possible if you do not feel well while you are

taking Actimol Tablet 650 mg.

Ask your doctor or pharmacist to answer any questions you may have.

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. If these reactions occur, stop use and seek medical assistance right away.

Side effects such as skin reactions including skin rashes, swelling of the lower layer of skin, serious skin conditions such as Stevens Johnson Syndrome/Toxic Epidermal Necrolysis, low platelet count, lowered white blood cell count, constriction of the air passages of the lung (more likely in asthmatics sensitive to aspirin or other nonsteroidal anti-inflammatory drugs (NSAIDs)) and sudden onset liver failure have been reported.

You are advised to consult doctor/ or pharmacist for further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Actimol Tablet 650 mg

-Storage

Keep all medicines out of sight and reach of children.

Do not take this medicine after the expiry date imprinted on the product packaging.

Keep your medicine in a cool dry place where temperature stays below 30°C.

Protect from light and moisture.

-Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

White, oblong-shaped tablets with Pharmaniaga icon on one side and breakline on the other.

-Ingredients

Active ingredient(s):

Each tablet contains Paracetamol 650 mg.

-MAL Number:

MAL16125044XZ

Manufacturer

Pharmaniaga Manufacturing Berhad (198001006232)
11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650, Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

Product Registration Holder

Pharmaniaga Manufacturing Berhad (198001006232)
11A, Jalan P/1, Kawasan Perusahaan Bangi, 43650, Bandar Baru Bangi, Selangor Darul Ehsan, Malaysia.

Date of revision:

-25/04/2022

Serial Number:

- NPRA(R2/1)220407/866