

BAXTER CONFIDENTIAL - INTERNAL USE ONLY

Part Number: BE3504172	Date: 30JUN22	Proofread No.: 05
Designer: Z.B.	Page: 1 of 1	
Colour Reference: BLACK		

IYDB3WH1E
Baxter

Central Vein

With Electrolytes

For single use only

Read the package leaflet before use

Keep out of the sight and reach of children

Jauhi Daripada Kanak-Kanak

Controlled Medicine /
Ubat Terkawal

Store below 30°C

Do not freeze

Store in the overpouch

Chemical and physical in-use stability has been demonstrated for 7 days at 2°C- 8°C followed by 48 hours at temperature not exceeding 30°C.

Do not infuse before, simultaneously or after administration of blood through the same equipment

Use only if the amino acid and glucose solutions are clear, the lipid emulsion and the final mixture after reconstitution are homogenous and with a milky appearance and the container undamaged

Do not connect in series

Do not infuse without opening the seals and mixing the contents of the 3 compartments **500**

After reconstitution, immediately administer the product

OLIMEL N12E Emulsion for Infusion

For central intravenous administration
1,000 mL of the final mixture provides the following:

Total calories approx.	950 kcal
Non-protein calories	640 kcal
Nitrogen	12.0 g
Non-protein calories / nitrogen ratio	53 kcal/g
Glucose	73.3 g
Lipids	35.0 g
Glucose / lipid calories ratio	45/55

Amino acids	75.9 g
Glucose calories	290 kcal
Lipid calories	350 kcal
Lipid / total calories	37%
Electrolytes:	
Sodium	35.0 mmol
Potassium	30.0 mmol
Magnesium	4.0 mmol
Calcium	3.5 mmol
Phosphate	15.0 mmol
Acetate	70 mmol
Chloride	45 mmol
pH	6.4
Osmolarity	1,270 mOsm/L

Bag

Nitrogen g	12.0
Total kcal	950
Non-protein calories	640

1,000 mL

Formula for 1,000 mL after mixing:

Refined olive oil + refined soybean oil 35.00 g, Alanine 10.99 g, Arginine 7.44 g, Aspartic acid 2.20 g, Glutamic acid 3.79 g, Glycine 5.26 g, Histidine 4.53 g, Isoleucine 3.79 g, Leucine 5.26 g, Lysine acetate 8.43g (equivalent to Lysine 5.97g), Methionine 3.79 g, Phenylalanine 5.26 g, Proline 4.53 g, Serine 3.00 g, Threonine 3.79 g, Tryptophan 1.26 g, Tyrosine 0.20 g, Valine 4.86 g, Sodium acetate, trihydrate 1.5 g, Sodium glycerophosphate, hydrated 3.67 g, Potassium chloride 2.24 g, Magnesium chloride, hexahydrate 0.81 g, Calcium chloride, dihydrate 0.52 g, Glucose Monohydrate 80.67g (equivalent to Glucose 73.33g)

Excipients:

purified egg phospholipids (source: chicken egg yolk), glycerol, sodium oleate, sodium hydroxide, water for injections, glacial acetic acid, hydrochloric acid.

After reconstitution: Emulsion for Infusion 1,000 mL in a 3-compartment bag

Glucose solution 27.5% : 267 mL

Amino acid solution 14.2% : 533 mL

Lipid emulsion 17.5% : 200 mL


Manufactured by: Baxter SA, 7860 Lessines, Belgium

Product Registration Holder: Baxter Healthcare (Malaysia) Sdn. Bhd., B-21-3A, The Ascent, Paradigm, No.1, Jalan SS7/26A, Kelana Jaya, 47301 PJ, Selangor, Malaysia

MAL22066005AZ

POM

BE-35-04-172



Lot:



Mfg Date

EXP:

IYDB3WH1E