

Baxter
IYDB3WH1E

6 x 1,000 mL

OLIMEL N12E, emulsion for infusion

FOR CENTRAL INTRAVENOUS ADMINISTRATION

	1,000 mL
27.5% Glucose solution (corresponding to 27.5 g/100 mL)	267 mL
14.2% Amino acid solution (corresponding to 14.2 g/100 mL)	533 mL
17.5% Lipid emulsion (corresponding to 17.5 g/100 mL)	200 mL

Excipients:

Lipid emulsion compartment:

purified egg phospholipids (source: chicken egg yolk), glycerol, sodium oleate, sodium hydroxide, water for injections.

Compartment of amino-acid solution with electrolytes :

glacial acetic acid, water for injections.

Compartment of glucose solution with calcium :

hydrochloric acid, water for injections.

After reconstitution

Emulsion for Infusion

For single use only.

Read the package leaflet before use.

Keep out of the sight and reach of children. Jauhi Daripada Kanak-Kanak

Controlled Medicine / Ubat Terkawal

BE2503492

EXP: 99/9999

Lot: XXXXXXXXXXXXX

Mfg Date: 99/9999



Baxter
IYDB3WH1E

6 x 1,000 mL

OLIMEL N12E, emulsion for infusion

FOR CENTRAL INTRAVENOUS ADMINISTRATION

With electrolytes.

Do not infuse before, simultaneously or after administration of blood through the same equipment.

Use only if the amino acid and glucose solutions are clear, the lipid emulsion and the final mixture after reconstitution are homogenous and with a milky appearance and the container undamaged.

Do not connect in series.

Do not infuse without opening the seals and mixing the contents of the 3 compartments. After reconstitution, immediately administer the product.

Do not freeze. Store Below 30°C.

Store in the overpouch.

Manufactured by:

Baxter SA,

7860 Lessines, Belgium

Product Registration Holder: Baxter Healthcare (Malaysia)

Sdn. Bhd., B-21-3A, The Ascent,

Paradigm, No.1, Jalan SS7/26A, Kelana

Jaya, 47301 PJ, Selangor, Malaysia

MAL22066005AZ

BE2503492

EXP: 99/9999

Lot: XXXXXXXXXXXXX

Mfg Date: 99/9999

