

BIO-BACTEDENE GEL 10%w/w

Povidone-Iodine 10g /100 g(equivalent to iodine 1 g)

What is in the leaflet

1. What Bio - Bactedene gel is used for
2. How Bio - Bactedene gel works
3. Before you use Bio-Bactedene gel
4. How to use Bio-Bactedene gel
5. While you are using Bio- Bactedene gel
6. Side Effects
7. Storage and Disposal of Bio -Bactedene gel
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of Revision

What Bio -Bactedene gel is used for

Bio - Bactedene gel is used for prevention of infection in cuts, abrasions, burns and use for pre-operative disinfection of the skin.

How Bio- Bactedene gel works

The active ingredient in Bio-Bactedene gel which is Povidone-iodine, is able to kill the broad range of germs.

Before you use Bio-Bactedene gel

-When you must not use it

Tell your doctor if any of the following are true for you as patients with these conditions should not use povidone-iodine

- if you are allergic (hypersensitive) to polyvinylpyrrolidone, iodine or any of the other ingredients of this medicine (listed in section Product Description)
- if you have former or current abnormal thyroid function or goiter
- if your age is less than 2 years old.
- If you are taking lithium

-Before you start to use it

Talk to your doctor or pharmacist before using Bio -Bactedene gel

- if you have former or current abnormal thyroid function or goiter
- if your kidneys are not working well

Check with your doctor if any of the following applies to you:

- if you have had a serious allergic reaction or a serious skin reaction to any medicine
- If you are taking lithium or sodium picosulfate
- If you would have your thyroid function tested or several types of tests for detection of occult blood in faeces or blood.
- You are younger than 2 years

-Using with other medicines

Some medicines can affect the way other medicines work. Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription:

-Lithium: Use with concurrent lithium therapy has been shown to exhibit additive hypothyroidic effects.

-Sodium picosulfate: Antibiotics may diminish the therapeutic effect of Sodium picosulfate.

Tell your doctor or pharmacist if you are taking, or have recently taken any of the medicines listed above or any other medicines.

How to use Bio- Bactedene gel

-How much to use

- Apply the proper amount of Bio- Bactedene gel on the affected areas of the skin 2-3 times daily

When to use it

Use as directed by your doctor or pharmacist.

-How long to use it

Do not use this product for longer than 1 week unless directed by a doctor because prolonged exposure may cause irritation, or, rarely, severe skin reactions.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to use the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have orally taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

Use this medicine exactly as the doctor or pharmacist has told you.
Tell all the doctors or pharmacists treating you that you are using Bactedene gel

-Things you must not do

- This product is for external use only. Do not use this product in the eyes or over large areas of the body.
- Do not heat prior to application
- Do not stop using the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist
- Do not give Bio -Bactedene gel to anyone else, even if they have the same symptoms or condition as you.

BIO-BACTEDENE GEL 10%w/w

Povidone-Iodine 10 grams (Iodine 1 gram)/100 grams

Side Effects

Like all medicines, Bio-Bactedene gel can have side effects, although not everybody gets them.

Some common side effects are listed below:

Common side effects: local irritation, skin burns and sensitivity reactions

Serious side effects: anaphylactic reactions, anaphylactoid reactions, anaphylactic shock, goiter, hypothyroidism, hyperthyroidism, metabolic acidosis and acute renal failure.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reaction directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my/ [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

MAL number: MAL20046134XZ

Manufacturer

MacroPhar Co., Ltd
89 Soi Pattanakarn 20 Yaek 4,
Pattanakarn Road., Suan Luang,
Suan Luang, Bangkok 10250, Thailand.

Product Registration Holder

TOP BIOPHARMA SDN. BHD.
No. 23A, Jalan Kesuma 3/11, Bandar Tasik
Kesuma, 43700 Beranang, Selangor D.E.
Malaysia..

Date of Revision

17/07/2020

Serial number

NPRA (R2) 20/024

Storage and Disposal of Bio-Bactedene gel

Storage:

- Store below 30 °C.
- Keep medicine out of reach of children.

Disposal:

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

Brown gel

-Ingredients

- * Active ingredient – Povidone iodine
- * Inactive ingredient – Sodium bicarbonate, purified water, polyethylene glycol 400, polyethylene glycol 4000