

BROVENT SYRUP

Bromhexine hydrochloride (8mg/5ml)

What is in this leaflet

1. What Brovent Syrup is used for
2. How Brovent Syrup works
3. Before you take Brovent Syrup
4. How to use Brovent Syrup
5. While you are using it
6. Side Effects
7. Storage and Disposal of Brovent Syrup
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Brovent Syrup is used for

Brovent Syrup is used to reduce the thickness of phlegm.

How Brovent Syrup works

Brovent Syrup contains bromhexine which is a mucolytic agent, that reduces the thickness of the phlegm and also increase the secretion of mucus. Bromhexine is also an expectorant, which means it can promote the expulsion of phlegm.

Before use Brovent Syrup

- When you must not use it :

Do not take Brovent Syrup if you are allergic to bromhexine or any other ingredients of this product.

- Before you start to use it:

- Consult with your doctor if you are pregnant, planning to get pregnant or breast-feeding your baby.
- Consult your doctor if you are currently taking any antibiotics as bromhexine may affect antibiotic absorption.
- Consult your doctor if you have the following medical conditions:
 - asthma
 - severe liver and kidney problems
 - stomach ulcer

-Taking other medicines:

- Tell the doctor if you are taking any other medicines including any medicine that you buy without

prescription from a pharmacy, supermarket or health food shop.

- Bromhexine may increase the amount of antibiotic absorption into the body.

How to take Brovent Syrup

- How much to take:

- Adult: 5 - 10 ml (1-2 teaspoons)
- Children (6 - 12 years) : 2.5 - 5 ml (half teaspoon to 1 teaspoon)

-When to take it :

To be taken 3 times a day, or as instructed by your doctor.

-How long to take it :

Use Brovent Syrup for the duration that the doctor recommends.

-If you forget to take it :

Consult your doctor or pharmacist on what you should do if you forget to use it. Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you take too much (overdose) :

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are taking Brovent Syrup

-Things you must do :

- Take your medicine exactly as your doctor has told you.
- Tell all doctors, dentists and pharmacists treating you that you are taking Brovent Syrup.
- Inform your doctor if you are pregnant while taking this medication.

-Things you must not do :

- Do not stop taking the medicine unless advised by your doctor.

- Do not take any new medicines without consulting your doctor.
- Do not give Brovent Syrup to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of :

- Taking bromhexine may cause an increase in the flow of secretions.
- Consult your doctor if the symptoms last longer than 14 days, or if the symptoms increase in spite of treatment with bromhexine.

Side Effects

Like all medicines, Brovent Syrup can cause side effects, although not everybody gets them. Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Brovent Syrup may cause severe allergy and serious skin reactions. Stop using Brovent Syrup and seek medical assistance immediately if you experience any of the following symptoms:

- allergic reaction, including swelling of the face, lips, mouth, tongue or throat which may cause difficulty in swallowing or breathing.
- severe allergy: breathing difficulties, light-headedness, skin swellings or rash.
- severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.
- hives and itchiness.
- nausea, stomach pain, vomiting, and diarrhoea.
- headache.
- involuntary urination at night.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my

[Consumers ► Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

BROVENT SYRUP

Bromhexine hydrochloride (8mg/5ml)

Storage and Disposal of Brovent Syrup

-Storage

Store below 30°C in a dry place, protected from direct sunlight.

Keep out of reach of children

Shake well before use.

-Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

Product Description

-What it looks like

Yellow-coloured clear syrup with a fruity odour and has a sweet taste.

-Ingredients

-Active Ingredient(s)

Bromhexine hydrochloride.

-Inactive Ingredients

Tween 80, Tutti fruity, Quinolline yellow, Propylene glycol, purified water, sugar, liquid glucose.

-MAL Number:

MAL19940582XCZ

Manufacturer

Terapeutics Sdn. Bhd. (590500-W)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat S.P.S. Penang, Malaysia

Product Registration Holder

Zontron Pharmaceuticals Sdn. Bhd.
(445695-T)

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat S.P.S. Penang, Malaysia

Date of Revision

02/09/2021

Serial Number

NPRA(R4/1)180821/00068