SCABICIDE LOTION

Crotamiton (10% w/v)

What is in this leaflet

1. What Scabicide Lotion is used for
2. How Scabicide Lotion works
3. Before you use Scabicide Lotion
4. How to use Scabicide Lotion
5. While you are using it
6. Side effects
7. Storage and Disposal of Scabicide Lotion
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Scabicide Lotion is used for

Scabicide Lotion is used for the treatment of scabies. Scabicide Lotion is for external use only.

How Scabicide Lotion works

Scabicide Lotion contains crotamiton, which is an anti-scabies and anti-itch agent, is highly effective in removing scabies and relief of itchiness.

Before you use Scabicide Lotion

- When you must not use it

Do not use Scabicide Lotion if:

- you are allergic or hypersensitive to crotamiton or any of the ingredients contained in this product.
- you are having skin disease or skin inflammation. Consult your doctor or pharmacist.
- it is not prescribed by a qualified doctor or pharmacist to you.

- Before you start to use it

Inform your doctor if:

- you are pregnant, trying to get pregnant or think you may be pregnant.
- you are breast-feeding.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Scabicide Lotion

- How much to use

For treatment of scabies:

- Adults and children: Rub onto the skin of the whole body from the chin down, paying particular attention to all folds and creases.
- Scabicide Lotion can be used in children. However, in children under three years of age, Scabicide Lotion should NOT be applied more than once in a day.

- When to use it

Use as directed by your doctor or pharmacist.

- The application should be repeated once daily, preferably in the evening,
- A second application is advisable 24 hours later.

DIRECTIONS OF USAGE:

After you have taken a warm bath, the skin should be well-dried. Rub Scabicide Lotion onto your entire body surface (except for your face and scalp) until there are no traces of Scabicide Lotion remains visible on your skin. Special attention should be given to areas that are particularly susceptible to infestation (e.g. between the fingers, wrists, under arm and around the genitals). Areas where there is pus (white-yellowish liquid discharge from wounds) should be covered with a dressing impregnated with Scabicide Lotion. During treatment with Scabicide Lotion, you may take a bath shortly before the next application. After treatment with Scabicide Lotion is finished, you are advised to take, a cleansing bath followed by a change of bed linen and underclothing.

- How long to use it

Treatment of scabies should last for a total of 3-5 days, depending on the response. Consult with your doctor or pharmacist if symptoms persist.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

In case of accidental consumption, go to the Emergency Department of your nearest hospital. Symptoms may be observed such as nausea, vomiting and irritation of the inside lining of the cheeks (inside the mouth), throat and stomach. There have been reported cases of loss of consciousness and seizure, but rare.

While you are using it

- Things you must do

- Use your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacist treating you that you are using Scabicide lotion.
- Tell your doctor immediately if you become pregnant while using this medication.
- Consult your doctor if your condition does not improve or if it becomes worse.
- Clothing and bed linen should be changed the next morning. A cleansing bath should be taken after the last application.

- Things you must not do

- Do not use any new medicines without consulting your doctor or pharmacist.
- Do not stop using medication until the treatment course is finished, even

SCABICIDE LOTION

Crotamiton (10% w/v)

if you are feeling better after the first few days.

- Do not use this medication for weeping skin conditions.
- Do not give Scabicide lotion to anyone else, even if they have the same symptom or condition as you.
- Scabicide lotion should not be used in the inside lining of the cheeks (inside the mouth) and in or around the eyes, as direct contact with the eyelids may lead to eye inflammation. In case of
- accidental contact with the eyes or inside of the mouth, please rinse thoroughly with running water.
- Scabicide Lotion should not be applied in the presence of wounds, eczema, broken skin, or very inflamed skin.
  Consult your doctor if you have any of the skin conditions mentioned above.
- If you are breastfeeding and your doctor has prescribed you Scabicide Lotion, do not apply Scabicide Lotion to the nipple area.

- Things to be careful of

- If you develop allergy or severe irritation, discontinue treatment and seek immediate medical attention from your doctor.

Side effects

Like all medicines, Scabicide Lotion can cause side effects, although not everybody gets them.
The most commonly reported side effect during treatment is itchiness.
Skin inflammation and hypersensitivity reactions like rash, eczema, skin redness, skin irritation and swelling beneath the skin may occur, but rare.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Scabicide Lotion

- Storage

Keep all medicines out of the reach of children.
Store below 30°C in a dry place protected from direct light.

- Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help to protect the environment.

Product Description

- What it looks like

Clear, colourless and odourless lotion

- Ingredients

- Active ingredient(s)
Crotamiton

- Inactive ingredients
Alcohol denatured, PEG-400

- MAL number:

MAL19940584XCZ

Manufacturer

Teraputics Sdn. Bhd**.**
(590500-W)
Lot 10 & 11, PERDA Industrial Park, Lorong IKS Simpang Ampat B, 14100 Simpang Ampat, S.P.S., Pulau Pinang, Malaysia

Product Registration Holder:

Zontron Pharmaceuticals Sdn. Bhd. (445695-T)
Lot 10 & 11, PERDA Industrial Park, Lorong IKS Simpang Ampat B, 14100 Simpang Ampat, S.P.S., Pulau Pinang, Malaysia

Date of revision

10/06/2021

Serial Number

NPRA(R4/1)200521/00024