

CANDINIL CREAM/LOTION

Clotrimazole (1% w/w) / Clotrimazole (1% w/v)

What is in this leaflet

1. What Candinil Cream/Lotion is used for?
2. How Candinil Cream/Lotion it works
3. Before you use Candinil Cream/Lotion
4. How to use Candinil Cream/Lotion
5. While you are using it
6. Side effects
7. Storage and Disposal of Candinil Cream/Lotion
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Candinil Cream/Lotion is used for

Candinil Cream/lotion is used for the treatment of skin infections caused by fungus.

How Candinil Cream/Lotion works

Candinil Cream/lotion contains the active ingredients clotrimazole. Clotrimazole is an antifungal which works by killing the fungi causing the skin infections.

Before you use Candinil Cream/Lotion

- When you must not use it

Do not apply Candinil Cream/lotion if:

- you are allergic (hypersensitive) to Clotrimazole or any other ingredients of this medicine (listed in Product Description).

- Before you start to use it

Stop using the medicine if you get any skin reactions.

Pregnancy and breast-feeding:

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Tell your doctor or pharmacist about all the other medicines and skin products you use, including prescribed and non-prescribed medicines, vitamins, herbal supplements, and cosmetics.

How to use Candinil Cream/Lotion

- How much to use

Apply a thin layer over the affected area.

- When to use it

Use 2 or 3 times a day or as directed by your doctor or pharmacist.

- How long to use it

Use Candinil Cream/Lotion for the duration that the doctor recommends.

- If you forget to use it

Apply the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to apply the medicine and skip the missed dose. Do not apply a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Ingesting this medicine may cause dizziness, nausea or vomiting.

While you are using it

- Things you must do

- Use your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are using Candinil Cream/Lotion.
- Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

- Do not stop using the medicine unless advised by your doctor.
- Do not use any new medicines without consulting your doctor or pharmacist.
- Do not give Candinil Cream/lotion to anyone else, even if they have the same symptoms or condition as you.

- Do not put the cream/lotion in your mouth or swallow it.
- If you accidentally get cream/lotion in your eyes or mouth, wash immediately with water.

- Things to be careful of

- The cream contains cetylalcohol and stearyl alcohol, which may cause local skin reactions. Stop using the medicine if you get any skin reactions.
- Clotrimazole cream/lotion may cause damage to latex contraceptive (condom).
- Use alternative precautions for at least five days after using this product.
- Candinil Cream/lotion is for external use only.

Side effects

Like all medicines, Candinil Cream/Lotion can cause side effects, although not everybody gets them.

You may experience the following:

- allergic reaction such as fainting, difficulty in breathing, low blood pressure and hives.
- skin or tissue problem such as blisters, discomfort or pain, swelling, irritation, skin peeling or itchiness, rash, stinging or burning sensation.

If you experience any of the above effects or react badly to the cream in any other way not listed in this leaflet, tell your doctor or pharmacist immediately.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Candinil Cream/Lotion

- Storage

Keep all medicines out of the reach of children.

CANDINIL CREAM/LOTION

Clotrimazole (1% w/w) / Clotrimazole (1% w/v)

Store below 30°C. Store in a dry place protected from direct light.

Do not use this product after the expiry date or if the tube is leaking or if the tube nozzle cover is broken.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Candinil Cream: Odourless white soft cream

Candinil Lotion: Odourless, clear, colourless lotion

- Ingredients

- Active ingredient(s)

Clotrimazole

- Inactive ingredients

Candinil Cream: Chlorocresol, tween 80, liquid paraffin, white soft paraffin, cetomagrogol 1000, glyceryl monostearate, propylene glycol, cetostearyl alcohol, purified water.

Candinil Lotion: Propylene glycol, glycerin, PEG 400, alcohol denatured

- MAL number:

Candinil Cream: MAL19900279XCZ

Candinil Lotion: MAL19900300XCZ

Manufacturer:

Terapeutics Sdn. Bhd.

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat, Pulau Pinang, Malaysia

Product Registration Holder:

Zontron Pharmaceuticals Sdn. Bhd.

Lot 10 & 11, PERDA Industrial Park,
Lorong IKS Simpang Ampat B, 14100
Simpang Ampat, Pulau Pinang, Malaysia

Date of revision

23/11/2021

Serial Number

NPRA(R4/1)221121/00088