

1. Similar artwork is applicable to all pack sizes
2. All pack size(s) stated may not be available locally
3. Batch No, mfg. date and expiry date will be inkjet printed on the label.
4. Statement 'KKM/Kontrak KKM/Kementerian Kesihatan Malaysia/Tender KKM' will be inkjet/printed/stickered for tender items when granted.

**Indication:**  
Pentoxifylline is used in the treatment of peripheral occlusive arterial disease and arteriovenous disorders (e.g. intermittent claudication or rest pain) and trophic disturbances (e.g. gangrene and leg ulcers). It is also used in cerebrovascular disorders (sequel to cerebral arteriosclerosis such as difficulties in concentration, vertigo, impairment of memory)

**Dosage:**  
Usual dose is one tablet two or three times daily with meals and swallow whole with some liquid. Impaired renal function creatinine clearance below 10ml/min, may be necessary to reduce the dose to 800mg or 400mg daily.  
Keep out of reach of children.  
*Jauhi daripada kanak-kanak.*

Lot : Mfg : Exp :

# Trenlin™

## 400

### S. R. Tablet

Each tablet contains  
400mg pentoxifylline.

100 Tablets



DUOPHARMA

**CONTROLLED MEDICINE**

Route of administration: Oral  
Refer enclosed package insert for Contraindication /  
Precautions and Warnings.  
Keep container tightly closed.  
Store in a dry place below 25°C.

MAL05032541AZ SIN 10727P BRU09090324P

1500010260 N.2

Product Registration Holder: **Duopharma Manufacturing (Bangi) Sdn. Bhd.**  
Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate,  
43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufactured By: **Duopharma Manufacturing (Bangi) Sdn. Bhd.**  
Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate,  
43650 Bandar Baru Bangi, Selangor, Malaysia.



Pantone 185C



Black



PANTONE 2035 C



PANTONE 7624 C

Size: 113mm(W) x 58mm(H)