

BEAZYME TABLET

Papain 150,000 USP UNITS

What is in the leaflet

1. What Beazyme Tablet is used for
2. How Beazyme Tablet works
3. Before you use Beazyme Tablet
4. How to use Beazyme Tablet
5. While you are using it
6. Side effects
7. Storage and disposal of Beazyme Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Beazyme Tablet is used for

Beazyme Tablet is used in conjunction with other physical or chemotherapeutic measures for treatment or prevention of swelling and inflammation.

How Beazyme Tablet works

Beazyme contains proteolytic enzyme from carica papaya as active ingredient. Beazyme acts as an anti-inflammatory enzyme.

Before you use Beazyme Tablet

- When you must not take it

Do not take Beazyme Tablet if:

- You have blood clotting disease.
- You have any generalised or systemic infections.

- Before you start to use it

Tell your doctor if:

- You are pregnant or breastfeeding.

- Taking other medicines

Tell your doctor or pharmacist if you are taking the following

medicine as it may interact with your Beazyme Tablet:

- Any anticoagulant medicines to prevent blood clots

It may still be all right for you to take Beazyme Tablet and your doctor will be able to decide what is suitable for you.

How to use Beazyme Tablet

- How much to use

For oral administration only. For adults: Chew or suck 2 tablets 4 times daily for the first day, followed by one tablet 4 times daily for at least 5 days.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

- Take as stated in the leaflet without exceeding the dosage.
- Follow any recommendations and advices from your doctor or pharmacist.

- If you forget to use it

Do not double up the dosage as Beazyme Tablet may give you unacceptable side effects.

- If you use too much (overdose)

It is advisable for you to contact and inform your doctor if you did take Beazyme Tablet too much.

While you are using it

- Things you must do

- Discuss with your doctor or pharmacists do's and don't's while taking this medicine.

- Tell your doctor if unexpectedly you need to take any other medicines.
- Take Beazyme Tablet according to the recommended dosage.
- Tell your doctor or pharmacist if you experience any side effects that are not listed in this leaflet.

- Things you must not do

- Do not take your medicine more than the suggested frequency in a day.
- Do not over dosage the medicine during any single consumption.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, Beazyme Tablet can cause side effects although not everybody gets them. You are advised to consult doctor/ pharmacist for further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Monitoring by calling Tel: 03-78835490, or visiting the website npra.moh.gov.my (Public → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)).

Storage and Disposal of Beazyme Tablet

- Storage

Keep Beazyme Tablet out of reach of children.

Store in a dry place below 30°C.

Keep container well closed and protect from light.

- Disposal

Do not use Beazyme Tablet after the expiry date which is stated on the carton.

Product Description

- What it looks like

A8mm,diameter,round,flat ,marked with "BZYM"and scored on one side,plain on the other side,light blue tablet.

- Ingredients

- Active ingredient

- Proteolytic Enzyme From Carica Papaya

- Inactive ingredients:

- Talcum
- Icing sugar
- Liquid glucose
- Povidone
- Sodium carboxy methyl cellulose
- FD and C Blue #1
- FD & C Yellow #6
- Methyl salicylate flavour
- Oil of peppermint
- Isopropyl Alcohol
- Magnesium stearate

- MAL number

MAL19912640X

Product Registration Holder

Duopharma Manufacturing (Bangi) Sdn. Bhd.

Lot 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate,

43650 Bandar Baru Bangi Selangor Darul Ehsan

Date of Revision

06/08/2019

Serial Number:

NPRA (R2/1)191108/279

Manufacturer

Duopharma Manufacturing (Bangi) Sdn. Bhd.

Lot 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor Darul Ehsan