

GLYPRIN TABLET

Aspirin/Glycine (100mg/45mg)

What is in the leaflet

1. What Glyprin is used for
2. How Glyprin works
3. Before you use Glyprin
4. How to use Glyprin
5. While you are using it
6. Side effects
7. Storage and Disposal of Glyprin
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Glyprin is used for

Glyprin is usually used in patients with unstable angina, in patients who have had a heart attack and in patients at risk of a stroke.

How Glyprin works

Aspirin helps to prevent the blood from clotting.

This effect is used to reduce the risk of heart attack or stroke or other problems that may occur when a blood vessel is blocked by blood clots.

Ask your doctor if you have any questions about why this medicine has been prescribed for you. Your doctor may have prescribed it for another reason.

Before you use Glyprin

- When you must not use it

- Do not use this medicine if you are allergic to aspirin.
- Inform your doctor if you have ever had an ulcer in the stomach or intestine, have any conditions that increases your risk of bleeding.
- Do not use this medicine if you are taking other medicines to prevent blood clots or gastric irritants.
 - Do not use this medicine if you are pregnant or breast-feeding.
 - Do not use aspirin in children or teenagers unless

directed to do so by your doctor.

Do not take it after the expiry date printed on the pack. If you take it after the expiry date has passed, it may not work as well.

Do not take it if the packaging is torn or shows signs of tampering.

If you are not sure whether you should start taking this medicine, talk to your doctor or pharmacist.

- Before you start to use it

Tell your doctor or pharmacist if you have allergies to any other medicines, foods, preservatives or dyes.

- Taking other medicines

Inform your doctor and pharmacist about other medicines or alcohol you are taking, especially other blood thinning agents and medicines to control sugar level, other pain-killers or medicines used for joint pain (fenbufen, indomethacin, piroxicam), Medicine to treat cancer (methotrexate), medicines to treat epilepsy (phenytoin, valproic acid) and medicines that promotes excretion of uric acid in the urine (probenecid and sulphinpyrazone)

How to use Glyprin

Follow the directions for using this medicine provided by your doctor or pharmacist.

- How much to use

Take one tablet once a day at the same time every day. Tablet can be dispersed in water immediately before taken or swallowed whole.

- When to use it

Take one tablet once a day at the same time every day.

- How long to use it

Continue taking your medicine for as long as your doctor tells you.

- If you forget to use it

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Otherwise, use it as soon as you remember, and then go back to taking your medicine as you would normally.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering to take your medicine, ask your pharmacist for some hints.

- If you use too much (overdose)

Immediately telephone your doctor for advice or go to Accident and Emergency at your nearest hospital if you think that you or anyone else may have taken too much Glyprin.

Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Symptoms of an overdose may includes feeling warm, flushing, sweating, restlessness, dizziness, weakness, rapid breathing and confusion.

While you are using it

- Things you must do

Always follow your doctor's instructions carefully.

Tell your doctor if, for any reason, you have not taken your medicine exactly as directed.

Tell all the doctors, dentists and pharmacists who are treating you that you are taking Glyprin.

Tell your doctor if you become pregnant while taking Glyprin.

If you are about to start on any new medicine, remind your doctor and pharmacist that you are taking Glyprin.

- Things you must not do
Do not take more than the recommended dose unless your doctor tells you to.

- Things to be careful of
Be careful driving or operating machinery until you know how Glyprin affects you.

Side effects

These possible side-effects may go away during treatment as your body adjusts to the medicine. If they continue, are severe or bother you, tell your doctor or pharmacist.

- stomach pain or discomfort, indigestion or heartburn.
- nausea or vomiting.

Although the following side effects are less common, if they do occur they need medical attention. Contact your doctor as soon as possible if you notice:

- unusual bleeding or bruising
- black stools
- Vomiting blood
- Chest tightness and difficulty in breathing, and may provoke asthma attack in susceptible individuals'

Stop taking Glyprin and contact your doctor immediately if you notice:

- skin rash or itching
- any difficulty in breathing
- swelling of the eyelids, face or lips

Not all side effects reported for this medicine are included in this leaflet. If your general state of health worsens while taking this medicine, please consult your doctor.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Glyprin

- Storage

Store Glyprin below 25 C in a cool dry place. Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Glyprin is a white, round tablet, marked with 'GLY' on one side and plain on the other side.

- Ingredients

Active ingredient

- Aspirin 100mg
- Glycine 45mg

Inactive ingredients:

- Sodium Croscarmellose
- Sodium Stearyl Fumarate
- Sodium Lauryl Sulphate
- Microcrystalline Cellulose

- MAL number
MAL20001570XZ

Manufacturer

Duopharma Manufacturing (Bangi) Sdn. Bhd
Lot 2 & 4, Jalan P/7,
Section 13,
Bangi Industrial Estate
43650 Bandar Baru Bangi.
Selangor Darul Ehsan.

Product Registration Holder

Duopharma Manufacturing (Bangi) Sdn. Bhd
Lot 2, 4, 6, 8 & 10 Jalan P/7,
Section 13,
Bangi Industrial Estate
43650 Bandar Baru Bangi.
Selangor Darul Ehsan.

Date of Revision

30/11/2021

Serial Number:

NPRA(R2/1)211115/787