

CORXEVA TABLET

Telmisartan (40mg and 80mg)

What is in this leaflet

1. What CORXEVA is used for
2. How CORXEVA works
3. Before you use CORXEVA
4. How to use CORXEVA
5. While you are using it
6. Side effects
7. Storage and Disposal of CORXEVA
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What CORXEVA is used for

Corxeva is used to treat essential hypertension (high blood pressure). 'Essential' means that the high blood pressure is not caused by any other condition.

High blood pressure, if not treated, can damage blood vessels in several organs, which could lead sometimes to heart attack, heart or kidney failure, stroke, or blindness. There are usually no symptoms of high blood pressure before damage occurs. Thus it is important to regularly measure blood pressure to verify if it is within the normal range.

Corxeva is also used to reduce cardiovascular events (i.e. heart attack or stroke) in patients who are at risk because they have a reduced or blocked blood supply to the heart or legs, or have had a stroke or have high risk diabetes. Your doctor can tell you if you are at high risk for such events.

How CORXEVA works

Corxeva belongs to a class of medicines known as angiotensin II receptor antagonists. Angiotensin II is a substance produced in your body which causes your blood vessels to narrow, thus increasing your blood pressure. Corxeva blocks the effect of angiotensin II so that the blood vessels relax, and your blood pressure is lowered.

Before you use CORXEVA

- When you must not use it

- if you are allergic (hypersensitive) to telmisartan or any other ingredients included in Corxeva tablets (listed in section Product Description).
- if you are more than 3 months pregnant. (It is also better to avoid Corxeva in early pregnancy – see pregnancy section.)
- if you have severe liver problems such as cholestasis or biliary obstruction (problems with drainage of the bile from the liver and gall bladder) or any other severe liver disease.

If any of the above applies to you, tell your doctor or pharmacist before taking Corxeva.

- Before you start to use it

Please tell your doctor if you are suffering or have ever suffered from any of the following conditions or illnesses:

- Kidney disease or kidney transplant.
- Renal artery stenosis (narrowing of the blood vessels to one or both kidneys).
- Liver disease.
- Heart trouble.
- Raised aldosterone levels (water and salt retention in the body along with imbalance of various blood minerals).
- Low blood pressure (hypotension), likely to occur if you are dehydrated (excessive loss of body water) or have salt deficiency due to diuretic therapy (water tablets), low-salt diet, diarrhoea, vomiting, or haemodialysis.
- Elevated potassium levels in your blood.
- Diabetes.

You must tell your doctor if you think you are (or might become)

pregnant. Corxeva is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

In case of surgery or anaesthesia, you should tell your doctor that you are taking Corxeva.

The use of Corxeva in children and adolescents up to the age of 18 years is not recommended.

As with all other angiotensin antagonists, Corxeva may be less effective in lowering the blood pressure in black patients.

- Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription. Your doctor may need to change the dose of these other medicines or take other precautions. In some cases you may have to stop taking one of the medicines. This applies especially to the medicines listed below taken at the same time with Corxeva:

- Lithium containing medicines to treat some types of depression.
- Medicines that may increase blood potassium levels such as salt substitutes containing potassium, potassium-sparing diuretics (certain 'water tablets'), ACE inhibitors, angiotensin II receptor antagonists, NSAIDs (non steroidal antiinflammatory medicines, e.g. aspirin or ibuprofen), heparin, immunosuppressives (e.g. cyclosporin or tacrolimus), and the antibiotic trimethoprim.
- Diuretics ('water tablets'), especially if taken in high doses together with Corxeva, may lead to excessive loss of body water and low blood pressure (hypotension).

CORXEVA TABLET

Telmisartan (40mg and 80mg)

As with other blood pressure lowering medicines, the effect of Corxeve may be reduced when you take NSAIDs (non steroidal anti-inflammatory medicines, 1 e.g. aspirin or ibuprofen) or corticosteroids.

Corxeve may increase the blood pressure lowering effect of other medicines used to treat high blood pressure.

How to use CORXEVA

- How much to use

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The usual dose of Corxeve is one tablet a day. Try to take the tablet at the same time each day.

You can take Corxeve with or without food. The tablets should be swallowed with some water or other non-alcoholic drink. It is important that you take Corxeve every day until your doctor tells you otherwise. If you have the impression that the effect of Corxeve is too strong or too weak, talk to your doctor or pharmacist.

For treatment of high blood pressure, the usual dose of Corxeve for most patients is one 40 mg tablet once a day to control blood pressure over the 24 hour period. However, sometimes your doctor may recommend a lower dose of 20 mg or a higher dose of 80 mg.

Alternatively, Corxeve may be used in combination with diuretics ('water tablets') such as hydrochlorothiazide which has been shown to have an additive blood pressure lowering effect with Corxeve.

For reduction of cardiovascular events, the usual dose of Corxeve is one 80 mg tablet once a day. At the beginning of the preventive therapy with Corxeve 80 mg, blood pressure should be frequently monitored.

If your liver is not working properly, the usual dose should not exceed 40 mg once daily.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Corxeve for as long as your doctor recommends.

- If you forget to use it

If you forget to take a dose, do not worry. Take it as soon as you remember then carry on as before. If you do not take your tablet on one day, take your normal dose on the next day. Do not take a double dose to make up for forgotten individual doses.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

- If you use too much (overdose)

If you accidentally take too many tablets, contact your doctor, pharmacist, or your nearest hospital emergency department immediately.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Corxeve.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give Corxeve to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Pregnancy and lactation

Pregnancy

You must tell your doctor if you think you are (or might become) pregnant. Your doctor will normally advise you to stop taking Corxeve before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Corxeve.

Corxeve is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

Breast-feeding

Tell your doctor if you are breastfeeding or about to start breastfeeding. Corxeve is not recommended for mothers who are breast-feeding, and your doctor may choose another treatment for you if you wish to breastfeed, especially if your baby is newborn, or was born prematurely.

Children and adolescents

The use of Corxeve in children and adolescents up to the age of 18 years is not recommended.

Driving and using machines

No information is available on the effect of Corxeve on the ability to drive or operate machinery. Some people feel dizzy or tired when they are treated for high blood pressure. If you feel dizzy or tired, do not drive or operate machinery.

Side effects

Like all medicines, Corxeve can cause side effects, although not everybody gets them.

Some side effects can be serious and need immediate medical attention:

You should see your doctor immediately if you experience any of the following symptoms: Sepsis* (often called "blood poisoning", is a severe infection with whole-body inflammatory response), rapid swelling of the skin

CORXEVA TABLET

Telmisartan (40mg and 80mg)

and mucosa (angioedema); these side effects are rare but are extremely serious and patients should stop taking the product and see their doctor immediately. If these effects are not treated they could be fatal.

Possible side effects of Corxeva:

Common side effects (may affect up to 1 in 10 people): Low blood pressure (hypotension) in users treated for reduction of cardiovascular events.

Uncommon side effects (may affect up to 1 in 100 people):

Urinary tract infections, upper respiratory tract infections (e.g. sore throat, inflamed sinuses, common cold), deficiency in red blood cells (anaemia), high potassium levels, difficulty falling asleep, feeling sad (depression), fainting (syncope), feeling of spinning (vertigo), slow heart rate (bradycardia), low blood pressure (hypotension) in users treated for high blood pressure, dizziness on standing up (orthostatic hypotension), shortness of breath, abdominal pain, diarrhoea, discomfort in the abdomen, bloating, vomiting, itching, increased sweating, drug rash, back pain, muscle cramps, muscle pain (myalgia), kidney impairment including acute kidney failure, pain in the chest, feeling of weakness, and increased level of creatinine in the blood.

Rare side effects (may affect up to 1 in 1,000 people): Sepsis* (often called "blood poisoning", is a severe infection with whole-body inflammatory response which can lead to death), increase in certain white blood cells (eosinophilia), low platelet count (thrombocytopenia), severe allergic reaction (anaphylactic reaction), allergic reaction (e.g. rash, itching, difficulty breathing, wheezing, swelling of the face or low blood pressure), low blood sugar levels (in diabetic patients), feeling anxious, impaired vision, fast heart beat

(tachycardia), dry mouth, upset stomach, abnormal liver function**, rapid swelling of the skin and mucosa which can also lead to death (angioedema also with fatal outcome), eczema (a skin disorder), redness of skin, hives (urticaria), severe drug rash, joint pain (arthralgia), pain in extremity, tendon pain, flu-like-illness, decreased haemoglobin (a blood protein), increased levels of uric acid, increased hepatic enzymes or creatine phosphokinase in the blood.

*In a long-term study involving more than 20,000 patients, more patients treated with telmisartan experienced sepsis compared with patients who received no telmisartan. The event may have happened by chance or could be related to a mechanism currently not known.

** Most cases of abnormal liver function and liver disorder from post-marketing experience with telmisartan occurred in Japanese patients. Japanese patients are more likely to experience this side effect.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Reporting of side effects:

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF or Vaccines (AEFI)].

Storage and Disposal of CORXEVA

- Storage

Keep out of the reach and sight of children.

Do not use Corxeva after the expiry date which is stated on the carton after "EXP". The expiry date refers

to the last day of that month.

Store below 30°C

Store in the original package in order to protect from moisture.

Corxeva is sensitive to moisture.

Remove your Corxeva tablet from the blister only directly prior to intake.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

Corxeva Tablet 40 mg: White, oblong tablets with LC debossed on one side. Dimensions 12.0 mm x 5.9 mm approximately.

Corxeva Tablet 80 mg: White, oblong tablets with LC debossed on one side. Dimensions 16.0 mm x 8.0 mm approximately.

- Ingredients

• Active Ingredients
Telmisartan

• Inactive Ingredients
Sodium hydroxide
Povidone (K25)
Meglumine
Mannitol (E421)
Magnesium stearate
Crospovidone (type IA)
Purified water

- *MAL Number*

MAL XXXXXXXX AZ

Manufacturer LABORATORIOS LICONSA, S.A.

Avda. Miralcampo, 7, Pol. Ind. Miralcampo, Azuqueca de Henares, 19200 Guadalajara Spain

Product Registration Holder

Duopharma Marketing Sdn. Bhd,

CORXEVA TABLET

Telmisartan (40mg and 80mg)

Lot No. 2,4,6,8,10, Jalan P/7,
Seksyen 13, Kawasan Perusahaan
Bandar Baru Bangi, 43650, Bandar
Baru Bangi, Selangor, Malaysia

Date of revision

May 2026

Serial Number

NPRA (R2) 25/113