

Avorius 5mg Tablet

Desloratadine 5mg

What is in this leaflet

1. What Avorius is used for
2. How it works
3. Before you use Avorius
4. How to use Avorius
5. While you are using it
6. Side effects
7. Storage and Disposal of Avorius
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of Revision

What Avorius is used for

Avorius is used for the rapid relief of symptoms associated with seasonal allergic rhinitis (inflammation of the nasal passages caused by an allergy, for example, hay fever or allergy to dust mites). These symptoms include sneezing, runny or itchy nose, itchy palate, and itchy, red or watery eyes.

Avorius is also used to relieve the symptoms associated with urticaria (a skin condition caused by an allergy). These symptoms include itching and hives.

How it works

Avorius is an antiallergy medicine that does not make you drowsy. It helps control your allergic reaction and its symptoms.

Before you use Avorius

When you must not use it

Do not use Avorius if you are allergic (hypersensitive) to desloratadine, to any of the other ingredients of Avorius or to loratadine.

Before you start to use it

Efficacy and safety of Avorius Tablets in children

under 12 years of age have not been established.

The use of desloratadine is not recommended in breast-feeding women and during pregnancy.

Talk to your doctor, pharmacist or nurse before taking desloratadine if you have medical or familial history of seizures.

Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Avorius

How much to use

Adults and adolescents (12 years of age and over): one tablet once a day.

When to use it

Swallow the tablet whole with water, with or without food.

How long to use it

Continue taking your medicine for as long as your doctor tells you.

Do not stop taking it unless your doctor tells you to - even if you feel better.

If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Avorius.

Tell your doctor immediately if you become pregnant while taking this medication.

Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give Avorius to anyone else, even if they have the same symptoms or condition as you.

Things to be careful of

When driving vehicles or operating machinery it should be taken into account that some people experience drowsiness, which may affect their ability to drive or use machines.

Avorius 5mg Tablet

Desloratadine 5mg

Side effects

Like all medicines, Avorius can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

The most frequent of adverse events reported were fatigue, dry mouth and headache. Other undesirable effects reported during the post-marketing period are listed in the following table:

System Organ Class	Adverse reactions seen with desloratadine
Metabolism and nutrition disorders	
Not known	Increased appetite
Mental disorders	
Very rare	Hallucinations
Not known	Abnormal behaviour, aggression
Nervous system disorders	
Common	Headache
Very rare	Dizziness, drowsiness, sleeping difficulty, restlessness with increased body movement, seizures
Heart disorders	
Very rare	Rapid heartbeat, irregular rapid pumping of the heart
Not known	Irregular heart rhythms
Gastrointestinal disorders	
Common	Dry mouth
Very rare	Abdominal

	pain, nausea, vomiting, indigestion, diarrhoea
Liver disorders	
Very rare	Liver inflammation and abnormal liver function tests
Not known	Jaundice
Skin and subcutaneous tissue disorders	
Not known	Sun allergy
Musculoskeletal and connective tissue disorders	
Very rare	Muscle pain
General disorders	
Common	Tiredness
Very rare	Severe allergic reactions (difficulty in breathing, swelling, itching, rash and hives)
Not known	Lack of energy
Investigations	
Not known	Weight increased

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Avorius

Storage

The tablet should be stored below 30°C, protect from moisture.

Disposal

Medicines should not be disposed of via wastewater

or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

What it looks like

Light blue colored, round shaped, biconvex, film coated tablets with both side plain.

Ingredients

Active ingredient

Desloratadine

Inactive ingredient

Maize starch, Microcrystalline Cellulose, Pregelatinized Starch, Disodium Hydrogen Phosphate Dihydrate, Purified Water, Sodium sulphite, Talc, Opadry Blue 03B50533, Isopropyl alcohol and Dichloromethane.

MAL number

MAL18116128ACZ

Manufacturer

Ind-Swift Laboratories Limited
Off NH-21, Village Jawaharpur, Tehsil Derabassi, District S.A.S. Nagar (Mohali), Punjab - 140 507, India.

Product Registration Holder

Apex Pharmacy Marketing Sdn Bhd
No. 2, Jalan SS 13/5, 47500 Subang Jaya, Selangor, Malaysia

Date of Revision

20 January 2026

Serial Number

NPRA(R2/1)260204/00014