

REPARIL® GEL N

Aescin, Diethylamine salicylate (1g/5g)

What is in the leaflet

1. What Reparil Gel N is used for
2. How Reparil Gel N works
3. Before you use Reparil Gel N
4. How to use Reparil Gel N
5. While you are using it
6. Side effects
7. Storage and Disposal of Reparil Gel N
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Reparil Gel N is used for

Reparil Gel N is used for the treatment of sprains, bruises, crush injuries, and swelling of a tendon (tendonitis). It can be used to treat neck pain and lower back pain (lumbago or sciatica). *Reparil Gel N* can also be used to treat vein swelling or injury and varicose veins.

How Reparil Gel N works

Reparil Gel N contains aescin and diethylamine salicylate which reduce inflammation. Aescin also improves blood flow problems.

Before you use Reparil Gel N

- When you must not use it

Do not use this medicine:

- If you are allergic to salicylates and/or aescin, or any of the ingredients of Reparil Gel N.
- On broken skin, mucous membranes or skin area that is exposed to radiotherapy.
- If you are pregnant or breast feeding.

This medicine is not recommended for children under 12 years.

- Before you start to use it

This product is for external use only. Do not apply to wounds or damaged skin.

- Taking other medicine

For patients taking warfarin, excessive application onto the skin for muscle or joint pains may increase the chances of bleeding.

Please inform your doctor or pharmacist if you are taking any other medicines.

How to use Reparil Gel N

- How much to use

Spread a thin layer of the gel onto the skin of the affected area once daily or more frequently. It is not necessary to rub the gel into the skin.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

Continue using Reparil Gel N for as long as your doctor recommends.

- If you forget to use it

If you forget to use this medicine, use it as soon as you remember. Then continue using it at your usual time.

- If you use too much (overdose)

It is advisable for you to inform your doctor if you applied too much Reparil Gel N.

While you are using it

- Things you must do

Inform your doctor of all the medications you are taking.

- Things you must not do

Do not allow the gel to come into contact with the eyes

- Things to be careful of

Reparil Gel N contains fragrances lavender oil and neroli oil/bitter orange flower oil with d-limonene, farnesol and linalool. These substances may cause allergic reactions.

Seek advice from your doctor or pharmacist if your condition worsens or does not improve.

Side Effects

Like all medicines, Reparil Gel N can cause side effects, although not everybody gets them.

Reparil Gel N might cause skin irritation, redness or swelling, burning or stinging sensation.

Dizziness and nausea might occur in rare cases.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Reparil Gel N

- Storage

Store below 30°C. Protect from light.

Keep medicines out of reach of children.

- Disposal

Medicine should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

REPARIL® GEL N

Aescin, Diethylamine salicylate (1g/5g)

Product Description

- What it looks like

A transparent, colourless yellowish gel with an aromatic odour.

- Ingredients

- Active ingredients

Aescin, Diethylamine salicylate

- Inactive ingredients

Disodium Edetate, Lavender Oil, 2Propanol, Macrogol 6 Glycerol Caprylocaprate, Trometamol, Carbomers, Citrus Aurantium Amara (Bitter Orange) Oil (Flower), Purified Water

- MAL number

MAL06030857XR

Manufacturer

Madaus GmbH
Lütticher Straße 5,
53842 Troisdorf, Germany

Product Registration Holder

Mylan Healthcare Sdn. Bhd.
15-03 & 15-04, Level 15,
Imazium, No. 8, Jalan SS 21/37,
Damansara Uptown, 47400
Petaling Jaya, Selangor,
Malaysia.

Date of Revision

10/04/2026

Serial Number

NPRA(R2/1)230207/1007