

Reparil® Gel N

1. What Reparil Gel N is and what it is used for

100g Reparil Gel N contains:

Aescin	1g
Diethylamine salicylate	5g

Reparil Gel N is used for the following:

- In cases of contusions, crush injuries, sprains, bruises, haematoma and tendonitis.
- Painful conditions of the vertebral column (disc lesions, neck pain, lumbago, sciatica).
- In superficial thrombophlebitis, varicose veins. For care of the veins after injections or infusions.

2. What you need to know before you use Reparil Gel N

Do not use Reparil Gel N:

- If you are allergic to the active substances or to any of the ingredients of Reparil Gel N.
- To broken skin, mucous membranes or skin areas exposed to radiotherapy.
- If you are pregnant or breast feeding.

Warnings and Precautions

- As a preparation for topical application, its use must be for external use only. Do not bandage tightly.
- The contact with the eyes must be avoided.
- Patients hypersensitive to salicylates are at a greater risk of developing asthma attacks, local skin and mucosal swelling (Quincke's oedema) or urticaria than other patients. In these patients, diethylamine salicylate/aescin must be used with caution and only under direct medical monitoring.
- Reparil Gel N contains diethylamine salicylate and when applied or rubbed onto the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application onto the skin for muscle or joint pains may increase the chances of bleeding.
- This medicine contains fragrances lavender oil and neroli oil/bitter orange flower oil with d-limonene, farnesol and linalool. These substances may cause allergic reactions.
- If condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days, discontinue use of this product and consult a doctor.

Special populations

The use in children under the age of 12 years is not recommended.

No dose adjustment is necessary in the elderly population.

No dose reduction is required in patients with impaired renal or liver function.

Using Reparil Gel N with other medicines

For patients taking warfarin, excessive application onto the skin for muscle or joint pains may increase the chances of bleeding.

3. How to use Reparil Gel N

Unless otherwise directed, spread a thin layer of the gel onto the skin of the affected area once daily or more frequently. It is not necessary to rub the gel into the skin, but this may be done if desired.

4. Possible side effects

Skin irritation, redness or swelling, burning or stinging sensation. Adverse reaction is rare, symptom of dizziness, nausea may occur.

5. How to store Reparil Gel N

Store below 30°C in a dry place.

Keep medicines out of reach of children.

6. Further information

Properties

- Reparil Gel N is a special formulation in an alcohol base with a threefold action: anti-oedematous, anti-inflammatory and pain-relieving.
- Aescin, in a form which penetrates the skin, inhibits the accumulation of fluid in the tissues and reduces tissue swelling. Local swelling, which causes sensations of heaviness and tension, is rapidly eliminated. It effectively combats inflammatory processes.
- The salicylic acid compound enhances the anti-inflammatory action and is extremely effective for relieving deep-seated pain.
- In addition, Reparil Gel N has the following advantages:
 - Reparil Gel N is transparent and penetrates rapidly and completely into the affected tissues.
 - Reparil Gel N does not make the skin greasy, does not smear and is therefore particularly clean and pleasant to use.

- Reparil Gel N cools the painful area, is kind to the skin and is outstandingly well tolerated even on sensitive parts of the skin.

What Reparil Gel N looks like and contents of the pack

Reparil Gel N is a transparent, colourless to yellowish gel with an aromatic odour.

Packs of 10g, 20g, and 40g. Not all pack sizes may be marketed.

Manufacturer

Madaus GmbH,
Lütticher Straße 5,
53842 Troisdorf,
Germany

This leaflet was last revised in Mar 2026.