

# REPARIL® GEL N

Aescin, Diethylamine salicylate (1g/5g)

What is in the leaflet

1. What *Reparil® Gel N* is used for
2. How *Reparil® Gel N* works
3. Before you use *Reparil® Gel N*
4. How to use *Reparil® Gel N*
5. While you are using it
6. Side effects
7. Storage and Disposal of *Reparil® Gel N*
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

## What *Reparil® Gel N* is used for

*Reparil Gel N* is used for the treatment of sprains, bruises, crush injuries, and swelling of a tendon (tendonitis). It can be used to treat neck pain and lower back pain (lumbago or sciatica). *Reparil Gel N* can also be used to treat vein swelling or injury and varicose veins.

## How *Reparil® Gel N* works

*Reparil-Gel N* contains aescin and diethylamine salicylate which reduce inflammation. Aescin also improves blood flow problems.

## Before you use *Reparil® Gel N*

- When you must not use it  
*Reparil® Gel N* should not be applied on broken skin, mucous membranes or skin area that is exposed to radiotherapy.

- Before you start to use it  
This product is for external use only. Do not apply to wounds or damaged skin.

- Taking other medicine  
Please inform your doctor or pharmacist if you are taking any other medicines.

## How to use *Reparil® Gel N*

- How much to use  
Spread a thin layer of the gel onto the skin of the affected area once daily or more frequently. It is not

necessary to rub the gel into the skin.

### When to use it

Please follow the advice of the doctor or pharmacist.

### - How long to use it

Continue using *Reparil® Gel N* for as long as your doctor recommends.

### - If you forget to use it

If you forget to use this medicine, use it as soon as you remember. Then continue using it at your usual time.

### - If you use too much (overdose)

It is advisable for you to inform your doctor if you applied too much *Reparil Gel N*.

## While you are using it

- Things you must do  
Inform your doctor of all the medications you are taking.

### - Things you must not do

Do not allow the gel to come into contact with the eyes.

### - Things to be careful of

Seek advice from your doctor or pharmacist if your condition worsens or does not improve.

## Side Effects

Like all medicines, *Reparil® Gel N* can cause side effects, although not everybody gets them.

*Reparil® Gel N* might cause skin irritation, redness or swelling, burning or stinging sensation.

Dizziness and nausea might occur in rare cases.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npdra.gov.my](http://npdra.gov.my) [Consumers→

Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of *Reparil® Gel N*

### - Storage

Store below 30°C. Protect from light.

### - Disposal

Medicine should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required.

## Product Description

### - What it looks like

A transparent, colourless to yellowish gel with an aromatic odour.

### - Ingredients

- Active ingredients  
Aescin, Diethylamine salicylate

- Inactive ingredients  
Disodium Edetate, Lavender Oil, 2-Propanol, Macrogol 6 Glycerol Caprylocaprate, Trometamol, Carbomers, Citrus Aurantium Amara (Bitter Orange) Oil (Flower), Purified Water

### - MAL number

MAL06030857X

## Manufacturer

MADAUS GmbH  
51101 Cologne, 51101, GERMANY

## Product Registration Holder

MYLAN HEALTHCARE SDN. BHD.  
15-03 & 15-04, Level 15, Imazium, No. 8, Jalan SS 21/37, Damansara Uptown, 47400 Petaling Jaya, Selangor, Malaysia.

## Date of Revision

- 04/01/2023

## Serial Number

NPRA(R2/1)230207/1007