

# REPARIL® TABLETS

Amorphous Aescin (20mg)

## What is in this leaflet

1. What is Reparil® Tablet is used for
2. How Reparil® Tablet works
3. Before you use Reparil® Tablet
4. How to use Reparil® Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of Reparil® Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

## What is Reparil® Tablet is used for

Reparil® Tablet is used for reduction of localized swelling following traumatic injuries.

## How Reparil® Tablet works

Reparil® Tablet contains aescin which reduces inflammation and improves blood flow to speed up recovery of localized swelling.

## Before you use Reparil® Tablet

### - When you must not use it

Do not take Reparil® Tablet if you have:

- hypersensitivity to aescin or any other ingredients of this medicine (listed in Product Description)
- kidney problems

### - Before you start use it

Reparil® Tablet are not indicated for children under 7 years of age.

### *Pregnancy and lactation*

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

Tell your doctor or pharmacist if you have any rare hereditary sugar malabsorption as Reparil® Tablet contains of lactose and sucrose and you should not take it.

### - Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Tell your doctor or pharmacist if you are taking the following medicines:

- Anticoagulants (blood thinning medicine)
- Antibiotics like gentamicin, cephalotin and ampicillin

## How to use Reparil® Tablet

### - How much to use

Always use Reparil® Tablet exactly as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure.

The dosage are as follows:

- Adults and adolescents over 14 years (initial dose): 2 tablets 3 times daily
- In milder cases or maintenance dose: 1 tablet 3 times daily
- Children between 7-14 years of age: 1 tablet taken 2 or 3 times daily.

### - When to use it

Use as directed by your doctor or pharmacist.

Take Reparil® Tablet after meals with some fluid.

### - How long to use it

Reparil® Tablet treatment should be continued for two to three months or for the duration that your doctor recommends.

### - If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose.

### - If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many tablets may cause excitation, anxiety, diarrhea, vomiting and others

## While you are using it

### - Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentist and pharmacists treating you that you are taking Reparil® Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

### - Things you must not do

Do not skip taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Reparil® Tablet to anyone else, even if they have the same symptoms or condition as you.

# REPARIL® TABLETS

Amorphous Aescin (20mg)

## Things to be careful of

### *Driving and using machines*

Be careful driving or operating machinery until you know how Reparil® Tablet affects you.

## Side effects

Like all medicines, Reparil® Tablet can cause side effects, although not everybody gets them.

In very rare cases, hypersensitivity reactions like hives can occur. Stop taking Reparil® Tablet if this happens.

Disorders of the gastrointestinal tract are uncommon.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers -> Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

## Storage and Disposal of Reparil® Tablet

### - Storage

Keep out of the reach and sight of children.

The expiry date is printed on the folding carton. Do not use this pack after the expiry date has passed.

Avoid storing product in direct sunlight.

Store below 30°C.

### - Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### - What it looks like

Reparil® Tablet are round, biconvex white, glossy coated-tablets of 7mm in diameter.

### - Ingredients

- Active ingredient:  
Amorphous aescin. Aescin is derived from horse chestnut (Aesculus hippocastanum).

- Inactive ingredient:  
Simethicone emulsion, white beeswax, sodium hydroxide, carmellose sodium, anhydrous colloidal silica, triethyl citrate, macrogol 8000, titanium dioxide, acacia, povidone, magnesium stearate, talc, sucrose, methacrylic acid-ethyl acrylate copolymer, lactose monohydrate, carnauba wax

- MAL number:  
MAL11030107XR

## Manufacturer

Madaus GmbH,  
Lütticher Straße 5,  
53842 Troisdorf,  
Germany.

## Product Registration Holder

Mylan Healthcare Sdn. Bhd. ,  
15-03 & 15-04, Level 15,  
Imazium,  
No. 8, Jalan SS 21/37,  
Damansara Uptown,  
47400 Petaling Jaya, Selangor,  
Malaysia.

## Date of revision

28/02/2025

## Serial Number

NPRA(R2/1)250310/1429