

VIARTRIL[®]-S 750mg FILM-COATED TABLETS

Glucosamine sulphate (750 mg)

What is in this leaflet

1. What Viartril-S 750mg Film-Coated Tablet is used for
2. How Viartril-S 750mg Film-Coated Tablet works
3. Before you use Viartril-S 750mg Film-Coated Tablet
4. How to use Viartril-S 750mg Film-Coated Tablet
5. While you are using Viartril-S 750mg Film-Coated Tablet
6. Side Effects
7. Storage and disposal of Viartril-S 750mg Film-Coated Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision
11. Serial Number

What Viartril-S 750mg Film-Coated Tablet is used for

Viartril-S 750mg Film-Coated Tablet is used as an adjuvant therapy for osteoarthritis (loss of cartilage in the joints).

How Viartril-S 750mg Film-Coated Tablet works

Viartril-S 750mg Film-Coated Tablet contains glucosamine sulphate as active ingredient. Glucosamine sulphate provides relief to joint disease by reducing joint swelling and pain.

Before you use Viartril-S 750mg Film-Coated Tablet

- When you must not use it

Do not take Viartril-S 750mg Film-Coated Tablet if you have:

- Hypersensitivity to glucosamine or any other ingredients of this medicine (listed in Product Description).
- Allergies to shellfish as the active ingredient is obtained from seafood (shellfish).

Pregnancy and lactation

Do not take Viartril-S 750mg Film-Coated Tablet if you are pregnant,

trying to get pregnant or think you may be pregnant.

Do not take Viartril-S 750mg Film-Coated Tablet if you are breast-feeding.

- Before you start to use it

Tell your doctor or pharmacist if you have:

- Asthma
- A controlled sodium diet
- Liver or kidney problems
- Diabetes or impaired glucose tolerance
- Another joint disease which may require alternative treatments

- Taking other medicines

Tell your doctor or pharmacist if you are taking:

- Coumarinic anticoagulants (blood thinner)
- Tetracyclines (antibiotics)

Pain relief or anti-inflammatory medicines can be taken together with Viartril-S 750mg Film-Coated Tablet.

Tell your doctor or pharmacist if you are taking any other medicines, including any that buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Viartril-S 750mg Film-Coated Tablet

- How much to use

Take one tablet.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

To be taken 2 times daily preferably

at meals or as directed by your doctor or pharmacist.

- How long to use it

Take Viartril-S 750mg Film-Coated Tablet for the duration that your doctor or pharmacist recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Viartril-S 750mg Film-Coated Tablet

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Viartril-S 750mg Film-Coated Tablet.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not give this medicine to children and adolescent under the age of 18 as the safety and efficacy have not been

VIARTRIL[®]-S 750mg FILM-COATED TABLETS

Glucosamine sulphate (750 mg)

established.

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give Viartril-S 750mg Film-Coated Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablet causes drowsiness, headache and visual disturbance, do not drive or use machines and contact your doctor immediately.

Side Effects

Like all medicines, Viartril-S 750mg Film-Coated Tablet can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

The side effects may include:

- *Heart and blood vessels:* swelling of limbs, fast heart rate
- *Central nervous system:* drowsiness, headache, sleepiness, difficulty in sleeping
- *Gastrointestinal:* nausea, vomiting, diarrhea, indigestion or pain in upper abdomen, passing gas, constipation, heartburn and eating disorder
- *Skin reactions:* rash, redness, itchiness, inflammation, hives, eczema (inflamed skin)
- *Others:* hair loss, allergic reactions, visual disturbances, swelling

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.moh.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Viartril-S 750mg Film-Coated Tablet

- Storage

Store below 30°C.
Keep out of the reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste.

Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Unmarked oblong white film-coated tablets without a score line.

- Ingredients

Each tablet contains:

- *Active ingredient*
Crystalline glucosamine sulphate 942 mg (equivalent to 750 mg glucosamine sulfate and 192 mg sodium chloride)
- *Inactive ingredients*
Microcrystalline cellulose, Povidone K 25, Croscarmellose sodium, Macrogol 6000, Magnesium stearate, Talc, Methacrylic acid - methyl methacrylate copolymer (1:1), Titanium dioxide, Ammonium methacrylate copolymer (type A), Triacetin

- MAL number:

MAL24096079X

Manufacturer

Madaus GmbH
Lütticher Straße 5
53842 Troisdorf
Germany

Product Registration Holder

Mylan Healthcare Sdn Bhd
15-03 & 15-04, Level 15, Imazium,
No. 8, Jalan SS 21/37, Damansara
Uptown,
47400 Petaling Jaya,
Selangor, Malaysia

Date of Revision

16/06/2025

Serial Number

NPRA (R2) 24/068

PIL-VIARTRILS TAB-0625