

'MPI' PYRIDOXINE TABLETS 10MG

Pyridoxine Hydrochloride (10mg)

What is in this leaflet:

1. What 'MPI' Pyridoxine Tablets 10mg is used for
2. How 'MPI' Pyridoxine Tablets 10mg works
3. Before you use 'MPI' Pyridoxine Tablets 10mg
4. How to use 'MPI' Pyridoxine Tablets 10mg
5. While you are using it
6. Side Effects
7. Storage and Disposal of 'MPI' Pyridoxine Tablets 10mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What 'MPI' Pyridoxine Tablets 10mg is used for

It is indicated as a health supplement.

How 'MPI' Pyridoxine Tablets 10mg works

Pyridoxine hydrochloride helps in protein, carbohydrate and fat metabolism.

Before you use 'MPI' Pyridoxine Tablets 10mg

- When you must not use it

- If you are allergic to pyridoxine hydrochloride or any of the other ingredients of 'MPI' Pyridoxine Tablets 10mg.

- Before you start to use it

You must tell your doctor if:

- you have allergies to any medicine which you have taken previously to treat your current condition.
- you are pregnant, trying to get pregnant, think you may be pregnant or breast-feeding.

- Taking other medicines

Tell your doctor if you are taking Levodopa, a medicine used in the treatment of Parkinson's disease.

Always inform your doctor and pharmacist if you are taking any other medicines, including any that you buy without prescription from a

pharmacy, supermarket or health food shop.

How to use 'MPI' Pyridoxine Tablets 10mg

'MPI' Pyridoxine Tablets 10mg should be swallowed whole, not chewed.

- How much to use

Usual adult dose: 1-2 tablets per day.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Take for 3 weeks followed by a multivitamin preparation or as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention. Taking large doses (more than 2g daily) of 'MPI' Pyridoxine Tablets 10mg for a long term (2 months or longer) has been associated with problems with the nerves, progressing from unstable gait and numb feet to numbness and clumsiness of hands.

While you are using it

- Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are

taking 'MPI' Pyridoxine Tablets 10mg.

Tell your doctor immediately if you become pregnant while taking this medication.

Contact your doctor if your symptoms worsen or they do not improve.

- Things you must not do

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give 'MPI' Pyridoxine Tablets 10mg to anyone else, even if they have same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side Effects

Like all medicines, 'MPI' Pyridoxine Tablets 10mg can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my

[Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccine (AEFI)]

Storage and Disposal of 'MPI' Pyridoxine Tablets 10mg

- Storage

Do not use after the expiry date stated on the pack. The expiry date refers to the last day of that month.

Keep in original packaging or amber container. Store below 30°C. Protect from light, heat and moisture.

Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A white, flat of diameter 6mm round tablet with 'MPI' logo.

- Ingredients

- Active ingredient:

Pyridoxine hydrochloride.

- Inactive ingredients:

Lactose, corn starch, colloidal silicon dioxide and magnesium stearate.

- MAL number:

MAL20034563XZ

Manufacturer and Product Registration Holder

Malaysian Pharmaceutical Industries Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas, Penang, Malaysia.

Date of Revision

25/08/2025

Serial Number

NPRA(R4/1)280219/00062