

MUCO-AID LIQUID

Lignocaine Hydrochloride/ Triamcinolone Acetonide/ Chlorhexidine Hydrochloride (2.5%w/v / 0.1%w/v / 0.5%w/v)

What is in this leaflet:

1. What Muco-Aid Liquid is used for
2. How Muco-Aid Liquid works
3. Before you use Muco-Aid Liquid
4. How to use Muco-Aid Liquid
5. While you are using it
6. Side effects
7. Storage and Disposal of Muco-Aid Liquid
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of Revision

What Muco-Aid Liquid is used for
Muco-Aid Liquid is indicated for the relief of discomfort and pain, for the reduction of inflammation and to promote healing in the following conditions:

- ulcer, lesions and soreness in the gums, tongue, lips or palate;
- trauma or lesions in the oral cavity caused by new denture, denture sore spots and denture inflammation.
- teething disorders.

How Muco-Aid Liquid works

Muco-Aid contains three types of active ingredients called lignocaine hydrochloride, chlorhexidine hydrochloride, and triamcinolone acetonide.

Lignocaine hydrochloride is a local anaesthetic agent which relieves pain. Chlorhexidine hydrochloride is an antimicrobial agent which inhibit the bacterial infection. Triamcinolone acetonide is an anti-inflammatory agent which helps to reduce swelling.

Before you use Muco-Aid Liquid

- When you must not use it:

Do not use this medicine if you are allergic (hypersensitive) to:

- Lignocaine hydrochloride
- Chlorhexidine hydrochloride
- Triamcinolone acetonide

- Any of the other ingredients of Muco-Aid Liquid.

Immediately stop using the product if you have any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate.

Do not use this medicine if you have lesions, blister and in the presence of infections in your mouth.

- Before you start to use it:

You must tell your doctor if:

- you have allergies to any medicine which you have used previously to treat your current condition.
- you have tuberculosis, gastric ulcer, or diabetes.
- you are pregnant especially during early pregnancy or nursing mothers.

Use with caution in children especially when extensive areas are treated since growth retardation has been reported.

- Taking other medicines:

Muco-Aid Liquid may interact with:

- Beta- blockers (for heart problems)
- Cimetidine (for heartburn and stomach ulcers)
- Non-steroidal Anti Inflammatory Drugs (NSAIDs) (pain killer).
- Thiazide, furosemide (water pills)

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use Muco-Aid Liquid

Always use this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

- How much to use:

Instill 2-4 drops of Muco-Aid Liquid to the lesion by using the tip of a clean cotton bud or finger. Rub gently to cover the lesion.

- When to use it:

Application may be repeated after 1-2 hours, preferably after meal, and at bedtime.

Use as directed by your doctor or pharmacist.

- How long to use it:

Use Muco-Aid Liquid for the duration that your doctor recommends.

- If you forget to use it:

If you forget to apply the medicine, you can apply it as soon as you remember.

- If you use too much (overdose):

Signs and symptoms of overdose may include:

- blurred or double vision, or both;
- dizziness;
- ringing sound in the ear;
- shivering or trembling;
- increased sweating;
- low blood pressure;
- slow or irregular heartbeat;
- unusual paleness,
- unusual anxiety, excitement, nervousness, restlessness and convulsion,
- drowsiness, not conscious, breathing difficulties and death.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have used too much of this medicine. Do this even if there are no signs of discomfort or

poisoning. You may need urgent medical attention.

While you are using it

- Things you must do:

Always follow your doctor's instruction carefully.

Tell your doctor if you are pregnant or breast-feeding.

- Things you must not do:

- Do not give Muco-Aid Liquid to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed Muco-Aid Liquid for you and your condition.

- Muco-Aid Liquid contains chlorhexidine and cannot be instilled into ear.

- Do not use Muco-Aid Liquid to treat any other complaints unless your doctor tells you to.

- Do not chew gum or food while numbness persists because of risk of biting your tongue.

- Do not eat for one hour following use of medication because it may impair swallowing leading to risk of aspiration.

- Things to be careful of:

- Be cautious when driving a car or operating machinery until you know how Muco-Aid Liquid affects you.

- Consult your doctor if there is no improvement after 7 days.

Side effects

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the following side effects:

- pain
- redness
- burning
- itching
- blisters and it can be filled with pus
- peeling

- Cushing's Syndrome (disorder caused by excessive cortisol hormone in body)
- High level of sugar in blood
- Presence of sugar in urine

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Muco-Aid Liquid

- Storage:

Do not use after the expiry date stated on the pack. The expiry date refers to the last day of that month.

Store below 30°C in well-closed containers. Protect from light, heat and freezing.

Keep out of the reach and sight of children.

- Disposal:

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like:

A white lotion with bubblegum and peppermint flavour.

- Ingredients:

- Active ingredient(s)
Lignocaine hydrochloride, Triamcinolone acetonide and Chlorhexidine hydrochloride
- Inactive ingredients
The other ingredients are Glycerin, Polysorbate 80, Xanthan Gum, Bubblegum Flavour and Peppermint Oil.

- MAL number

MAL05041940AZ

Manufacturer and Product Registration Holder

Malaysian Pharmaceutical Industries Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas, Penang, Malaysia.

Date of Revision

15/11/2021

Serial Number

NPRA(R4/1)180621/00052