

BISCOMIN TABLETS 8MG

Bromhexine HCl 8mg

What is in this leaflet

1. What Biscomin Tablets 8mg is used for
2. How Biscomin Tablets 8mg works
3. Before you use Biscomin Tablets 8mg
4. How to use Biscomin Tablets 8mg
5. While you are using Biscomin Tablets 8mg
6. Side effects
7. Storage and Disposal of Biscomin Tablets 8mg
8. Product description
9. Manufacturer and Product Registration holder
10. Date of Revision

What Biscomin Tablets 8mg is used for

Bromhexine HCl is the active ingredient of Biscomin Tablets 8mg. It is a mucolytic agent and is indicated in the treatment of respiratory disorders associated with viscid or excessive mucus.

How Biscomin Tablets 8mg works

Bromhexine alters the structure of mucus to decrease its viscosity and therefore facilitate its removal by ciliary action or expectoration.

Before you use Biscomin Tablets 8mg

- When you must not use it

- If you have known hypersensitivity to Bromhexine or any of its ingredients

- Before you start to use it

You must tell your doctor if:

- You have allergies to any medicine which you have taken previously to treat your current condition.

- Taking other medicines

Always inform your doctor and pharmacist if you are taking any other medicines, including herbal tonics, supplements and medicines that you buy without prescription.

How to use Biscomin Tablets 8mg

Always take this medicine exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

- How much to use

Adults: 8-16mg (1-2 tablets) three times a day

Children 6-12 years old: 4mg (half tablet) three times a day

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Biscomin Tablets 8mg for as long as your doctor recommends.

- If you forget to use it

- If you forget a dose, take it as soon as you remember it.
- However, if it is nearly time for the next dose, skip the missed dose.
- Do not take a double dose to make up for a forgotten dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using Biscomin Tablets 8mg

- Things you must do

Contact your doctor if your symptoms does not improve. Take the medicine after food.

- Things you must not do

Do not give Biscomin Tablets 8mg to anyone else, even if their symptoms seem similar or they have same condition as you. Your doctor has prescribed Biscomin Tablets 8mg for you and your condition.

- Things to be careful of

Special care is recommended during pregnancy. The benefits must be assessed against possible effects on the child.

Side effects

Like all medicines, Biscomin Tablets 8mg can have side effects, although not everybody gets them. The following side effects may happen with this medicine:

Gastro-intestinal side effects may occur occasionally (nausea, vomiting and diarrhoea) and skin rash.

Biscomin Tablets 8mg may cause severe allergy and serious skin reactions. Stop using Biscomin Tablets 8mg and seek medical assistance immediately if you experience any of the following symptoms:

- 1) Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- 2) Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

Consult your doctor or pharmacist if you are in doubt or for any further information.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website npra.moh.gov.my (Public→Reporting Medicinal

Problems/ Side Effects/ AEFI/
Vaccine Safety)

**Storage and Disposal of Biscomin
Tablets 8mg**

- Storage

Do not use after the expiry date stated on the pack. The expiry date refers to the last day of that month.

Store below 30°C. Protect from heat and moisture.

Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

An orange, scored, flat of diameter 7mm round tablet with 'MPI' logo.

- Ingredients

- Active ingredient – Bromhexine HCl
- Inactive ingredient - The other ingredients are Potato Starch, Lactose, Corn Starch, Sunset Yellow FCF, Magnesium Stearate.

- MAL number:

MAL19920348XZ

**Manufacturer and Product
Registration Holder**

Malaysian Pharmaceutical Industries
Sdn. Bhd.

Plot 14, Lebuhraya Kampung Jawa,
11900 Bayan Lepas, Penang,
Malaysia.

Date of Revision

26/06/2018