

U-IODIN LOTION 10%W/V

A deep brown liquid with a characteristic odor.

COMPOSITION:

Povidone-Iodine 10% w/v.

INDICATION:

As a skin disinfectant.

ROUTE OF ADMINISTRATION:

To apply topically.

RECOMMENDED DOSAGE:

For External Use Only.

“U-Iodin Lotion” is applied topically by dabbing gently to the affected areas or after spreading on a dressing, when necessary.

CONTRAINDICATIONS:

It should not be used on patients who are hypersensitive to iodides and on patients with non-toxic nodular colloid goiter.

WARNINGS AND PRECAUTIONS:

Application to large areas of broken skin should be avoided as excessive absorption of iodine may occur. Absorption of iodine may interfere tests of thyroid function.

ADVERSE EFFECTS / UNDESIRABLE EFFECTS:

Local irritation and sensitivity may occur. Discontinue treatment if local irritation occurs.

OVERDOSE AND TREATMENT:

Symptoms : Irritation of the skin not present before therapy.

Treatment : Discontinue treatment if local irritation occurs.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Plastic bottle : 500 ml, 1L

NAME AND ADDRESS OF MANUFACTURER / PRODUCT REGISTRATION HOLDER:

MALAYSIAN PHARMACEUTICAL INDUSTRIES SDN BHD (101323-U)

Plot 14, Lebuhraya Kampung Jawa, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

DATE OF REVISION:

[day/month/year]

Date of Revision will be based on BPFK approval date