

SERRIN TABLET

Consumer Medication Information Leaflet (RiMUP)

Serratiopeptidase (5 mg)

What is in this leaflet

1. What SERRIN Tablet is used for
2. How SERRIN Tablet works
3. Before you use SERRIN Tablet
4. How to use SERRIN Tablet
5. While you are using it
6. Side effects
7. Storage and Disposal of SERRIN Tablet
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What is SERRIN Tablet used for

SERRIN Tablet is used to relief of inflammation in the following diseases:

- After surgery and trauma.
- Engorgement of breast.
- Inflammation of the
 - Bladder.
 - Tube that carries sperm from the testes.
 - Gum tissue of the wisdom tooth.
 - Nasal sinus.
- Tooth abscess (pus).

It is also used in the treatment of difficulty of expectoration in:

- Bronchitis (inflammation of the airways that carry air to lungs).
- Pulmonary tuberculosis (infection of the lungs).
- Asthma.
- Disease after anesthesia

How SERRIN Tablet works

Serratiopeptidase is an enzyme which helping the body in breakdown of protein. It has the ability to reduce pain by blocking the release of pain inducing chemicals from inflamed tissues.

Before you use SERRIN Tablet

-When you must not use it

Do not take SERRIN Tablet if:

- You are allergy (hypersensitive) to serratiopeptidase or any of the other ingredients in the product.

- You are pregnant, trying to get pregnant or think you may be pregnant.
- If you are breast-feeding, please ask your doctor or pharmacist for advice before taking any medicine.

-Before you start to use it

Tell your doctor if:

- You are having history of medicine allergy and blood abnormalities such as bleeding from the nose and bloody mucus.
- You are having severe liver or kidney dysfunction.

-Taking other medications

Tell your doctor if you are taking the following medicine as it may interact with SERRIN Tablet:

- Anticoagulant (for blood thinning).

It may still be all right for you to take SERRIN Tablet and your doctor will be able to decide what is suitable for you.

How to use SERRIN Tablet

-How much to use

For oral administration only. To be taken 1-2 tablets three times a day after meals without chewing. Dosage may be adjusted according to the individual age and condition.

-When to use it

Take SERRIN Tablet three times daily after meal or as directed by physician. Your doctor will decide when it is appropriate for you to use SERRIN Tablet according to the type of condition you have.

-How long to use it

Continue taking SERRIN Tablet for as long as doctor recommends.

-If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

-If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking SERRIN Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

-Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give SERRIN Tablet to anyone else, even if they have the same symptoms or condition as you.

-Things to be careful of

- No interference with the capability of driving or operating heavy machinery.

Side effects

Like all medicines, SERRIN Tablet can cause side effects although not everybody gets them. You are advised to consult doctor/ pharmacist for further information. The following occasional side effects have been reported:

- Rash or redness.
- Diarrhoea.
- Loss of appetite.
- Gastric discomfort.

SERRIN TABLET

Consumer Medication Information Leaflet (RiMUP)

Serratiopeptidase (5 mg)

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of SERRIN Tablet

-Storage

Keep SERRIN Tablet out of reach of children. Store below 30°C. Keep container well closed. Protect from light and moisture.

-Disposal

Do not use SERRIN Tablet after the expiry date which is stated on the carton. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

SERRIN Tablet is orange, convex round enteric film-coated tablets with letters "AM 005" in black colour on one side.

-Ingredients

- Active ingredient(s)
Serratiopeptidase
- Inactive ingredient(s)
Lactose Monohydrate, Corn starch, Polyethylene Glycol 6000, Magnesium Stearate, Talcum, Eudragit L 100, Sunset yellow dye FCF, Sunset Yellow lake FCF, Purified water, Isopropyl alcohol, Acetone

-MAL number

MAL19991391X

Manufacturer

MacroPhar Co., Ltd.
89 Soi Pattanakarn 20 Yaek 4,
Pattanakarn Road, Suan Luang,
Suan Luang, Bangkok 10250

Thailand. Tel.: (662)314-6671

Fax: (662)318-6091

Product Registration Holder

Healol Pharmaceuticals Sdn. Bhd.
74-3, Jalan Wangsa Delima 6, KLSC
Wangsa Maju, 53300 Kuala Lumpur,
Malaysia

Tel.: 03-412 7106, 412 7107

Fax: 03-4149 8108

Date of revision

22/02/2021

Serial Number

NPRA(R2/1)210219/628