

12.7 cm

CLOMAZ[®]

(CLOTRIMAZOLE 1% CREAM)

LF-0189

COMPOSITION: Each 10 g of **CLOMAZ[®] (CLOTRIMAZOLE 1% CREAM)** contains Clotrimazole 0.1 g**PROPERTIES & ACTIONS:** Clotrimazole inhibits or kills many fungi, including yeasts and dermatophytes. Clotrimazole exhibit fungistatic and fungicidal activity against isolates of *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Epidermophyton floccosum*, *Microsporium canis*, and *Candida* species, including *Candida albicans*.**PHARMACOLOGY / PHARMACOKINETICS:** Clotrimazole is a broad-spectrum antifungal agent that is used for the treatment of dermal infections caused by various species of pathogenic dermatophytes, yeasts, and *Malassezia furfur*. The primary action of Clotrimazole is against dividing and growing organisms. Only very small amounts of clotrimazole appear to be absorbed systemically following topical application to the skin. Following application to the skin, highest concentrations of clotrimazole are present in the stratum spinosum and the papillary and reticular dermis.**INDICATIONS:** **CLOMAZ[®] (CLOTRIMAZOLE 1% CREAM)** is indicated for dermatomycoses caused by dermatophytes, yeasts, moulds, eg., tinea pedum, tinea manuum, tinea corporis, tinea inguinalis pityriasis versicolor, erythrasma, *Candida balanitis* and *Candida vulvitis*.**DOSAGE & APPLICATION:** Gently massage sufficient **CLOMAZ[®] (CLOTRIMAZOLE 1% CREAM)** into the affected surrounding skin areas 2-3 times daily. A small amount of cream is usually sufficient for an area about the size of the palm. In fungal infection of the feet, after washing the feet should be thoroughly dried (spaces between toes).**General duration of treatment:** Dermatomycoses : 3-4 weeks

Candida balanitis and Candida vulvitis : 1-2 weeks


Erythrasma and pityriasis versicolor : about 3 weeks

To prevent relapses, treatment should be continued for approximately 2 weeks beyond the disappearance of all signs of disease. For the treatment to become a full success, reliable and sufficiently long application of clotrimazole cream is important.

PRECAUTIONS / WARNINGS:**General:** If irritation occurs or if the patient's dermatologic condition does not improve within 2 weeks for tinea cruris or within 4 weeks for tinea pedis or tinea corporis during self-medication with clotrimazole cream or solution, treatment should be discontinued and the patient should consult a physician or pharmacist. Clotrimazole topical cream is not intended for ophthalmic use.**Pediatric:** For self-medication, clotrimazole topical cream should not be used in children younger than 2 years of age unless otherwise instructed by a physician.**Pregnancy and lactation:** Clotrimazole topical cream should be used during the first trimester of pregnancy only when the drug is considered essential to the welfare of the patient. Since it is not known whether clotrimazole is distributed into milk, the drug should be used with caution in nursing women.**Drug Interactions:** Synergism or antagonism between clotrimazole and nystatin, or amphotericin B, or flucytosine against strains of *C. albicans* has not been reported.**SIDE EFFECTS / ADVERSE REACTIONS:** Although clotrimazole is usually well tolerated when administered topically to the skin, blistering, erythema, edema, pruritus, burning, stinging, peeling, urticaria, skin fissures, and general irritation of the skin have occasionally occurred. If irritation or sensitization occurs following topical application of clotrimazole, the drug should be discontinued.**CONTRAINDICATION:**

Clotrimazole products are contraindicated individual who have shown hypersensitivity to any of their component.

OVERDOSAGE: Acute overdosage with topical application of clotrimazole is unlikely and would not be expected to lead to a life-threatening situation.**STORAGE CONDITIONS:** Store below 30°C**SHELF-LIFE:** 4 years**DESCRIPTION OF PRODUCT:** Homogeneous, white cream.**PACKING SIZE:** Aluminium tube 15 g.

Manufactured by:
 **L.B.S. LABORATORY LTD., PART.**
 602 Soi Panichanant, Sukhumvit 71 Rd., Bangkok 10110 Thailand.
 Tel. (66) 2392-4082 Fax. (66) 2391-7122

Product registration holder:
HEALOL PHARMACEUTICALS SDN. BHD.
 74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju,
 53300 Kuala Lumpur, Malaysia
 Tel : (60) 3-4142-7106, 4142-7107 Fax : (60) 3-4149-7108

18.8 cm

Code: R 347/24
 Project: CLOMAZ (CLOTRIMAZOLE 1% CREAM)
LF Eng
 Date: 17-09-2024
 Printed Size: 12.7x18.8 cm (WxH)
 Special Colours: **K=100**