

 ZIM LABORATORIES LIMITED B-21/22, MIDC Area, Kalmeshwar, Nagpur 441501, Maharashtra State, India	Mod. 01
Product Name : Carton of KIDIMAC (Azithromycin For Oral Suspension USP 200 mg)	
Material : GSM NLT 300 ITC Cyber XL paper board	
Date of Preparation : 26/06/2020	
Size : (L) 38 x (W) 38 x (H) 84 mm	
Artwork Code : *PC190X/XX/XX	
Carton Type : Open top Carton. Bottom Flaps are automatically-Locked.	
Dossier Code : R000051/00	
Packing Type : HDPE Bottle	
Language : English	
Other Instruction : NA	
Finished : UV Coating	
Pack Size :15 mL	
Blister Size : NA	
Embossing : NA	
Foil Stamping : NA	
Colour : P. Process Cyan C, P. Red 032 C, P. 354 C, P. 2945 C and Black P. Process Cyan C P. Red 032 C P. 354 C P. 2945 C Black	
*PC190/X/XX/XX Where X is variable and it will be change at the time of commercialised. All dimension in mm	

	15 ml Azithromycin for Oral Suspension USP 200 mg/5 ml		15 ml Azithromycin for Oral Suspension USP 200 mg/5 ml	MAL No. : MAL 20057001ACZ Mfg. Lic. No. : 1036 Batch No. Mfg. Date Exp. Date Controlled Medicine / Ubat Terkawal Jauhi ubat dari kanak-kanak Keep medicine out of reach of children WARNING: Unsuitable for phenylketonurics. Product registration holder: Healol Pharmaceuticals Sdn Bhd Healol pharmaceuticals Sdn Bhd 74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju, 53300 Kuala Lumpur, Malaysia. Manufactured by : ZIM LABORATORIES LIMITED B-21/22, MIDC Area, Kalmeshwar, Nagpur, Maharashtra State, 441501, India
	Composition : Each 5 ml of reconstituted suspension contains : Azithromycin Dihydrate USP 200 mg Eq. to Azithromycin 200 mg Preservatives Methyl Hydroxybenzoate 0.2 % w/v & Propyl Hydroxybenzoate 0.02 % w/v. Dosage : As directed by the Physician. Direction For Use : 1. Tap the bottle to loosen the powder. 2. Add 10ml of boiled and cooled water in bottle. 3. Shake vigorously to mix medicine properly. 4. Not to be injected. Storage : Before reconstitution store below 30°C. After reconstitution store the suspension at 25°C to 30°C. Keep taking the medicine until the course is complete. Reconstituted suspension should be used within 10 days. Discard the reconstituted suspension if more than 10 days. Do not freeze.			

History Sheet		
Revision No.	Reason	Date
Mod. No.: 01	Change colour scheme same as like 30ml as per customer email on dated 16/12/2020.	17/12/2020