

## What is in this leaflet

1. What ACEVIZ is used for
2. How ACEVIZ works
3. Before you use ACEVIZ
4. How to use ACEVIZ
5. While you are using it
6. Side effects
7. Storage and Disposal of ACEVIZ
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

## What is ACEVIZ used for

ACEVIZ is indicated for all respiratory tract diseases leading to the formation of thick secretions difficult to be expectorated, such as acute and chronic bronchitis, laryngitis, sinusitis, tracheitis, influenza, bronchial asthma and (as complementary treatment) mucoviscidosis.

## How ACEVIZ works

Each sachet contains the active substance acetylcysteine. Acetylcysteine belongs to a group of medicines called mucolytics which work by thinning the mucus (phlegm) so that it can be coughed up more easily.

## Before you use ACEVIZ

### -When you must not use it

Do not take ACEVIZ if:

- you are allergic to the ingredients of this product
- you are pregnant; trying to get pregnant or think you may be pregnant.
- you are breast-feeding.

### -Before you start to use it

Ask your doctor or pharmacist for advice before taking any medicine. Please make sure you have understood the explanation given by your doctor or pharmacist before taking this medicine. Speak to your doctor /pharmacist if you have any other health related problems.

### -Taking other medications

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Tell your doctor if you are taking the following medicines before starting ACEVIZ:

- medicines to treat coughs (e.g. cough syrup)
- activated charcoal, used to treat poisoning
- drugs used to treat bacterial infections, antibiotics (e.g. cefalexin, cefuroxime, cefadroxil, cefaclor, ceftriaxone)
- medicines to treat angina attacks, (tightness in your chest, neck and arm) (e.g. glyceryl trinitrate)

## How to use ACEVIZ

### -How much to use

*Usual posology for acute diseases:*

Adults & children above 12 years old: 600 mg daily, divided into one or more administrations (e.g. 200 mg sachets 3 times daily).

Children from 2 to 12 years of age: 200 mg twice daily.

*Special posologies:*

Long-term treatment: 400–600 mg daily, divided into one or more administrations, with a maximum treatment duration of 3 - 6 months.

If the excessive mucus production and the consequent cough do not disappear after a two- week treatment, the diagnosis should be re-evaluated in order to exclude a possible malignant disease of the respiratory tract.

Not recommended in children below two (2) years of age.

### -When to use it

Use as directed by your doctor or pharmacist.

### -How long to use it

Continue taking ACEVIZ for as long as your doctor recommends.

### -If you forget to use it

Try to take ACEVIZ as prescribed. However, if you misses a dose, just resume the usual schedule of one sachet once daily.

Do not take a double dose to make up for a forgotten dose.

### -If you use too much (overdose)

Contact your doctor immediately for advice or go to the Emergency Department of your nearest hospital. An acute overdose of ACEVIZ can cause gastrointestinal symptoms such as nausea, vomiting and diarrhoea.

## While you are using it

### -Things you must do

Take your medicine exactly as your doctor has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking ACEVIZ.

Tell your doctor immediately if you become pregnant while taking this medication.

### -Things you must not do

Do not breastfeed when taking ACEVIZ.

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give ACEVIZ to anyone else, even if they have the same symptoms or condition as you.

### -Things to be careful of

If your condition's symptoms become worse, contact your doctor immediately.

## Side effects

ACEVIZ may cause severe allergy and serious skin reactions. Stop using ACEVIZ and seek medical assistance

# ACEVIZ

Acetylcysteine Powder for Oral Solution 200 mg

Consumer Medication Information Leaflet (RiMUP)

immediately if you experience any of the following symptoms:

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

## Storage and Disposal of ACEVIZ

### -Storage

Store at temperature not exceeding 30°C.

Keep out of reach of children.

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product Description

### -What it looks like

White to off-white, granular powder after reconstitution in water forms a clear solution.

### -Ingredients

- Active ingredient(s)  
Acetylcysteine
- Inactive ingredient(s)  
Vitamin C, Sodium  
Saccharin, Sucrose, Flavour  
Orange.

### -MAL number:

## Manufacturer

XL Laboratories Pvt. Ltd.  
E-1223, Phase-I Extn., (Ghatal)  
RIICO Indl. Area, Bhiwadi-301019,  
Rajasthan, India.

## Product Registration Holder

Healol Pharmaceuticals Sdn.Bhd  
74-3, Jalan Wangsa Delima 6,  
KLSC Wangsa Maju,  
53300 Kuala Lumpur, Malaysia.

## Date of revision

-12/03/2020

## Serial Number

NPRA(R2)20/039