

Pharmacode position will vary
as per width of PI

FRONT SIDE

340.00 mm



Package Insert

1. NAME OF THE MEDICINAL PRODUCT

Flamant 90 (Etoricoxib 90 mg film-coated tablets)
Flamant 120 (Etoricoxib 120 mg film-coated tablets)

2. QUALITATIVE AND QUANTITATIVE COMPOSITION:

Etoricoxib 90 mg film coated tablets is White to off white, apple-shaped, biconvex, film-coated tablets debossed "149" on one side and plain on other side.
Etoricoxib 120 mg film coated tablets is Light green to green, apple-shaped, biconvex, filmcoated tablets debossed "150" on one side and plain on other side.

3. PHARMACEUTICAL FORM

Tablets

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Etoricoxib Film Coated Tablets is indicated for:

- Acute and chronic treatment of the signs and symptoms of osteoarthritis (OA) and rheumatoid arthritis (RA)
- Treatment of ankylosing spondylitis (AS)
- Treatment of acute gouty arthritis
- Chronic low back pain (30 mg and 60 mg only)
- Treatment of acute pain, including that related to primary dysmenorrhoea and minor dental procedures.

The decision to prescribe a selective COX-2 inhibitor should be based on an assessment of the individual patient's overall risks.

4.2 Posology and method of administration

Posology

Etoricoxib Film Coated Tablets is administered orally. Etoricoxib Film Coated Tablets may be taken with or without food. Etoricoxib Film Coated Tablets should be administered for the shortest duration possible and the lowest effective daily dose should be used.

Osteoarthritis

The recommended dose is 30 mg or 60 mg once daily.

Rheumatoid arthritis

The recommended dose is 60 mg once daily. In some patients with insufficient relief from symptoms, an increased dose of 90 mg once daily may increase efficacy. Once the patient is clinically stabilised, down-titration to a 60 mg once daily dose may be appropriate. In the absence of an increase in therapeutic benefit, other therapeutic options should be considered.

Ankylosing spondylitis

The recommended dose is 60 mg once daily. In some patients with insufficient relief from symptoms, an increased dose of 90 mg once daily may increase efficacy. Once the patient is clinically stabilised, down-titration to a 60 mg once daily dose may be appropriate. In the absence of an increase in therapeutic benefit, other therapeutic options should be considered.

Chronic low back pain

The recommended dose is 60 mg once daily

Acute Pain

In the following acute painful conditions, Etoricoxib Film Coated Tablets should be used only for the acute symptomatic period, limited to a maximum of 8 days treatment:

Acute Gouty Arthritis

The recommended dose is 120 mg once daily.

Primary Dysmenorrhoea

The recommended dose is 120 mg once daily.

Minor Dental Procedures

The recommended dose is 90 mg once daily.

Doses greater than those recommended for each indication have either not demonstrated additional efficacy or have not been studied. Therefore:

- The dose for QA should not exceed 60 mg daily.
- The dose for RA should not exceed 90 mg daily.
- The dose for ankylosing spondylitis should not exceed 90 mg daily.
- The dose for acute gout should not exceed 120 mg daily.
- The dose for acute pain and primary dysmenorrhoea should not exceed 120 mg daily.
- The dose of chronic low back pain should not exceed 60 mg daily.
- The dose for minor dental procedures should not exceed 90 mg daily.

As the cardiovascular risks of selective COX-2 inhibitors may increase with dose and duration of exposure, the shortest duration possible and the lowest effective daily dose should be used.

The patient's need for symptomatic relief and response to therapy should be re-evaluated periodically.

Elderly, Gender, Race

No dosage adjustment in Etoricoxib Film Coated Tablets is necessary for the elderly or based on gender or race.

Hepatic Insufficiency

In patients with mild hepatic insufficiency (Child-Pugh score 5-6), a dose of 60 mg once daily should not be exceeded. In patients with moderate hepatic insufficiency (Child-Pugh score 7-9), the dose should be reduced; a dose of 60 mg every other day should not be exceeded, administration of 30 mg once daily can also be considered. There are no clinical or pharmacokinetic data in patients with severe hepatic insufficiency (Child-Pugh score >9).

Renal Insufficiency

In patients with advanced renal disease (creatinine clearance <30 mL/min), treatment with Flamant is not recommended. No dosage adjustment is necessary for patients with lesser degrees of renal insufficiency (creatinine clearance =30 mL/min).

Flamant 90 & Flamant 120 consist of 90 & 120mg tablet. Flamant 90 & Flamant 120 are not able to deliver all approved dose regimens of etoricoxib. Therefore, other approved dosage forms and strengths of etoricoxib should be used in such cases.

Mode of administration: Oral

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
Active peptic ulceration or active gastro-intestinal (GI) bleeding.

Patients who, after taking acetylsalicylic acid or NSAIDs including COX-2 (cyclooxygenase-2) inhibitors, experience bronchospasm, acute rhinitis, nasal polyps, angioneurotic oedema, urticaria, or allergic-type reactions.

Pregnancy and lactation (see sections 4.6 and 5.3).

Severe hepatic dysfunction (serum albumin <25g/l or Child-Pugh score ≥10).

Estimated renal creatinine clearance <30 ml/min.
Children and adolescents under 16 years of age.

Inflammatory bowel disease.

Congestive heart failure (NYHA II-IV).
Contraindication for Etoricoxib in patients with hypertension (high blood pressure) whose blood pressure is not under control.

Established ischaemic heart disease, peripheral arterial disease, and/or cerebrovascular disease.

Patients who have increased risk of cardiovascular disease (ischaemic heart disease and stroke).

4.4 Special warnings and precautions for use

Gastrointestinal effects

Upper gastrointestinal complications [perforations, ulcers or bleedings (PUBs)], some of them resulting in fatal outcome, have occurred in patients treated with etoricoxib.

Caution is advised with treatment of patients most at risk of developing a gastrointestinal complication with NSAIDs; the elderly, patients using any other NSAID or acetylsalicylic acid concomitantly or patients with a prior history of gastrointestinal disease, such as ulceration and GI bleeding. There is a further increase in the risk of gastrointestinal adverse effects (gastrointestinal ulceration or other gastrointestinal complications) when etoricoxib is taken concomitantly with acetylsalicylic acid even at low doses. A significant difference in GI safety between selective COX-2 inhibitors + acetylsalicylic acid vs. NSAIDs + acetylsalicylic acid has not been demonstrated in long-term clinical trials (see section 5.1).

Cardiovascular effects

Clinical trials suggest that the selective COX-2 inhibitor class of drugs may be associated with a risk of thrombotic events (especially myocardial infarction (MI) and stroke), relative to placebo and some NSAIDs, including etoricoxib, can be associated with new onset or recurrent congestive heart failure. For information regarding a dose related response for etoricoxib Caution should be exercised in patients with a history of cardiac failure, left ventricular dysfunction, or hypertension and in patients with pre-existing oedema from any other reason. If there is clinical evidence of deterioration in the condition of these patients, appropriate measures including discontinuation of etoricoxib should be taken.

Patients with significant risk factors for cardiovascular events (e.g. hypertension, hyperlipidaemia, diabetes mellitus, smoking) should only be treated with etoricoxib after careful consideration.

COX-2 selective inhibitors are not a substitute for acetylsalicylic acid for prophylaxis of cardiovascular thrombo-embolic diseases because of their lack of antiplatelet effect. Therefore, anti-platelet therapies should not be discontinued.

Renal effects

Renal prostaglandins may play a compensatory role in the maintenance of renal perfusion. Therefore, under conditions of compromised renal perfusion, administration of etoricoxib may cause a reduction in prostaglandin formation and, secondarily, in renal blood flow, and thereby impair renal function. Patients at greatest risk of this response are those with pre-existing significantly impaired renal function, uncompensated heart failure, or cirrhosis. Monitoring of renal function in such patients should be considered.

Fluid retention, oedema and hypertension

As with other medicinal products known to inhibit prostaglandin synthesis, fluid retention, oedema and hypertension have been observed in patients taking etoricoxib. All Nonsteroidal Anti-inflammatory Drugs (NSAIDs), including etoricoxib, can be associated with new onset or recurrent congestive heart failure. For information regarding a dose related response for etoricoxib Caution should be exercised in patients with a history of cardiac failure, left ventricular dysfunction, or hypertension and in patients with pre-existing oedema from any other reason. If there is clinical evidence of deterioration in the condition of these patients, appropriate measures including discontinuation of etoricoxib should be taken.

Etoricoxib may be associated with more frequent and severe hypertension than some other NSAIDs and selective COX-2 inhibitors, particularly at high doses. Therefore, hypertension should be controlled before treatment with etoricoxib and special attention should be paid to blood pressure monitoring during treatment with etoricoxib. Blood pressure should be monitored within two weeks after initiation of treatment and periodically thereafter. If blood pressure rises significantly, alternative treatment should be considered.

Hepatic effects

Elevations of alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST) (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials treated for up to one year with etoricoxib 30, 60 and 90 mg daily. Any patients with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver function test has occurred, should be monitored. If signs of hepatic insufficiency occur, or if persistently abnormal liver function tests (three times the upper limit of normal) are detected, etoricoxib should be discontinued.

General

If during treatment, patients deteriorate in any of the organ system functions described above, appropriate measures should be taken and discontinuation of etoricoxib therapy should be considered. Medically appropriate supervision should be maintained when using etoricoxib in the elderly and in patients with renal, hepatic, or cardiac dysfunction. Caution should be used when initiating treatment with etoricoxib in patients with dehydration. It is advisable to rehydrate patients prior to starting therapy with etoricoxib.

Serious skin reactions, some of them fatal, including exfoliative dermatitis, Stevens-Johnson syndrome, and toxic epidermal necrolysis, have been reported very rarely in association with the use of NSAIDs and some selective COX-2 inhibitors during post-marketing surveillance (see section 4.8). Patients appear to be at highest risk for these reactions early in the course of therapy with the onset of the reaction occurring in the majority of cases within the first month of receiving etoricoxib (see section 4.8). Some selective COX-2 inhibitors have been associated with an increased risk of skin reactions in patients with a history of any drug allergy. Etoricoxib should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of hypersensitivity.

Etoricoxib may mask fever and other signs of inflammation.

Caution should be exercised when co-administering etoricoxib with warfarin or other oral anticoagulants.

The use of etoricoxib, as with any medicinal product known to inhibit cyclooxygenase / prostaglandin synthesis, is not recommended in women attempting to conceive.

Etoricoxib Film Coated Tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Warning when prescribing COX-2 Inhibitors to patients with risk factors of heart disease, hypertension (high blood pressure), hyperlipidemia, diabetes, smoking patient and patient with peripheral arterial disease.

Given the association between cardiovascular risk and exposure to COX-2 Inhibitors, doctors are advised to use the lowest effective dose for the shortest possible duration of treatment.

Risk of GI Ulceration, Bleeding and Perforation with NSAID

Serious GI toxicity such as bleeding, ulceration and perforation can occur at any time, with or without warning symptoms, in patients treated with NSAID therapy. Although minor upper GI problems (e.g. dyspepsia) are common, usually developing early in therapy, prescribers should remain alert for ulceration and bleeding in patients treated with NSAIDs even in the absence of previous GI tract symptoms. Studies to date have not identified any subset of patients not at risk of developing peptic ulceration and bleeding. Patients with prior history of serious GI events and other risk factors associated with peptic ulcer disease (e.g. alcoholism, smoking, and corticosteroid therapy) are at increased risk. Elderly or debilitated patients seem to tolerate ulceration or bleeding less than other individuals and account for most spontaneous reports for fatal GI events.

4.5 Interaction with other medicinal products and other forms of interaction

Pharmacodynamic interactions

Oral anticoagulants: In subjects stabilised on chronic warfarin therapy, the administration of etoricoxib 120 mg daily was associated with an approximate 13% increase in prothrombin time International Normalised Ratio (INR). Therefore, patients receiving oral anticoagulants should be closely monitored for their prothrombin time INR, particularly in the first few days when therapy with etoricoxib is initiated or the dose of etoricoxib is changed.

Diuretics, ACE inhibitors and Angiotensin II Antagonists: NSAIDs may reduce the effect of diuretics and other antihypertensive drugs. In some patients with compromised renal function (e.g. dehydrated patients or elderly patients with compromised renal function) the co-administration of an ACE inhibitor or Angiotensin II antagonist and agents that inhibit cyclo-oxygenase may result in further deterioration of renal function, including possible acute renal failure, which is usually reversible. These interactions should be considered in patients taking etoricoxib concomitantly with ACE inhibitors or angiotensin II antagonists. Therefore, the combination should be administered with caution, especially in the elderly. Patients should be adequately hydrated and consideration should be given to monitoring of renal function after initiation of concomitant therapy, and periodically thereafter.

Acetylsalicylic Acid: In a study in healthy subjects, at steady state, etoricoxib 120 mg once daily had no effect on the antiplatelet activity of acetylsalicylic acid (81 mg once daily). Etoricoxib can be used concomitantly with acetylsalicylic acid at doses used for cardiovascular prophylaxis (low-dose acetylsalicylic acid). However, concomitant administration of low dose acetylsalicylic acid with etoricoxib may result in an increased rate of GI ulceration or other complications compared to use of etoricoxib alone. Concomitant administration of etoricoxib with doses of acetylsalicylic acid above those for cardiovascular prophylaxis or with other NSAIDs is not recommended.

Cyclosporin and tacrolimus: Although this interaction has not been studied with etoricoxib, coadministration of cyclosporin or tacrolimus with any NSAID may increase the nephrotoxic effect of cyclosporin or tacrolimus. Renal function should be monitored when etoricoxib and either of these drugs is used in combination.

Pharmacokinetic interactions

The effect of etoricoxib on the pharmacokinetics of other drugs

Lithium: NSAIDs decrease lithium renal excretion and therefore increase lithium plasma levels. If necessary, monitor blood lithium closely and adjust the lithium dosage while the combination is being taken and when the NSAID is withdrawn.

Methotrexate: Two studies investigated the effects of etoricoxib 60, 90 or 120 mg administered once daily for seven days in patients receiving once weekly methotrexate doses of 7.5 to 20 mg for rheumatoid arthritis. Etoricoxib at 60 and 90 mg had no effect on methotrexate plasma concentrations or renal clearance. In one study, etoricoxib 120 mg had no effect, but in the other study, etoricoxib 120 mg increased methotrexate plasma concentrations by 28% and reduced renal clearance of methotrexate by 13%. Adequate monitoring for methotrexate-related toxicity is recommended when etoricoxib and methotrexate are administered concomitantly.

Oral contraceptives: Etoricoxib 60 mg given concomitantly with an oral contraceptive containing 35 micrograms ethinyl estradiol (EE) and 0.5 to 1mg norethindrone for 21 days increased the steady state AUC_{0-24hr} of EE by 37%. Etoricoxib 120 mg given with the same oral contraceptive concomitantly or separated by 12 hours, increased the steady state AUC_{0-24hr} of EE by 50 to 60%. This increase in EE concentration should be considered when selecting an oral contraceptive for use with etoricoxib. An increase in EE exposure can increase the incidence of adverse events associated with oral contraceptives (e.g., venous thrombo-embolic events in women at risk).

Hormone Replacement Therapy (HRT): Administration of etoricoxib 120 mg with hormone replacement therapy consisting of conjugated estrogens (0.625 mg PREMARINTM) for 28 days, increased the mean steady state AUC_{0-24hr} of unconjugated estrone (41%), equin (76%), and 17-β-estradiol (22%). The effect of the recommended chronic doses of etoricoxib (30, 60, and 90 mg) has not been studied. The effects of etoricoxib 120 mg on the exposure (AUC_{0-24hr}) to these estrogenic components of PREMARIN were less than half of those observed when PREMARIN was administered alone and the dose was increased from 0.625 to 1.25 mg. The clinical significance of these increases is unknown and higher doses of PREMARIN were not studied in combination with etoricoxib. These increases in estrogenic concentration should be taken into consideration when selecting post-menopausal hormone therapy for use with etoricoxib because the increase in oestrogen exposure might increase the risk of adverse events associated with HRT.

Prednisone/prednisolone: In drug-interaction studies, etoricoxib did not have clinically important effects on the pharmacokinetics of prednisone/prednisolone.

Digoxin: Etoricoxib 120 mg administered once daily for 10 days to healthy volunteers did not alter the steady-state plasma AUC_{0-24hr} or renal elimination of digoxin. There was an increase in digoxin C_{max} (approximately 33%). This increase is not generally important for most patients. However, patients at high risk of digoxin toxicity should be monitored for this when etoricoxib and digoxin are administered concomitantly.

Effect of etoricoxib on drugs metabolised by sulfotransferases

Etoricoxib is an inhibitor of human sulfotransferase activity, particularly SUL1E1, and has been shown to increase the serum concentrations of ethinyl estradiol. While knowledge about effects of multiple sulfotransferases is presently limited and the clinical consequences for many drugs are still being examined, it may be prudent to exercise care when administering etoricoxib concurrently with other drugs primarily metabolised by human sulfotransferases (e.g., oral salbutamol and minoxidil).

Effect of etoricoxib on drugs metabolised by CYP isoenzymes

Based on *in vitro* studies, etoricoxib is not expected to inhibit cytochromes P450 (CYP) 1A2, 2C9, 2C19, 2D6, 2E1 or 3A4. In a study in healthy subjects, daily administration of etoricoxib 120 mg did not alter hepatic CYP3A4 activity as assessed by the erythromycin breath test.

Effects of other drugs on the pharmacokinetics of etoricoxib

The main pathway of etoricoxib metabolism is dependent on CYP enzymes. CYP3A4 appears to contribute to the metabolism of etoricoxib *in vivo*. *In vitro* studies indicate that CYP2D6, CYP2C9, CYP1A2 and CYP2C19salsocatalyse the main metabolic pathway, but their quantitative roles have not been studied *in vivo*.

Ketoconazole: Ketoconazole, a potent inhibitor of CYP3A4, dosed at 400 mg once a day for 11 days to healthy volunteers, did not have any clinically important effect on the single-dose pharmacokinetics of 60 mg etoricoxib (43% increase in AUC).

Rifampicin: Co-administration of etoricoxib with rifampicin, a potent inducer of CYP enzymes, produced a 65% decrease in etoricoxib plasma concentrations. This interaction may result in recurrence of symptoms when etoricoxib is coadministered with rifampicin. While this information may suggest an increase in dose, doses of etoricoxib greater than those listed for each indication have not been studied in combination with rifampicin and are therefore not recommended.

Antacids: Antacids do not affect the pharmacokinetics of etoricoxib to a clinically relevant extent.

4.6 Fertility, pregnancy and lactation

Pregnancy

The use of etoricoxib, as with any drug substance known to inhibit COX-2, is not recommended in women attempting to conceive. No clinical data on exposed pregnancies are available for etoricoxib. Studies in animals have shown reproductive toxicity (see section 5.3). The potential for human risk in pregnancy is unknown. Etoricoxib, as with other medicinal products inhibiting prostaglandin synthesis, may cause uterine inertia and premature closure of the ductus arteriosus during the last trimester. Etoricoxib is contraindicated in pregnancy (see section 4.3). If a woman becomes pregnant during treatment, etoricoxib must be discontinued.

Lactation

It is not known whether etoricoxib is excreted in human milk. Etoricoxib is excreted in the milk of lactating rats. Women who use etoricoxib must not breast feed (see sections 4.3 and 5.3).

4.7 Effects on ability to drive and use machines

Patients who experience dizziness, vertigo or somnolence while taking etoricoxib should refrain from driving or operating machinery.

4.8 Undesirable effects

In clinical trials, etoricoxib was evaluated for safety in 7152 individuals, including 4614 patients with OA, RA, chronic low back pain or ankylosing spondylitis (approximately 600 patients with OA or RA were treated for one year or longer).

In clinical studies, the undesirable effects profile was similar in patients with OA or RA treated with etoricoxib for one year or longer.

In a clinical study for acute gouty arthritis, patients were treated with etoricoxib 120 mg once daily for eight days. The adverse experience profile in this study was generally similar to that reported in the combined OA, RA, and chronic low back pain studies.

In a cardiovascular safety outcomes program of pooled data from three active comparator controlled trials, 17, 412 patients with OA or RA were treated with etoricoxib (60 mg or 90 mg) for a mean duration of approximately 18 months. The safety data and details from this program are presented in section 5.1.

The following undesirable effects were reported at an incidence greater than placebo in clinical trials in patients with OA, RA, chronic low back pain or ankylosing spondylitis treated with etoricoxib 30 mg, 60 mg or 90 mg for up to 12 weeks, or in the MEDAL Program studies, or in post-marketing experience:

[Very Common (≥1/10) Common (≥1/100 to <1/10) Uncommon (≥1/1000 to <1/100) Rare (≥1/10,000 to <1/1,000) Very rare (<1/10,000), not known (cannot be estimated from the available data)]

Infections and infestations:

Uncommon: gastroenteritis, upper respiratory infection, urinary tract infection.

Immune system disorder:

Very rare: hypersensitivity reactions, including angioedema, anaphylactic/anaphylactoid reactions including shock.

Metabolism and nutrition disorders:

Common: oedema/fluid retention
Uncommon: appetite increase or decrease, weight gain.

Psychiatric disorders:

Uncommon: anxiety, depression, mental acuity decreased.
Very rare: confusion, hallucinations.

Nervous system disorder:

Common: dizziness, headache.
Uncommon: dysgeusia, insomnia, paresthesia/hypaesthesia, somnolence.

Eye disorders:

Uncommon: blurred vision, conjunctivitis.

Ear and labyrinth disorders:

Uncommon: tinnitus, vertigo.

Cardiac disorders:

Common: palpitations.
Uncommon: atrial fibrillation, congestive heart failure, non-specific ECG changes, myocardial infarction*.

Vascular disorders:

Common: hypertension.
Uncommon: flushing, cerebrovascular accident*, transient ischaemic attack.
Very rare: hypertensive crisis.

Respiratory, thoracic and mediastinal disorders:

Uncommon: cough, dyspnoea, epistaxis.
Very rare: bronchospasm.

Gastrointestinal disorders:

Common: gastrointestinal disorders (e.g., abdominal pain, flatulence, heartburn), diarrhea, dyspepsia, epigastric discomfort, nausea.
Uncommon: abdominal distention, acid reflux, bowel movement pattern change, constipation, dry mouth, gastroduodenal ulcer, irritable bowel syndrome, oesophagitis, oral ulcer, vomiting, gastritis.
Very rare: peptic ulcers including gastrointestinal perforation and bleeding (mainly in the elderly).

Hepatobiliary disorders:

Very rare: hepatitis.

Skin and subcutaneous tissue disorders:

Common: ecchymosis.
Uncommon: facial oedema, pruritus, rash.

610.00 mm

Very rare: urticaria, Stevens-Johnson syndrome, toxic epidermal necrolysis.

Musculoskeletal, connective tissue and bone disorders:
 Uncommon: muscular cramp/spasm, musculoskeletal pain/stiffness.

Renal and urinary disorders:
 Uncommon: proteinuria.
 Very rare: renal insufficiency, including renal failure, usually reversible upon discontinuation of treatment (see section 4.4).

General disorders and administration site conditions:
 Common: asthenia/fatigue, flu-like disease.
 Uncommon: chest pain.

Investigations:
 Common: ALT increased, AST increased.
 Uncommon: blood urea nitrogen increased, creatine phosphokinase increased, haematocrit decreased, haemoglobin decreased, hyperkalaemia, leukocytes decreased, platelets decreased, serum creatinine increased, uric acid increased.
 Rare: blood sodium decreased.

The following serious undesirable effects have been reported in association with the use of NSAIDs and cannot be ruled out for etoricoxib: nephrotoxicity including interstitial nephritis and nephrotic syndrome; hepatotoxicity including hepatic failure, jaundice and pancreatitis.
 * Based on analyses of long-term placebo and active controlled clinical trials, selective COX-2 inhibitors have been associated with an increased risk of serious thrombotic arterial events, including myocardial infarction and stroke. The absolute risk increase for such events is unlikely to exceed 1% per year based on existing data (uncommon).

4.9 Overdose
 In clinical studies, administration of single doses of etoricoxib up to 500 mg and multiple doses up to 150 mg/day for 21 days did not result in significant toxicity. There have been reports of acute overdose with etoricoxib, although adverse experiences were not reported in the majority of cases. The most frequently observed adverse experiences were consistent with the safety profile for etoricoxib (e.g. gastrointestinal events, cardiovascular events).
 In the event of overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the GI tract, employ clinical monitoring, and institute supportive therapy, if required.
 Etoricoxib is not dialysable by haemodialysis; it is not known whether etoricoxib is dialysable by peritoneal dialysis.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties
Pharmacotherapeutic group: Anti-inflammatory and antirheumatic products, non-steroids, coxibs.
ATC code: M01 Ah05

Mechanism of Action
 Etoricoxib is an oral, selective cyclo-oxygenase-2 (COX-2) inhibitor within the clinical dose range.
 Across clinical pharmacology studies, Etoricoxib Film Coated Tablets produced dose-dependent inhibition of COX-2 without inhibition of COX-1 at doses up to 150 mg daily. Etoricoxib did not inhibit gastric prostaglandin synthesis and had no effect on platelet function.
 Cyclooxygenase is responsible for generation of prostaglandins. Two isoforms, COX-1 and COX-2 have been identified. COX-2 is the isoform of the enzyme that has been shown to be induced by pro-inflammatory stimuli and has been postulated to be primarily responsible for the synthesis of prostanoid mediators of pain, inflammation, and fever. COX-2 is also involved in ovulation, implantation and closure of the ductus arteriosus, regulation of renal function, and central nervous system functions (fever induction, pain perception and cognitive function). It may also play a role in ulcer healing. COX-2 has been identified in tissue around gastric ulcers in man but its relevance to ulcer healing has not been established.

Clinical efficacy and safety

Efficacy
 In patients with osteoarthritis (OA), etoricoxib 60 mg once daily provided significant improvements in pain and patient assessments of disease status. These beneficial effects were observed as early as the second day of therapy and maintained for up to 52 weeks. Studies with etoricoxib 30 mg once daily demonstrated efficacy superior to placebo over a 12-week treatment period (using similar assessments as the above studies). In a dose ranging study, etoricoxib 60 mg demonstrated significantly greater improvement than 30 mg for all 3 primary endpoints over 6 weeks of treatment. The 30 mg dose has not been studied in osteoarthritis of hands.
 In patients with rheumatoid arthritis (RA), etoricoxib 60 mg and 90 mg once daily both provided significant improvements in pain, inflammation, and mobility. In studies evaluating the 60 mg and 90 mg dose, these beneficial effects were maintained over the 12-week treatment periods.
 In patients experiencing attacks of acute gouty arthritis, etoricoxib 120 mg once daily over an eight-day treatment period, relieved moderate to extreme joint pain and inflammation comparable to indomethacin 50 mg three times daily. Pain relief was observed as early as four hours after initiation of treatment.
 In patients with ankylosing spondylitis, etoricoxib 90 mg once daily provided significant improvements in spine pain, inflammation, stiffness and function. The clinical benefit of etoricoxib was observed as early as the second day of therapy after initiation of treatment and was maintained throughout the 52-week treatment period.
 In studies specifically designed to measure the onset of action of etoricoxib, the onset of action occurred as early as 24 minutes after dosing.

Safety
Multinational Etoricoxib and Diclofenac Arthritis Long-term (MEDAL) Program
 The MEDAL Program was a prospectively designed Cardiovascular (CV) Safety Outcomes Program of pooled data from three randomized, double-blind active comparator-controlled trials, the MEDAL study, EDGE II and EDGE.
 The MEDAL Study was an endpoint driven CV Outcomes study in 17,804 OA and 5,700 RA patients treated with etoricoxib 60 (OA) or 90 mg (OA and RA) or diclofenac 150 mg daily for a mean period of 20.3 months (maximum of 42.3 months, median 21.3 months). In this trial, only serious adverse events and discontinuations due to any adverse events were recorded.
 The EDGE and EDGE II studies compared the gastrointestinal tolerability of etoricoxib versus diclofenac. The EDGE study included 7,111 OA patients treated with a dose of etoricoxib 90 mg daily (1.5 times the dose recommended for OA) or diclofenac 150 mg daily for a mean period of 9.1 months (maximum 16.6 months, median 11.4 months). The EDGE II study included 4,086 RA patients treated with etoricoxib 90 mg daily or diclofenac 150 mg daily for a mean period of 19.2 months (maximum 33.1 months, median 24 months).
 In the pooled MEDAL Program, 34,701 patients with OA or RA were treated for a mean duration of 17.9 months (maximum 42.3 months, median 16.3 months) with approximately 12,600 patients receiving treatment for more than 24 months. Patients enrolled in the Program had a wide range of cardiovascular and gastrointestinal risk factors at baseline. Patients with a recent history of myocardial infarction, coronary artery bypass grafting or percutaneous coronary intervention within 6 months preceding enrolment were excluded. Use of gastroprotective agents and low dose aspirin were permitted in the studies.

Overall Safety:
 There was no significant difference between etoricoxib and diclofenac in the rate of cardiovascular thrombotic events.
 Cardiovascular adverse events were observed more frequently with etoricoxib than with diclofenac, and this effect was dose-dependent (see specific results below). Gastrointestinal and hepatic adverse events were observed significantly more frequently with diclofenac than etoricoxib. The incidence of adverse experiences in EDGE and EDGE II and of adverse experiences considered serious or resulting in discontinuation in the MEDAL study was higher with etoricoxib than diclofenac.

Cardiovascular safety results:
 The rate of confirmed thrombotic cardiovascular serious adverse events (consisting of cardiac, cerebrovascular, and peripheral vascular events) was comparable between etoricoxib and diclofenac, and data are summarized in the table below. There were no statistically significant differences in thrombotic event rates between etoricoxib and diclofenac across all subgroups analyzed including patient categories across a range of baseline cardiovascular risk. When considered separately, the relative risks for confirmed thrombotic cardiovascular serious adverse events with etoricoxib 60 mg or 90 mg compared with diclofenac 150 mg were similar.

	Etoricoxib (N=16,819) 25,836 Patient-Years	Diclofenac (N=16,483) 24,766 Patient-Years	Between Treatment Comparison
	Rate† (95% CI)	Rate† (95% CI)	Relative Risk (95% CI)
Confirmed Thrombotic Cardiovascular Serious Adverse Events			
Per-protocol	1.24 (1.11, 1.38)	1.30 (1.17, 1.45)	0.95 (0.81, 1.11)
Intent-to-treat	1.25 (1.14, 1.36)	1.19 (1.08, 1.30)	1.05 (0.93, 1.19)
Confirmed Cardiac Events			
Per-protocol	0.71 (0.61, 0.82)	0.78 (0.68, 0.90)	0.90 (0.74, 1.10)
Intent-to-treat	0.69 (0.61, 0.78)	0.70 (0.62, 0.79)	0.99 (0.84, 1.17)
Confirmed Cerebrovascular Events			
Per-protocol	0.34 (0.28, 0.42)	0.32 (0.25, 0.40)	1.08 (0.80, 1.46)
Intent-to-treat	0.33 (0.28, 0.39)	0.29 (0.24, 0.35)	1.12 (0.87, 1.44)
Confirmed Peripheral Vascular Events			
Per-protocol	0.20 (0.15, 0.27)	0.22 (0.17, 0.29)	0.92 (0.63, 1.35)
Intent-to-treat	0.24 (0.20, 0.30)	0.23 (0.18, 0.28)	1.08 (0.81, 1.44)

†Events per 100 Patient-Years; CI=confidence interval
 N=total number of patients included in Per-protocol population

Per-protocol: all events on study therapy or within 14 days of discontinuation (excluded: patients who took < 75% of their study medication or took non-study NSAIDs >10% of the time).

Intent-to-treat: all confirmed events up to the end of the trial (included patients potentially exposed to non-study interventions following discontinuation of study medication). Total number of patients randomised, n= 17,412 on etoricoxib and 17,289 on diclofenac. CV mortality, as well as overall mortality, was similar between the etoricoxib and diclofenac treatment groups.

Cardiorenal Events:
 Approximately 50% of patients enrolled in the MEDAL study had a history of hypertension at baseline. In the study, the incidence of discontinuations due to hypertension-related adverse events was statistically significantly higher for etoricoxib than for diclofenac. The incidence of congestive heart failure adverse events (discontinuations and serious events) occurred at similar rates on etoricoxib 60 mg compared to diclofenac 150 mg but was higher for etoricoxib 90 mg compared to diclofenac 150 mg (statistically significant for 90 mg etoricoxib vs. 150 mg diclofenac in MEDAL OA cohort).

The incidence of confirmed congestive heart failure adverse events (events that were serious and resulted in hospitalisation or a visit to an emergency department) was non-significantly higher with etoricoxib than diclofenac 150 mg, and this effect was dose-dependent. The incidence of discontinuations due to oedema-related adverse events was higher for etoricoxib than diclofenac 150 mg, and this effect was dose-dependent (statistically significant for etoricoxib 90 mg, but not for etoricoxib 60 mg).

The cardiorenal results for EDGE and EDGE II were consistent with those described for the MEDAL Study.
 In the individual MEDAL Program studies, for etoricoxib (60 mg or 90 mg), the absolute incidence of discontinuation in any treatment group was up to 2.6% for hypertension, up to 1.9% for oedema, and up to 1.1% for congestive heart failure, with higher rates of discontinuation observed with etoricoxib 90 mg than etoricoxib 60 mg.

MEDAL Program Gastrointestinal Tolerability Results:
 A significantly lower rate of discontinuations of treatment for any clinical (e.g., dyspepsia, abdominal pain, ulcer) GI adverse event was observed with etoricoxib compared with diclofenac within each of the three component studies of the MEDAL Program. The rates of discontinuations due to adverse clinical GI events per hundred patient-years over the entire period of study was as follows: 3.23 for etoricoxib and 4.96 for diclofenac in the MEDAL Study; 9.12 with etoricoxib and 12.28 with diclofenac in the EDGE study; and 3.71 with etoricoxib and 4.81 with diclofenac in the EDGE II study.

MEDAL Program Gastrointestinal Safety Results:
 Overall upper GI events were defined as perforations, ulcers and bleeds. The subset of overall upper GI events considered complicated included perforations, obstructions, and complicated bleeding; the subset of upper GI events considered uncomplicated included uncomplicated bleeds and uncomplicated ulcers. A significantly lower rate of overall upper GI events was observed with etoricoxib compared to diclofenac. There was no significant difference between etoricoxib and diclofenac in the rate of complicated events. For the subset of upper GI haemorrhage events (complicated and uncomplicated combined), there was no significant difference between etoricoxib and diclofenac. The upper GI benefit for etoricoxib compared with diclofenac was not statistically significant in patients taking concomitant low-dose aspirin (approximately 33% of patients).

The rates per hundred patient-years of confirmed complicated and uncomplicated upper GI clinical events (perforations, ulcers and bleeds (PUBs)) were 0.67 (95% CI 0.57, 0.77) with etoricoxib and 0.97 (95% CI 0.85, 1.10) with diclofenac, yielding a relative risk of 0.69 (95% CI 0.57, 0.83).

The rate for confirmed upper GI events in elderly patients was evaluated and the largest reduction was observed in patients ≥ 75 years of age (1.35 (95% CI 0.94, 1.87) vs. 2.78 (95% CI 2.14, 3.56) events per hundred patient-years for etoricoxib and diclofenac, respectively).

The rates of confirmed lower GI clinical events (small or large bowel perforation, obstruction, or haemorrhage, (POBs)) were not significantly different between etoricoxib and diclofenac.
MEDAL Program Hepatic Safety Results:
 Etoricoxib was associated with a statistically significantly lower rate of discontinuations due to hepatic-related adverse experiences than diclofenac. In the pooled MEDAL Program, 0.3% of patients on etoricoxib and 2.7% of patients on diclofenac discontinued due to hepatic-related adverse experiences. The rate per hundred patient-years was 0.22 on etoricoxib and 1.84 for diclofenac (p-value was <0.001 for etoricoxib vs. diclofenac). However, most hepatic adverse experiences in the MEDAL Program were non-serious.

Additional Thrombotic Cardiovascular Safety Data
 In clinical studies excluding the MEDAL Program Studies, approximately 3,100 patients were treated with etoricoxib ≥60 mg daily for 12 weeks or longer. There was no discernible difference in the rate of confirmed serious thrombotic cardiovascular events between patients receiving etoricoxib ≥60 mg, placebo, or non-naproxen NSAIDs. However, the rate of these events was higher in patients receiving etoricoxib compared with those receiving naproxen 500 mg twice daily. The difference in antiplatelet activity between some COX-1 inhibiting NSAIDs and selective COX-2 inhibitors may be of clinical significance in patients at risk of thrombo-embolic events. Selective COX-2 inhibitors reduce the formation of systemic (and therefore possibly endothelial) prostacyclin without affecting platelet thromboxane. The clinical relevance of these observations has not been established.

Additional Gastrointestinal Safety Data

In two 12-week double-blind endoscopy studies, the cumulative incidence of gastroduodenal ulceration was significantly lower in patients treated with etoricoxib 120 mg once daily than in patients treated with either naproxen 500 mg twice daily or ibuprofen 800 mg three times daily. Etoricoxib had a higher incidence of ulceration as compared to placebo.

Renal Function Study in the Elderly
 A randomized, double-blind, placebo-controlled, parallel-group study evaluated the effects of 15 days of treatment of etoricoxib (90 mg), celecoxib (200 mg bid), naproxen (500 mg bid) and placebo on urinary sodium excretion, blood pressure, and other renal function parameters in subjects 60 to 85 years of age on a 200-mEq/day sodium diet.
 Etoricoxib, celecoxib, and naproxen had similar effects on urinary sodium excretion over the 2 weeks of treatment. All active comparators showed an increase relative to placebo with respect to systolic blood pressures; however, etoricoxib was associated with a statistically significant increase at Day 14 when compared to celecoxib and naproxen (mean change from baseline for systolic blood pressure: etoricoxib 7.7 mmHg, celecoxib 2.4 mmHg, naproxen 3.6 mmHg).

5.2 Pharmacokinetics properties

Absorption
 Orally administered etoricoxib is well absorbed. The absolute bioavailability is approximately 100%. Following 120 mg once-daily dosing to steady state, the peak plasma concentration (geometric mean C_{max} = 3.6 µg/ml) was observed at approximately 1 hour (T_{max}) after administration to fasted adults. The geometric mean area under the curve (AUC_{0-24hr}) was 37.8 µg^h/ml. The pharmacokinetics of etoricoxib are linear across the clinical dose range.
 Dosing with food (a high-fat meal) had no effect on the extent of absorption of etoricoxib after administration of a 120-mg dose. The rate of absorption was affected, resulting in a 36% decrease in C_{max} and an increase in T_{max} by 2 hours.
 These data are not considered clinically significant. In clinical trials, etoricoxib was administered without regard to food intake.

Distribution

Etoricoxib is approximately 92% bound to human plasma protein over the range of concentrations of 0.05 to 5 µg/ml. The volume of distribution at steady state (V_{ss}) was approximately 1.20l in humans.
 Etoricoxib crosses the placenta in rats and rabbits, and the blood-brain barrier in rats.

Biotransformation

Etoricoxib is extensively metabolised with <1% of a dose recovered in urine as the parent drug. The major route of metabolism to form the 6'-hydroxymethyl derivative is catalyzed by CYP enzymes. CYP3A4 appears to contribute to the metabolism of etoricoxib in vivo. In vitro studies indicate that CYP2D6, CYP2C3, CYP1A2 and CYP2C19 also can catalyse the main metabolic pathway, but their quantitative roles in vivo have not been studied.
 Five metabolites have been identified in man. The principal metabolite is the 6'-carboxylic acid derivative of etoricoxib formed by further oxidation of the 6'-hydroxymethyl derivative. These principal metabolites either demonstrate no measurable activity or are only weakly active as COX-2 inhibitors. None of these metabolites inhibit COX-1.

Elimination

Following administration of a single 25-mg radiolabeled intravenous dose of etoricoxib to healthy subjects, 70% of radioactivity was recovered in urine and 20% in faeces, mostly as metabolites. Less than 2% was recovered as unchanged drug.
 Elimination of etoricoxib occurs almost exclusively through metabolism followed by renal excretion. Steady state concentrations of etoricoxib are reached within seven days of once daily administration of 120 mg, with an accumulation ratio of approximately 2, corresponding to a half-life of approximately 22 hours. The plasma clearance after a 25-mg intravenous dose is estimated to be approximately 50 ml/min.

Characteristics in patients

Elderly patients: Pharmacokinetics in the elderly (65 years of age and older) are similar to those in the young.

Gender: The pharmacokinetics of etoricoxib are similar between men and women.

Hepatic impairment: Patients with mild hepatic dysfunction (Child-Pugh score 5-6) administered etoricoxib 60 mg once daily had an approximately 16% higher mean AUC as compared to healthy subjects given the same regimen. Patients with moderate hepatic dysfunction (Child-Pugh score 7-9) administered etoricoxib 60 mg every other day had similar mean AUC to the healthy subjects given etoricoxib 60 mg once daily, etoricoxib 30 mg once daily has not been studied in this population. There are no clinical or pharmacokinetic data in patients with severe hepatic dysfunction (Child-Pugh score ≥10). (See sections 4.2 and 4.3.)

Renal impairment: The pharmacokinetics of a single dose of etoricoxib 120 mg in patients with moderate to severe renal insufficiency and patients with end-stage renal disease on haemodialysis were not significantly different from those in healthy subjects. Haemodialysis contributed negligibly to elimination (dialysis clearance approximately 50 ml/min). (See sections 4.3 and 4.4.)

Paediatric patients: The pharmacokinetics of etoricoxib in paediatric patients (<12 years old) have not been studied.
 In a pharmacokinetic study (n=16) conducted in adolescents (aged 12 to 17) the pharmacokinetics in adolescents weighing 40 to 60 kg given etoricoxib 60 mg once daily and adolescents >60 kg given etoricoxib 90 mg once daily were similar to the pharmacokinetics in adults given etoricoxib 90 mg once daily. Safety and effectiveness of etoricoxib in paediatric patients have not been established (see section 4.2).

5.3 Preclinical safety data

Not applicable

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients:

Calcium hydrogen phosphate (A-TAB), Microcrystalline Cellulose (Avicel PH 101), Croscarmellose sodium (Ac-Di-Sol), Povidone (Kollidon 30), Purified water, Magnesium stearate

Composition of Opadry II 32K510088 Green for Flamant 120 (Etoricoxib 120 mg film-coated tablets)

%w/w	Ingredients/Compendial Reference
39.000	HPMC 2910/HYPROMELLOSE (USP, PhEur, JP, ChP, GB)
28.000	LACTOSE MONOHYDRATE (NF, PhEur, JP, ChP)
22.400	TITANIUM DIOXIDE (USP,FCC,PhEur,JP,JSFA, ChP,GB)
8.000	TRICETIN (USP, FCC, PhEur, JPE,ChP)
1.700	FD&C BLUE #2/INDIGO CARMINE ALUMINUM LAKE (JECFA, JP MO)
0.900	IRON OXIDE YELLOW (NF,JPE, ChP,JECFA)

Composition of Opadry II 32K180001 White for Flamant 90 (Etoricoxib 90 mg film-coated tablets)

%w/w	Ingredients/Compendial Reference
39.000	HPMC 2910/HYPROMELLOSE (USP, PhEur, JP, ChP, GB)
28.000	LACTOSE MONOHYDRATE (NF, PhEur, JP, ChP)
25.000	TITANIUM DIOXIDE (USP,FCC,PhEur,JP,JSFA, ChP,GB)
8.000	TRICETIN (USP, FCC, PhEur, JPE,ChP)

6.2 Incompatibilities

Not applicable

6.3 Shelf life

24 months

6.4 Special precautions for storage

Store below 30°C. Protect from light and moisture. Keep out of reach of children. Store in the original package.

6.5 Nature and contents of container

ALU-PVC/PVDC Blister of 10 Tablets
BOX CONTAINING 30 TABLETS.
 Read the instructions carefully before use.

6.6 Special precautions for disposal

Not applicable

7. MANUFACTURED BY:

Inventia
Inventia Healthcare Limited,
 F1-F1/1-F75/1, Additional Ambernath M.I.D.C.,
 Ambernath (East), Thane 421506, Maharashtra State, India.

8. PRODUCT REGISTRATION HOLDER:

Healio Pharmaceuticals Sdn Bhd
 74-3, Jalan Wangsa Delima 6,
 KLSC Wangsa Maju, 53300,
 Kuala Lumpur, Malaysia

9. DATE OF REVISION

November 2025

Product Name: FLAMANT 90/120- Etoricoxib 90 mg film-coated tablets		Packaging Material: Booklet (Front & Back Side)		Location: Ambernath	
Dimension: 340 x 610 mm (As per Drawing)		Artwork No.: 2012359		SAP Code: 2012359	
Drawing No.: PGB195230692 BK		Pharma Code: 3096		Country/Client: Malaysia_Healio	
Substrate: ITC Superfine Paper		Specification: 40 GSM		Pack Size: 30 Tablets	
Any other Requirement: Folding Size : 30 x 76 & Transparency Required, Booklet to be closed with tear here adhesive tape					
No. of Colour: 01 ■ Black					
Version No.: 1.0		Change History : 1) New Artwork- CC/BD/2025/136			Date: 20-11-2025
Signature and Date					
Department	Client	Packaging Development - R&D	Regulatory Affairs	Corporate Quality	