

# INTASZORYL-1,2,3&4

(Glimepiride Tablets 1 mg, 2 mg, 3 mg & 4 mg)

---

## What is in this leaflet

1. What Intaszoryl Tablet is used for
2. How Intaszoryl works
3. Before you use Intaszoryl
4. How to take Intaszoryl
5. While you are using Intaszoryl
6. Side effects
7. Storage and disposal of Intaszoryl
8. Product description
9. Manufacturer and product registration holder
10. Date of revision

## What Intaszoryl Tablets is used for

Intaszoryl tablets are used for the treatment of non –insulin dependent (type II) diabetes, where blood sugar levels cannot be controlled adequately by diet, physical exercise and weight reduction alone.

Intaszoryl may also be used in combination with an oral antidiabetic containing metformin or with insulin.

## How Intaszoryl works

Intaszoryl stimulates your pancreas to make more insulin which helps to control blood sugar levels.

## Before you use Intaszoryl

### -When you must not take it

- You are allergic to Glimepiride or any of the other ingredients of Intaszoryl tablets.
- If you have insulin dependent (type I) diabetes
- If you have been in a diabetic coma

- If you have ketoacidosis
- If you have severe liver or kidney problems
- If you are pregnant or breast feeding

### -Before you start to take it

Check with your doctor or pharmacist before taking your medicine if:

- You are recovering from an injury, operation, infections with fever, or from other forms of stress, inform your doctor as temporary change of treatment may be necessary.
- You have severe liver or kidney disorder.

### -Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without a prescription.

## Taking Intaszoryl with food and drink:

- Alcohol intake may increase and decrease the blood sugar lowering action of Intaszoryl in an unpredictable way.

## Pregnancy and breast-feeding:

- Intaszoryl are not recommended during pregnancy and breast-feeding.

## How to take Intaszoryl

### -How much to use it

The usual initial dose is 1 mg Intaszoryl once daily. If necessary, the daily dose can be increased. Any increase should be based on regular blood sugar

monitoring and should be gradual.

## Children and adolescents (below 18 years)

Intaszoryl is not to be used in children or adolescents.

## Older people (age 65 years and over)

No dosage adjustment is necessary for elderly patients.

### -When to use it

Always take Intaszoryl exactly as your doctor has told you.

Taking Intaszoryl at the same time each day will help you remember when to take your tablet.

### -How long to use it

Continue taking Intaszoryl tablets every day for as long as your doctor tells you.

### -If you forget to take it

Take the missed dose as soon as you remember. Skip the missed dose if it is almost time for your next scheduled dose. Do not take extra medicine to make up the missed dose.

### -If you take too much (Overdose)

Contact your doctor immediately or go the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

# INTASZORYL-1,2,3&4

(Glimepiride Tablets 1 mg, 2 mg, 3 mg & 4 mg)

---

## While you are using Intaszoryl

### -Things you must do

You should not take this medicine while you are pregnant. Follow your doctor's instructions carefully.

If you do not follow your doctor's instructions, your treatment may not help or you may have unwanted side effects.

The level of sugar in your blood or urine should be checked regularly. Your doctor may also take blood tests to monitor your blood cell levels and liver function.

### -Things you must not do

Do not use Intaszoryl to treat any other complaints unless your doctor says you can.

Do not give this medicine to anyone else, even if their symptoms seem to be similar to yours.

### -Things to be careful of

Your ability to concentrate or react may be impaired if your blood sugar is lowered or raised or if you develop visual problems as a result of such conditions.

Bear in mind that you could endanger yourself or others. Please ask your doctor whether you can drive a car if you:

- Have frequent episodes of hypoglycaemia
- Have fewer or no warning signals of hypoglycaemia

## Side effects

Like all medicines, Intaszoryl tablets can cause side effects,

although not everybody gets them.

### Common side effects:

- Allergic reactions (including Inflammation of blood vessels, often with skin rash) which may develop into serious reactions with difficulty in breathing, fall in blood pressure and sometimes progressing to shock.
- Abnormal liver function including yellowing of the skin and eyes (jaundice), problems with the bile flow cholestasis, inflammation of the liver (hepatitis) or liver failure.
- Allergy (hypersensitivity) of the skin such as itching, rash, hives and increased sensitivity to sun. Some mild allergic reactions may develop into serious reactions.
- Severe hypoglycaemia including loss of consciousness, seizures or coma.

### Rare side effects

- Lower blood sugar than normal (hypoglycaemia)
- Decrease in the number of blood cells:  
Blood platelets (which increases risk of bleeding or bruising)  
White blood cells (which makes infections more likely)  
Red blood cells (which can make the skin pale and cause weakness or breathlessness)  
These problems generally get better after you stop taking Intaszoryl.

### Very rare side effects

- Allergic reactions (including

inflammation of blood vessels, often with skin rash) which may develop into serious reactions with difficulty in breathing, fall in blood pressure and sometimes progressing to shock. If you experience any of these symptoms, tell your doctor immediately.

- Abnormal liver function including yellowing of the skin and eyes (jaundice), impairment of the bile flow (cholestasis), inflammation of the liver (hepatitis) or liver failure.
- Feeling or being sick, diarrhea, feeling full or bloated, and abdominal pain.
- Decrease in the amount of sodium level in your blood (shown by blood tests)

### Other side effects include:

- Allergy (hypersensitivity) of the skin may occur such as itching, rash, hives and increased sensitivity to sun. Some mild allergic reactions may develop into serious reactions with swallowing or breathing problems, swelling of your lips, throat or tongue, tell your doctor immediately.
- Allergic reactions with sulfonylureas, sulfonamides, or related drugs may occur
- Problems with your sight may occur when beginning treatment with Intaszoryl. This is due to changes in blood sugar levels and should soon improve
- Increased liver enzymes

If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

# INTASZORYL-1,2,3&4

(Glimepiride Tablets 1 mg, 2 mg , 3 mg & 4 mg)

---

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website [npra.gov.my](http://npra.gov.my) [Consumers →Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

## Storage and disposal of Intaszoryl

### -Storage

Store below 30°C

Keep this medication in the container it came in, tightly closed, and out of reach of children. Store it away from excess heat and moisture.

### -Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

## Product description

### -What it looks like

Oval shaped, uncoated tablets with score line on one side and plain on other side of tablet. Intaszoryl are packed in PVC/PVDC Alu –Blister of 10 tablets. Each Carton contains 3 such blister. (3x10)

### -Ingredients

-Active Ingredient:

Glimepiride

-Inactive Ingredient:

Lactose Monohydrate  
Sodium Starch Glycolate  
Povidone (K-30)

Purified Water  
Sodium Starch Glycolate  
Magnesium Stearate

### **Coloring agent**

For 1 mg: Iron Oxide red  
For 2 mg: Iron Oxide yellow and Lake of Indigo carmine  
For 3 mg: Iron Oxide Yellow  
For 4 mg: Lake of Indigo carmine

### -MAL Numbers

Intaszoryl-1mg tablets  
MAL10063005AZ  
Intaszoryl -2mg tablets  
MAL10063006AZ  
Intaszoryl – 3mg tablets  
MAL10063007AZ  
Intaszoryl-4mg tablets  
MAL10063008AZ

## Manufacturer

INTAS PHARM LTD.  
PLOT NO. 457/458, VILLAGE  
MATODA TALIKA: SANAND  
382210, INDIA

## Product Registration Holder

Jetpharma Sdn Bhd  
No 13 Jalan Rajawali 2,  
Bandar Puchong Jaya,  
47100 Puchong,  
Selangor, Malaysia.

## Date of revision

05/11/2024

## Serial Number

NPRA(R2/1)220316/851