

LABEL

 Manufacturer & Product Registration Holder: SUNWARD PHARMACEUTICAL SDN. BHD. No. 9, 11 & 17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia		Batch No.: _____ Mfg. Date: _____ Exp. Date: _____	  9 555275 700045
ALLERSIN SYRUP 4MG/5ML 60ml ALLERGIES Each 5ml contains : Chlorpheniramine Maleate 4mg Sodium Benzoate 9.94mg Route of administration: Oral Store at or below 30°C. Protect from light. Keep out of reach of children. Jauhkan dari kanak-kanak. MAL19861154AZ SIN3348P BRU19052595P Controlled Medicine S300C-R12		Please scan QR code for Indication/Dosage/ Precautions/Warnings/Contraindications/ Side Effects/Treatment for overdose. Caution: This may cause drowsiness. If affected do not drive or operate heavy machinery. Avoid alcoholic drink. Warning: When used for treatment of cough and cold: a) Not to be used in children less than 2 years of age. b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.	
Hosp. / Clinic / Pharmacy : _____ Patient Name : _____ R/N : _____ Date : _____ Take / Minum : _____ ml _____ Times daily Before / After meal Kali sehari Sebelum / Selepas makan			

 Manufacturer & Product Registration Holder: SUNWARD PHARMACEUTICAL SDN. BHD. No. 9, 11 & 17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia		Batch No.: _____ Mfg. Date: _____ Exp. Date: _____	  9 555275 700052
ALLERSIN SYRUP 4MG/5ML 100ml ALLERGIES Each 5ml contains : Chlorpheniramine Maleate 4mg Sodium Benzoate 9.94mg Route of administration: Oral Store at or below 30°C. Protect from light. Keep out of reach of children. Jauhkan dari kanak-kanak. MAL19861154AZ SIN3348P Controlled Medicine S300B-R3		Please scan QR code for Indication/Dosage/ Precautions/Warnings/Contraindications/ Side Effects/Treatment for overdose. Caution: This may cause drowsiness. If affected do not drive or operate heavy machinery. Avoid alcoholic drink. Warning: When used for treatment of cough and cold: a) Not to be used in children less than 2 years of age. b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.	
Hosp. / Clinic / Pharmacy : _____ Patient Name : _____ R/N : _____ Date : _____ Take / Minum : _____ ml _____ Times daily Before / After meal Kali sehari Sebelum / Selepas makan			

 Manufacturer & Product Registration Holder: SUNWARD PHARMACEUTICAL SDN. BHD. No. 9, 11 & 17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia		Batch No.: _____ Mfg. Date: _____ Exp. Date: _____	  9 555275 700069
ALLERSIN SYRUP 4MG/5ML 120ml ALLERGIES Each 5ml contains : Chlorpheniramine Maleate 4mg Sodium Benzoate 9.94mg Route of administration: Oral Store at or below 30°C. Protect from light. Keep out of reach of children. Jauhkan dari kanak-kanak. MAL19861154AZ SIN3348P BRU19052595P Controlled Medicine S300D-R11		Please scan QR code for Indication/Dosage/ Precautions/Warnings/Contraindications/ Side Effects/Treatment for overdose. Caution: This may cause drowsiness. If affected do not drive or operate heavy machinery. Avoid alcoholic drink. Warning: When used for treatment of cough and cold: a) Not to be used in children less than 2 years of age. b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.	
Hosp. / Clinic / Pharmacy : _____ Patient Name : _____ R/N : _____ Date : _____ Take / Minum : _____ ml _____ Times daily Before / After meal Kali sehari Sebelum / Selepas makan			

LABEL

		Manufacturer & Product Registration Holder: SUNWARD PHARMACEUTICAL SDN. BHD. No. 9, 11 & 17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia	
ALLERSIN SYRUP 4MG/5ML		3.8Litre	
ALLERGIES			
Each 5ml contains : Chlorpheniramine Maleate 4mg Sodium Benzoate 9.94mg		Please scan QR code for Indication/Dosage/ Precautions/Warnings/Contraindications/ Side Effects/Treatment for overdose.	
Route of administration: Oral		Caution: This may cause drowsiness. If affected do not drive or operate heavy machinery. Avoid alcoholic drink. Warning: When used for treatment of cough and cold: a) Not to be used in children less than 2 years of age. b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.	
Store at or below 30°C. Protect from light.			
Keep out of reach of children. Jauhi dari kanak-kanak.		 	
MAL19861154AZ SIN3348P BRU19052595P			
Controlled Medicine		Batch No.: Mfg. Date: Exp. Date:	
S300F-R7			

3.8L- FOR EXPORT ONLY