

SUNITON TABLET

Orphenadrine Citrate / Paracetamol (35mg / 450mg)

What is in this leaflet

1. What Suniton Tablet is used for
2. How Suniton Tablet works
3. Before you use Suniton Tablet
4. How to use Suniton Tablet
5. While you are using it
6. Side effects
7. Storage and disposal of Suniton Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Suniton Tablet is used for

Suniton Tablet is used for the relief of painful muscle ache and headache.

How Suniton Tablet works

Orphenadrine is a skeletal muscle relaxant that acts in the nervous system to produce muscle relaxant effects. Paracetamol is an analgesic (pain killer) and antipyretic (relieve fever).

Before you use Suniton Tablet

- When you must not use it

Do not take Suniton Tablet if you:

- are hypersensitive to chlorpheniramine citrate, paracetamol or any of the other ingredients of this medicine (listed in Product Description).
- have glaucoma (increase pressure in eyes).
- have gastric outflow obstruction.
- have ulcer in stomach.
- have enlargement of prostate or bladder neck obstruction.
- have severe muscle weakness.
- have liver problem or kidney problem.

- Before you start to use it

Tell your doctor before taking this medicine if you have:

- rapid heart rate or other heart disease.
- impaired liver or kidney function.

This preparation contains Paracetamol. Do not take any other paracetamol containing medicines at the same time.

Pregnancy and breastfeeding

You should check with your doctor if you are or think you may be pregnant, planning to become pregnant or if you are breastfeeding.

Laboratory

Tell your doctor that you are taking Suniton Tablet before doing any medical tests.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. You should talk to your doctor or pharmacist if you are taking:-

- medicines to treat pain and inflammation such as phenylbutazone, aminopyrine, griseofulvin and hexobarbital.
- medicines for seizure such as phenobarbital.
- alcohol.
- medicines to treat cancer (doxorubicin).
- contraceptives, medicine to prevent pregnancy.
- medicine contains anticholinergic drug to treat antidepressant and antihistamine.

How to use Suniton Tablet

- How much to use

You should take Suniton Tablet exactly as directed by your doctor or pharmacist.

Take one to two tablets.

- When to use it

Take three times daily or directed by your doctor or pharmacist.

- How long to use it

Take Suniton Tablet for the duration that your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of Suniton Tablet may cause excitement, confusion, delirium leading to coma. Seizure and rapid heart rate with dilated pupils, urinary retention and acute liver damage may occur.

While you are using it

- Things you must do

- Take Suniton Tablet only with a doctor's prescription.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Suniton Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Suniton Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

This medicine may affect your ability to drive or use machines. If the tablets make you feel light-headedness, dizzy or fainting, do not drive or use machines and contact your doctor immediately.

Allergy alert: Paracetamol may cause severe skin reactions.

Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Side effects

Like all medicines, Suniton Tablet can cause side effects, although not everybody gets them.

Side effects due to orphenadrine citrate are:

- mouth dryness
- rapid heart rate / pounding heart beat
- urinary retention
- blurred vision
- dilated pupils
- increased eye pressure
- weakness
- nausea
- vomiting
- headache
- constipation
- allergic reaction
- itchiness
- hallucinations

- agitation / shaking
- gastric pain
- rarely hives and other skin problem.

Side effects due to paracetamol which has been reported are:

- Blood disorder: low count of red blood cells, white blood cells, and platelets.
- Skin: rashes, severe skin reactions such as Stevens Johnson Syndrome or toxic epidermal necrolysis (Lyell's syndrome).

Any other effects not listed above may also occur in some people. Please refer to your doctor or pharmacist if you experience any other side effects while taking this medication.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Suniton Tablet

- Storage

Store at below 30 °C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A round, flat, white, tablet with single score on one side only.

- Ingredient(s)

- Active Ingredient

- Orphenadrine citrate
- Paracetamol

- Inactive Ingredients

- Corn Starch
- Sodium Starch Glycolate
- Magnesium Stearate
- Isopropyl alcohol
- Polyvinylpyrrolidone (Kolidon 90F)
- Polyvinylpyrrolidone (K90)

- MAL number

MAL20012742AZ

Manufacturer

Sunward Pharmaceutical Sdn. Bhd.
No. 9,11 & 17, Jalan Kempas 4, Taman
Perindustrian Tampoi Indah, 81200
Johor Bahru, Johor, Malaysia.

Product Registration Holder

Sunward Pharmaceutical Sdn. Bhd.
No. 9,11 & 17, Jalan Kempas 4, Taman
Perindustrian Tampoi Indah, 81200
Johor Bahru, Johor, Malaysia.

Date of revision

20/04/2023

Serial Number

NPRA(R2/1)230419/1047