

CONMY TABLETS

Terazosin Hydrochloride (2mg)

What is in this leaflet

1. What Conmy Tablets 2mg is used for
2. How Conmy Tablets 2mg works
3. Before you use Conmy Tablets 2mg
4. How to use Conmy Tablets 2mg
5. While you are using it
6. Side effects
7. Storage and Disposal of Conmy Tablets 2mg
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Conmy Tablets 2mg is used for

It is used as treatment to control high blood pressure and to improve urination in men due to enlargement of prostate gland or known as symptomatic benign prostatic hyperplasia (BPH).

How Conmy Tablets 2mg works

Terazosin hydrochloride relax the muscles in prostate and bladder neck leads to improvement in urine flow.

Before you use Conmy Tablets 2mg

- When you must not use it

Do not take Conmy Tablets 2mg if you:-

- have heart failure due to mechanical obstruction.
- are hypersensitivity to Terazosin hydrochloride or any ingredients listed at the end of this leaflet.

- Before you start to use it

Talk to your doctor before taking Conmy Tablets 2mg, if you have:

- low blood pressure

Pregnancy and breastfeeding

You should check with your doctor if you are pregnant or think you may be pregnant, planning to become pregnant or if you are breastfeeding. The content and effects of Terazosin hydrochloride during breastfeeding are unknown.

Children

Safety and effectiveness in children have not been determined.

- Taking other medicines

There are few medicines which may interact with Conmy Tablets 2mg. It is important to tell your doctor or pharmacist about all the medicines that you are taking, including those obtained without doctor's prescription.

You should talk to your doctor or pharmacist if you are taking:-

- medicines used for pain killer/anti-inflammatory (such as acetaminophen, aspirin, codeine, ibuprofen, indomethacin).
- medicines contains control of high blood pressure agents (calcium channel blocker such as verapamil).
- antibiotics (example, erythromycin, trimethoprim and sulfamethoxazole).
- medicines used to control nose stuffiness (such as phenylephrine hydrochloride, phenylpropanolamine and pseudoephedrine).
- medicine to treat gout (allopurinol).
- medicine for heart illness or high blood pressure (such as atenolol, hydrochlorothiazide, methyclothiazide and propranolol).
- medicines contains steroids.
- medicine used for stomach and intestinal treatment (such as antacids).
- medicine used to treat diabetes.
- medicines contain sedatives and tranquilizers for depressant (example, diazepam).
- medicine contain chlorpheniramine.

How to use Conmy Tablets 2mg

- How much to use

You should take Conmy Tablets 2mg as advised by your doctor or pharmacist. Dosage requirements are variable and must be individualized on the basis of the disease and the response of the individual. Do not take more or less than what your doctor prescribes. Initial recommended dosage should be start at 1mg at bed time.

Oral administration

Benign Prostatic Hyperplasia

The dosage is gradually increase to 2mg, 5mg and 10mg once daily to achieve the desire improvement of symptoms.

High blood pressure

The recommended dosage is 1-5mg once daily at bedtimes.

Your doctor may assess by slowly increased to achieve the benefits from dose as high as 20mg per day to achieve desired blood pressure response and monitor the dosing interval to be sure the control is maintained.

- When to use it

Use Conmy Tablets 2mg according to doctor's prescription.

- How long to use it

It is important to take Conmy Tablets 2mg as long as your doctor prescribes.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too many Conmy Tablets 2mg may cause low blood pressure.

While you are using it

- Things you must do

- Tell all the doctors, dentists and pharmacists treating you that you are taking Conmy Tablets 2mg
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Conmy Tablets 2mg to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, Conmy Tablets 2mg can cause side effects, although not everybody gets them. The most common side effects reported such as feeling tiredness or lack of energy in body, strong pounding of heartbeat, nausea, excessive of fluids in lower legs or hand, sleepiness, runny nose or stuffiness and blurred vision.

The side effects may include dizziness and fainting when standing up quickly from a lying or sitting position (syncopal episodes), back pain, headache, low blood pressure, rapid heartbeat, weight gain, pain around arms and legs, low sexual desires, depression, nervousness, abnormal sensation such as tingling or pricking, difficulty in breathing, sinus infection and impotence.

Post marketing experience: Lack of platelet in blood and prolong of erection in penis have been reported.

Laboratory test: Decrease in haematocrit (ratio of total volume red blood cell), haemoglobin, white blood cells, total protein, and albumin.

Any other effects not listed above may also occur in some people. Please refer to your doctor or pharmacist if you experience any side effects while taking this medication.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835490, or visiting the website npra.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of Conmy Tablets 2mg- Storage

Store at below 30 °C. Protect from light and keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description- What it looks like

Round, flat tablet, orange in colour with “BH”
2
score on one side and “S” on the other side.

- Ingredient(s)- Active Ingredient

- Terazosin (2mg) equivalent to Terazosin Hydrochloride Dihydrate 2.37mg.

- Inactive Ingredients

- Corn Starch
- Iron Oxide (Yellow)
- Lactose Monohydrate
- Magnesium Stearate
- Povidone
- Sodium Starch Glycolate
- Purified Water

- MAL number

MAL09100245AZ

Manufacturer

Taiwan Biotech Co., Ltd.
22, Chieh-Shou Road,
Taoyuan, Taiwan.

Product Registration Holder

Sunward Pharmaceutical Sdn. Bhd.
No. 9,11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

Date of revision

21/05/2019

Serial Number

NPRA(R4/1)150519/00130