

MUCOCLEAR TABLET 30MG

Ambroxol Hydrochloride (30mg)

What is in this leaflet

1. What Mucoclear Tablet is used for
2. How Mucoclear Tablet works
3. Before you use Mucoclear Tablet
4. How to Mucoclear Tablet
5. While you are using it
6. Side effects
7. Storage and disposal of Mucoclear Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Mucoclear Tablet is used for

It is indicated for acute and chronic lung infection associated abnormal mucus secretion and and impaired mucus transport.

How Mucoclear Tablet works

The active ingredient in this medicine Ambroxol hydrochloride. It enhance mucus transport by reducing viscosity and excessive mucus improves expectoration and eases cough. The ambroxol is a metabolite of bromhexine, hence it has the similar action and uses of bromhexine.

Before you use Mucoclear Tablet

- When you must not use it

Do not take this medicine if you are hypersensitive to ambroxol hydrochloride or any of the other ingredients listed at the end of the leaflet.

Pregnancy and breastfeeding

You should check with your doctor if you are or think you may be pregnant, planning to become pregnant or if you are breastfeeding.

It is not recommended for pregnant women and nursing mothers unless the expected benefit outweighs any potential risks.

- Before you start to use it

Tell your doctor before taking this medicine if you have:

- Breastfeeding
- Pregnant

- Taking other medicines

There are a few medicines which may interact with Mucoclear Tablet. It is important to tell your doctor or pharmacist about all the medicines that you are taking, including those obtained without doctor's prescription.

How to use Mucoclear Tablet

- How much to use

You should take Mucoclear Tablet exactly as advised by your doctor or pharmacist. The amount you take each day will depend on your condition. Do not take more or less than what your doctor prescribes.

Oral administration

Adults: 2 tablets twice daily after meals. If in maintenance therapy, 1 tablet three times a day.

Children 5 years and above: One tablet three times daily.

- When to use it

Use Mucoclear Tablet when needed and according to doctor's prescription.

- How long to use it

It is important to take Mucoclear Tablet as long as your doctor prescribes.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

- Things you must do

- Take your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are taking Mucoclear Tablet.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Mucoclear Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Be careful when engaging in activities requiring mental alertness such as driving a car or operating machinery until you know how this medicine affects you.

Side effects

Like all medicines, Mucoclear Tablet may cause side effects such as severe allergy and serious skin reactions. Stop using Mucoclear Tablet and seek for medical assistance immediately if you experience any of the following symptoms:-

Severe allergy: Breathing difficulties, light headedness, skin swellings or rash.

Severe skin reaction: Skin reddening, blisters, rash, fever, sore throat or eye irritation.

Any other effects such as mild burning sensation in the chest, indigestion, nausea and vomiting may also occur in some people. Please refer to your doctor or pharmacist if you experience any other symptoms while taking this medication.

You may report any side effect or adverse drug reaction directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website www.npra.gov.my (Consumers→Reporting).

Storage and disposal of Mucoclear Tablet

- Storage

Store at or below 30°C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Round, convex pink plain tablet with single score on one side and "SW" on the other side.

- Ingredient(s)

- Active Ingredient

- Ambroxol Hydrochloride

- Inactive Ingredients

- Lactose
- Corn Starch
- Cochineal Red E124
- Magnesium Stearate

- MAL number

MAL20001060X

Manufacturer

Sunward Pharmaceutical Sdn. Bhd.
No. 9,11 & 17, Jalan Kempas 4, Taman
Perindustrian Tampoi Indah, 81200
Johor Bahru, Johor, Malaysia.

Product Registration Holder

Sunward Pharmaceutical Sdn. Bhd.
No. 9,11 & 17, Jalan Kempas 4, Taman
Perindustrian Tampoi Indah, 81200
Johor Bahru, Johor, Malaysia.

Date of revision

29/03/2022

Serial Number

NPRA(R2/1)220323/855