

SW HYOSCINE BUTYLBROMIDE TABLET

Hyoscine Butylbromide (10mg)

What is in this leaflet

1. What SW Hyoscine Butylbromide Tablet is used for
2. How SW Hyoscine Butylbromide Tablet works
3. Before you use SW Hyoscine Butylbromide Tablet
4. How to use SW Hyoscine Butylbromide Tablet
5. While you are using it
6. Side effects
7. Storage and disposal of SW Hyoscine Butylbromide Tablet
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision

What SW Hyoscine Butylbromide Tablet is used for

It is used for relief spasm and increased movement of gastrointestinal, gall bladder and urinary tract. It also used to relaxes the lower part of the uterus and relieve painful menstruation.

How SW Hyoscine Butylbromide Tablet works

The active ingredient in the medicine is hyoscine butylbromide. It helps reduces the movement and tone of the gastrointestinal tract and may even reduce the volume of its various secretions.

Before you use SW Hyoscine Butylbromide Tablet

- When you must not use it

Do not take this medicine if you:-

- are allergic to hyoscine butylbromide or any other ingredients of this medicine (listed in Product Description).
- have increased pressure in the eye, esophagus (food tube) movement disorder, prostate enlargement or obstruction of the bladder neck.
- have muscle weakness.

- Before you start to use it

You should check with your doctor if you :-

- have gastric outlet obstruction.
- are elderly person.
- have metabolic, kidney or liver problem.
- are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

There are few medicines which may interact with SW Hyoscine Butylbromide Tablet. Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

You should talk to your doctor or pharmacist if you are taking on the following medicines:-

- Tricyclic anti-depressants, for depression treatment.
- Antihistamines, treatment to relieve symptoms of different types of allergies.
- Quinidine, Disopyramide medication use to treat irregular heartbeat.
- Amantadine, medication for certain type of flu infection.
- Dopamine antagonist, such as Metoclopramide to prevent nausea and vomiting.

How to use SW Hyoscine Butylbromide Tablet

- How much to use

Oral intake.

Adults: One to two tablets, three to four times daily.

Children (6-12 years): One tablet, three times daily.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Use 3-4 times daily or as directed by your doctor or pharmacist.

- How long to use it

Use SW Hyoscine Butylbromide Tablet for the duration that the doctor recommends.

- If you forget to use it

Do not take extra dose to make up for a missed dose. Just take your next dose at usual the time. Consult your doctor or pharmacist on what you should do if you forget to use it.

- If you use too much (overdose)

If you take too much of the medication by mistake, contact your doctor immediately or seek medical attention.

Taking too many tablets may increase the signs and symptoms such as disorientation memory disturbances, dizziness, restlessness, hallucinations or confusions.

While you are using it

- Things you must do

- Use your medicine exactly as your doctor has told you.
- Tell all the doctors, dentists and pharmacists treating you that you are using SW Hyoscine Butylbromide Tablet.
- Tell your doctor immediately if you become pregnant while using this medication.

- Things you must not do

- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give SW Hyoscine Butylbromide Tablet to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, SW Hyoscine Butylbromide Tablet can cause side effects, although not everybody gets them.

The side effects of hyoscine butylbromide include dryness of the mouth with difficulty in swallowing, enlargement of eye pupil, eye vision unable to focus and sensitive to light, increased eye pressure, flushing, dryness of the skin and short term low heart rate followed by increased heart rate with an irregular heart beat and abnormal feeling of pounding heart.

Desire to urinate with the inability to do, as well as reduction in the tone and movement of the gastro-intestinal tract leading to constipation, occasionally vomiting, giddiness and staggering may occur.

Any other effects not listed above may also occur in some people. Please refer to your doctor or pharmacist if you experience any other side effects while taking this medication.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers→ Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and disposal of SW Hyoscine Butylbromide Tablet

- Storage

Store at or below 30 °C. Protect from light.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Round, white, film-coated tablet.

- Ingredient(s)

- Active Ingredient

- Hyoscine Butylbromide

- Inactive Ingredients

- Lactose
- Corn Starch
- Magnesium Stearate
- Povidone-K90
- Isopropyl Alcohol
- Hydroxypropyl Methylcellulose
- Deionised Water
- Titanium Dioxide
- Calcium Carbonate (Light)
- Talcum
- PEG 6000

- MAL number

MAL19921152AZ

MAL19992703ASZ

Manufacturer

(MAL19921152AZ)

Sunward Pharmaceutical Sdn. Bhd.

No. 9,11 & 17, Jalan Kempas 4, Taman

Perindustrian Tampoi Indah, 81200

Johor Bahru, Johor, Malaysia.

(MAL19992703ASZ)

Sunward Pharmaceutical Pte.Ltd

No.11,Wan Lee Road

627943, Singapore.

Product Registration Holder

Sunward Pharmaceutical Sdn. Bhd.

No. 9,11 & 17, Jalan Kempas 4, Taman

Perindustrian Tampoi Indah, 81200

Johor Bahru, Johor, Malaysia.

Date of revision

26/01/2023

Serial Number

NPRA(R4/1)221122/00104