

PACK INSERT FOR MALAYSIA

Recormon® pre-filled syringes



Epoetin beta

Information as set forth in this label only applies to Recormon

1. DESCRIPTION

1.1 Therapeutic / Pharmacologic Class of Drug

Antianaemic agent
ATC code: B03XA

1.2 Type of Dosage Form

Solution for injection (pre-filled syringes) (s.c. or i.v.).

1.3 Route of Administration

Solution for intravenous (i.v.) or subcutaneous (s.c.) injection.

1.4 Sterile / Radioactive Statement

Sterile Product.

1.5 Qualitative and Quantitative Composition

Active ingredient: epoetin beta (recombinant human erythropoietin: produced by recombinant DNA technology in CHO cell line).

Recormon is provided as solution for injection in pre-filled syringes.

The solution is a colourless, clear to slightly opalescent.

Solution for injection in pre-filled syringe:

2000 IU = 16.6 micrograms epoetin beta with 0.3 ml solution for injection

4000 IU = 33.2 micrograms epoetin beta with 0.3 ml solution for injection.

5000 IU = 41.5 micrograms epoetin beta with 0.3 ml solution for injection.

List of excipients:

All presentations contain up to 0.3 mg phenylalanine per pre-filled syringe (see section 2.4.1, Warnings and Precautions, General).

All other excipients include: urea, sodium chloride, polysorbate 20, sodium dihydrogen phosphate, sodium monohydrogen phosphate, calcium chloride, glycine, leucine, isoleucine, threonine, and glutamic acid.

2. CLINICAL PARTICULARS

2.1 Therapeutic Indication(s)

Recormon is indicated for:

- Treatment of anemia associated with chronic renal failure (renal anemia) in patients on dialysis.
- Treatment of symptomatic renal anemia in patients not yet undergoing dialysis.
- Prevention of anemia of prematurity in infants with a birth weight of 750g to 1500 g and a gestational age of less than 34 weeks.
- Treatment of anemia in adult patients with solid tumours and treated with platinum-based chemotherapy prone to induce anemia (cisplatin: 75 mg/m²/cycle; carboplatin: 350 mg/m²/cycle).
- Treatment of symptomatic anemia in adult patients with multiple myeloma, low grade non-Hodgkin's lymphoma or chronic lymphocytic leukemia, who have a relative erythropoietin deficiency and are receiving anti-tumor therapy. Deficiency is defined as an inappropriately low serum erythropoietin level in relation to the degree of anemia.
- Increasing the yield of autologous blood from patients in a pre-donation program.

Its use in this indication must be balanced against the reported increased risk of thromboembolic events. Treatment should only be given to patients with moderate anemia (Hb 10 to 13 g/dl [6.21 to 8.07 mmol/l], no iron deficiency) if blood conserving procedures are not available or insufficient when the scheduled major elective surgery requires a large volume of blood (4 or more units of blood for females or 5 or more units for males).

2.2 Dosage and Administration

Therapy with Recormon should be initiated by physicians experienced in the above mentioned indications. As anaphylactoid reactions were observed in isolated cases, it is recommended that the first dose be administered under medical supervision.

Substitution by any other biological medicinal product requires the consent of the prescribing physician.

The Recormon pre-filled syringe is ready for use.

Under no circumstances should more than one dose be administered per syringe; the medicinal product is for single use only (see section 4.2, Special Instructions for Use, Handling and Disposal).

Treatment of patients with anaemia due to chronic renal failure:

The solution can be administered subcutaneously or intravenously. In case of intravenous administration, the solution should be injected over approximately 2 minutes, e.g. in haemodialysis patients via the arteriovenous fistula at the end of dialysis. For non-haemodialysed patients, subcutaneous administration should always be preferred in order to avoid puncture of peripheral veins.

In CKD patients, the aim of treatment is to reach a target Hb level of 10-12 g/dl. An Hb level of 12 g/dl should not be exceeded. If the rise in haemoglobin is greater than 2 g/dl (1.3 mmol/L) in 4 weeks, an appropriate dose reduction should be considered. In the presence of hypertension or existing cardiovascular, cerebrovascular or peripheral vascular diseases, the weekly increase in Hb and the target Hb should be determined individually taking into account the clinical picture. Patients should be monitored closely to ensure that the lowest dose of Recormon is used to provide adequate control of the symptoms of anaemia.

Treatment with Recormon is divided into two stages:

1. Correction phase

– Subcutaneous administration

The initial dosage is 3 x 20 IU/kg body weight per week. The dosage may be increased every 4 weeks by 3 x 20 IU/kg per week if the increase of Hb is not adequate (< 0.25 g/dl per week). The weekly dose can also be divided into daily doses.

– Intravenous administration

The initial dosage is 3 x 40 IU/kg per week. The dosage may be raised after 4 weeks to 80 IU/kg - three times per week - and by further increments of 20 IU/kg if needed, three times per week, at monthly intervals.

For both routes of administration, the maximum dose should not exceed 720 IU/kg per week.

2. Maintenance phase

To maintain a target Hb value of approximately 10-12 g/dl, the dosage is initially reduced to half of the previously administered amount. Subsequently, the dose is adjusted at intervals of two to four weeks individually for the patient (maintenance dose). In the case of subcutaneous administration, the weekly dose can be given as one injection per week or in divided doses three or seven times per week. Patients who are stable on a once weekly dosing regimen may be switched to once every two weeks administration. In this case dose increases may be necessary.

Treatment with Recormon is normally a long-term therapy. It can, however, be interrupted, if necessary, at any time. Data on the once weekly dosing schedule are based on clinical studies with a treatment duration of 24 weeks.

Treatment of symptomatic anaemia in cancer patients receiving chemotherapy:

The solution is administered subcutaneously; the weekly dose can be given as one injection per week or in divided doses 3 to 7 times per week.

The recommended initial dose is 30,000 IU per week (corresponding to approximately 450 IU/kg body weight per week, based on an average weighted patient).

Recormon treatment is indicated if the haemoglobin value is ≤ 11 g/dl (6.83 mmol/L). Haemoglobin levels should not exceed 13 g/dl (8.07 mmol/L) (see section 3.1.2 Clinical / Efficacy Studies).

If, after 4 weeks of therapy, the haemoglobin value has increased by at least 1 g/dl (0.62 mmol/L), the current dose should be continued. If the haemoglobin value has not increased by at least 1 g/dl (0.62 mmol/L), a doubling of the weekly dose should be considered. If, after 8 weeks of therapy, the haemoglobin value has not increased by at least 1 g/dl (0.62 mmol/L), response is unlikely and treatment should be discontinued. The therapy should be continued for up to 4 weeks after the end of chemotherapy.

The maximum dose should not exceed 60,000 IU per week.

Once the therapeutic objective for an individual patient has been achieved, the dose should be reduced by 25 to 50% in order to maintain haemoglobin at that level. If required, further dose reduction may be instituted to ensure that haemoglobin level does not exceed 13 g/dl.

If the rise in haemoglobin is greater than 2 g/dl (1.3 mmol/L) in 4 weeks, the dose should be reduced by 25 to 50%.

Treatment for increasing the amount of autologous blood:

The solution is administered intravenously over approximately 2 minutes, or subcutaneously.

Recormon is administered twice weekly over 4 weeks. On those occasions where the patient's PCV allows blood donation, i.e. PCV $\geq 33\%$, Recormon is administered at the end of blood donation. During the entire treatment period, a PCV of 48% should not be exceeded.


The dosage must be determined by the surgical team individually for each patient as a function of the required amount of pre-donated blood and the endogenous red cell reserve:

1. The required amount of pre-donated blood depends on the anticipated blood loss, use of blood conserving procedures and the physical condition of the patient.
This amount should be that quantity which is expected to be sufficient to avoid homologous blood transfusions. The required amount of pre-donated blood is expressed in units whereby one unit in the nomogram is equivalent to 180 ml red cells.
2. The ability to donate blood depends predominantly on the patient's blood volume and baseline PCV. Both variables determine the endogenous red cell reserve, which can be calculated according to the following formula.
Endogenous red cell reserve = blood volume [ml] x (PCV - 33) ÷ 100
Women: blood volume [ml] = 41 [ml/kg] x body weight [kg] + 1200 [ml]
Men: blood volume [ml] = 44 [ml/kg] x body weight [kg] + 1600 [ml]
(body weight ≥ 45 kg)

The indication for Recormon treatment and, if given, the single dose should be determined from the required amount of pre-donated blood and the endogenous red cell reserve according to the following graphs.


Female patients

Required amount of pre-donated blood [units]



Male patients

Required amount of pre-donated blood [units]



The single dose thus determined is administered twice weekly over 4 weeks. The maximum dose should not exceed 1600 IU/kg body weight per week for intravenous or 1200 IU/kg per week for subcutaneous administration.

Prevention of anaemia of prematurity:

The solution is administered subcutaneously at a dose of 3 x 250 IU/kg body weight per week. Recormon treatment should start as early as possible, preferably by day 3 of life. Premature infants who have received a transfusion before starting Recormon treatment are not likely to benefit as much as infants who have not had a transfusion. The treatment should last for 6 weeks.

2.2.1 Special Dosage Instructions

Pediatric use

Results of pediatric clinical studies in children have shown that, on average, the younger the patients, the higher the Recormon doses required. Nevertheless, the recommended dosing schedule should be followed as the individual response cannot be predicted (see section 2.5.4 Pediatric use).

Geriatric use

No dedicated studies in geriatric patients were performed. A large proportion of geriatric patients were included in clinical trials with Recormon. A need for special dose adjustments in the geriatric population was not identified.

Hepatic impairment

No dedicated clinical trials were conducted in patients with hepatic impairment. No special dosage instructions are available.

2.3 Contraindications

Recormon is contraindicated in patients with:

- Known hypersensitivity to the active substance or to any of the excipients.
- Poorly controlled hypertension

In the indication "increasing the yield of autologous blood", Recormon must not be used in patients who, in the month preceding treatment, have suffered a myocardial infarction or stroke, patients with unstable angina pectoris, or patients who are at increased risk of deep venous thrombosis such as those with a history of venous thromboembolic disease.

2.4 Warnings and Precautions

2.4.1 General

In order to improve the traceability of biological medicinal products, the trade name and the batch number of the administered product should be clearly recorded (or stated) in the patient file.

Cancer

- ESAs shortened overall survival and/or time to tumour progression in clinical studies in patients with breast, non-small cell lung, head and neck, lymphoid and cervical cancers when dosed to target a hemoglobin of ≥ 12 g/dl
- The risks of shortened survival and tumor progression have not been excluded when ESAs are dosed to target a hemoglobin of <12g/dl
- To minimize these risks, as well as the risk of serious cardio- and thrombovascular events, use the lowest dose needed to avoid red blood cell transfusions.
- Use only for treatment of anemia due to concomitant myelosuppressive chemotherapy
- Discontinue following the completion of a chemotherapy course.

Recormon should be used with caution in the presence of refractory anaemia with excess blasts in transformation, epilepsy, thrombocytosis and chronic liver failure. Folic acid and vitamin B₁₂ deficiencies should be ruled out as they reduce the effectiveness of Recormon.

In order to ensure effective erythropoiesis, iron status should be evaluated for all patients prior to and during treatment, and supplementary iron therapy may be necessary and conducted in accordance with therapeutic guidelines.

Recormon contains phenylalanine as an excipient. Therefore, this should be taken into consideration in patients affected with severe forms of phenylketonuria.

The indication for Recormon treatment of nephrosclerotic patients not yet undergoing dialysis should be defined individually as a possible acceleration of progression of renal failure cannot be ruled out with certainty.

Lack of effect: The most common reasons for incomplete response to ESAs are iron deficiency and chronic inflammation (e.g. due to uraemia or advanced metastatic cancer). The following conditions may also compromise the effectiveness of ESAs therapy: chronic blood loss, bone marrow fibrosis, severe aluminium overload due to treatment of renal failure, folic acid or vitamin B₁₂ deficiencies, and haemolysis. If all the conditions mentioned are excluded and the patient has a sudden drop of haemoglobin associated with reticulocytopenia and anti-erythropoietin antibodies, examination of the bone marrow for the diagnosis of Pure Red Cell Aplasia (PRCA) should be considered. If PRCA is diagnosed, therapy with epoetin beta must be discontinued and patients should not be switched to another ESA.

Pure red cell aplasia caused by neutralising anti-erythropoietin antibodies has been reported in association with erythropoietin therapy, including Recormon. These antibodies have been shown to cross-react with all erythropoietic proteins, and patients suspected or confirmed to have neutralising antibodies to erythropoietin should not be switched to Recormon (see section 2.6 Undesirable Effects).

Effect on tumour growth

Epoetins are growth factors that primarily stimulate red blood cell production. Erythropoietin receptors may be expressed on the surface of a variety of tumour cells. As with all growth factors, there is a concern that epoetins could stimulate the growth of any type of malignancy.

A controlled clinical study in which epoetin beta was administered to patients with head and neck cancer, has shown a shorter locoregional progression free survival in patients receiving epoetin beta. Another clinical study in breast cancer designed to show a positive effect of epoetin beta on overall survival compared to untreated controls, showed no statistically significant effects in terms of overall survival or tumor progression. Furthermore, meta-analysis data from randomised, controlled clinical studies with epoetin beta in treatment of anaemia in cancer patients (12 studies, 2301 patients; including the two studies mentioned above) did not show any statistically significant negative effects on survival or tumor progression (see section 3.1.2 Clinical / Efficacy Studies).

In CKD patients and patients with cancer receiving chemotherapy an increase in blood pressure (hypertensive episodes) or aggravation of existing hypertension, especially in cases of rapid Hb increase can occur. Increases in blood pressure can be treated with antihypertensive drugs. If blood pressure rises cannot be controlled by drug therapy, a transient interruption of Recormon therapy is recommended. Particularly at the beginning of therapy, regular monitoring of the blood pressure is recommended, including between dialyses in patients with renal anaemia. In patients with CKD, hypertensive crisis with encephalopathy-like symptoms may also occur in individual patients with otherwise normal or low blood pressure. This requires the immediate attention of a physician and intensive medical care. Particular attention should be paid to sudden stabbing migraine-like headaches as a possible warning sign.

Severe aluminium overload due to treatment of renal failure may compromise the effectiveness of Recormon.

In CKD patients an increase in heparin dose during haemodialysis is frequently required during the course of therapy with Recormon as a result of the increased Hb. Occlusion of the dialysis system is possible if heparinisation is not optimal. Early shunt revision and thrombosis prophylaxis by administration of acetylsalicylic acid, for example, should be considered in CKD patients at risk of shunt thrombosis.

In CKD patients there may be a moderate dose-dependent rise in the platelet count within the normal range during treatment with Recormon, especially after intravenous administration. This regresses during the course of continued therapy. It is recommended that the platelet count be monitored regularly during the first 8 weeks of therapy.

In patients in an autologous blood pre-donation programme there may be an increase in platelet count, mostly within the normal range. Therefore, it is recommended that the platelet count be determined at least once a week in these patients. If there is an increase in platelets of more than 150 x 10⁹/L or if platelets rise above the normal range, treatment with Recormon should be discontinued.

For use of Recormon in an autologous pre-donation programme, the official guidelines on principles of blood donation must be considered, in particular:

- only patients with a PCV $\geq 33\%$ (haemoglobin ≥ 11 g/dl [6.83 mmol/L]) should donate;
- special care should be taken with patients below 50 kg weight;
- the single volume drawn should not exceed approx. 12 % of the patient's estimated blood volume.

Treatment should be reserved for patients in whom it is considered of particular importance to avoid homologous blood transfusion taking into consideration the risk/benefit assessment for homologous transfusions.

In patients treated for anaemia of prematurity, there may be a slight rise in platelet counts, particularly up to day 12 - 14 of life, therefore platelets should be monitored regularly.

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), which can be life-threatening or fatal, have been reported in association with epoetin treatment. More severe cases have been observed with long-acting epoetins. At the time of prescription patients should be advised of the signs and symptoms and monitored closely for skin reactions. If signs and symptoms suggestive of these reactions appear, Recormon should be withdrawn immediately and an alternative treatment considered. If the

patient has developed a severe cutaneous skin reaction such as SJS or TEN due to the use of Recormon, treatment with Recormon must not be restarted in this patient at any time.

Laboratory tests

Platelet counts and haematocrit/haemoglobin levels should be monitored at regular intervals in all patients.

In patients with chronic kidney disease, serum potassium elevation has been reported in patients receiving Recormon, though causality has not been established. If an elevated or rising potassium level is observed, then consideration should be given to interrupting Recormon administration until the level has been corrected.

2.4.2 Drug Abuse and Dependence

Misuse by non-anaemic persons may lead to an excessive increase in Hb. This may be associated with life-threatening complications of the cardiovascular system.

There are no reports on dependence when using epoetin beta.

2.4.3 Ability to Drive and Use Machines

Recormon has no or negligible influence on the ability to drive and use machines.

2.5 Use in Special Populations

2.5.1 Pregnancy

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 3.3 Nonclinical Safety).

For epoetin beta, all safety information with regard to exposure to Recormon during pregnancies has been gained from post marketing experience. A review of the available post marketing data does not show evidence of a causal association between harmful effects with respect to pregnancy, embryonal/foetal development or postnatal development and treatment with Recormon. However in the absence of clinical study data, caution should be exercised when prescribing to pregnant women.

2.5.2 Labor and Delivery

Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, embryonal/foetal development, parturition or postnatal development (see section 3.3 Nonclinical Safety).

For epoetin beta, all safety information with regard to exposure during labor and delivery has been gained from post marketing experience. No evidence of harmful effects with respect to labor and delivery have been observed.

However in the absence of clinical study data, caution should be exercised when prescribing to pregnant women in labor.

2.5.3 Lactation

Only limited experience in human lactation has been gained. Endogenous erythropoietin is excreted in breast milk and readily absorbed by the neonatal gastrointestinal tract. A decision on whether to continue or discontinue breastfeeding or to continue or discontinue therapy with epoetin beta should be made taking into account the benefit of breastfeeding to the child and the benefit of epoetin beta therapy to the woman.

2.5.4 Pediatric Use

Clinical registration trials have been performed in children and adolescents with anemia due to chronic kidney disease and in neonates for prevention of anaemia due to prematurity.

In the indication anemia due to chronic kidney disease, Recormon should not be used in infants (i.e. below 2 years of age) (see sections 2.2.1 Special Dosage Instructions, 2.4.1 Warnings and Precautions, General).

In the indications anaemia in cancer patients receiving chemotherapy and treatment for increasing the amount of autologous blood, Recormon is not indicated in the paediatric population.

2.5.5 Geriatric Use

(see section 2.2.1 Special Dosage Instructions).

2.5.6 Renal Impairment

(see section 2.4.1 General, Warnings and Precautions).

2.5.7 Hepatic Impairment

(see section 2.2.1 Special Dosage Instructions).

2.6 Undesirable Effects

2.6.1 Clinical Trials

Based on results from clinical trials including 1725 patients approximately 8 % of patients treated with Recormon are expected to experience adverse drug reactions.

Tabulated summary of adverse drug reactions from clinical trials:

Adverse drug reactions from clinical trials (Table 1, Table 2 and Table 3) are listed by MedDRA system organ class. The corresponding frequency category for each adverse drug reaction is based on the following convention: very common (≥1/10), common (≥1/100 to <1/10), uncommon (≥1/1,000 to <1/100), rare (≥1/10,000 to <1/1000), very rare (<1/10,000).

– [Anaemic patients due to chronic kidney disease](#)

The most frequent adverse drug reactions (common 1-10%), in particular during the early treatment phase with Recormon are hypertensive events including hypertension, hypertensive crisis with or without encephalopathy-like symptoms (e.g. headaches and confused state, sensorimotor disorders - such as speech disturbance or impaired gait - up to tonic-clonic seizures). These increases in blood pressure can occur in normotensive patients or can be an aggravation of existing hypertension (see section 2.4.1 Warnings and Precautions, General).

Shunt thromboses may occur, especially in patients who have a tendency to hypotension or whose arteriovenous fistulae exhibit complications (e.g. stenoses, aneurisms) (see section 2.4.1 Warnings and Precautions, General). In most cases, a fall in serum ferritin values simultaneous with a rise in Hb is observed. In addition, transient increases in serum potassium and phosphate levels have been observed in isolated cases. These parameters should be monitored regularly.

In very rare cases, neutralising anti-erythropoietin antibodies with or without pure red cell aplasia (PRCA) occurred during rHuEPO therapy. In case PRCA is diagnosed, therapy with erythropoietin must be discontinued and patients should not be switched to another erythropoietic substance.

The incidences of adverse drug reactions in clinical trials are shown in the table below. The table shows the difference in frequencies of adverse events between patients receiving Recormon and control.

Table 1: Adverse drug reactions occurring in anaemic patients due to chronic kidney disease, treated with Recormon

System Organ Class	Frequency Category
Vascular Disorders	
Hypertension	Common
Hypertensive crisis	Uncommon
Nervous System Disorders	
Headache	Common
Blood and Lymphatic System Disorders	
Shunt thrombosis	Rare
Thrombocytosis	Very rare

– [Cancer patients receiving chemotherapy with symptomatic anaemia](#)

Hypertensive events are common (1-10%) adverse drug reactions, in particular during the early phase of treatment.

In some patients, a fall in serum iron parameters is observed.

Clinical studies have shown a higher frequency of thromboembolic events in cancer patients treated with Recormon compared to untreated controls or placebo. In patients treated with Recormon, this incidence is 7 % compared to 4 % in controls (both “common”); this is not associated with any increase in thromboembolic mortality compared with controls.

The incidences of adverse drug reactions in clinical trials are shown in the table below. The table shows the difference in frequencies of adverse events between patients receiving Recormon and control.

Table 2: Summary of adverse drug reactions occurring in cancer patients receiving chemotherapy with symptomatic anaemia treated with Recormon

System Organ Class	Frequency Category
Vascular Disorders	
Hypertension	Common
Blood and Lymphatic System Disorders	
Thromboembolic event	Common
Nervous System Disorders	
Headache	Common

– [Patients in an autologous blood pre-donation programme](#)

Patients in an autologous blood pre-donation program have been reported to show a slightly higher frequency of thromboembolic events. However, a causal relationship with treatment with Recormon could not be established.

A temporary iron deficiency may occur, (see section 2.4.1 Warnings and Precautions, General).

The incidences of adverse drug reactions in clinical trials are shown in the table below. The table shows the difference in frequencies of adverse events between patients receiving Recormon and control.

Table 3: Summary of adverse drug reactions occurring in patients in autologous blood pre-donation programme treated with Recormon

System Organ Class	Frequency Category
Nervous System Disorders	
Headache	Common

– [Premature infants](#)

A fall in serum ferritin values is very common (>10%) (see section 2.4.1 Warnings and Precautions, General).

All indications

Rarely (≥1/10,000 to ≤1/1,000), skin reactions such as rash, pruritus, urticaria or injection site reactions may occur. In very rare cases (≤1/10,000) anaphylactoid reactions have been reported. However, in controlled clinical studies no increased incidence of hypersensitivity reactions was found.

In very rare cases (≤1/10,000), particularly when starting treatment, flu-like symptoms such as fever, chills, headaches, pain in the limbs, malaise and/or bone pain have been reported. These reactions were mild or moderate in nature and subsided after a couple of hours or days.

Laboratory Abnormalities

(see section 2.4.1 Warnings and Precautions, General).

2.6.2 Postmarketing Experience

The following adverse drug reactions have been identified from postmarketing experience with Recormon (Table 4). Adverse drug reactions are listed according to system organ classes in MedDRA and the corresponding frequency category estimation for each adverse drug reaction is based on the following convention: very common (≥1/10), common (≥1/100 to <1/10), uncommon (≥1/1,000 to <1/100), rare (≥1/10,000 to <1/1000), very rare (<1/10,000), unknown (cannot be estimated from the available data).

Table 4: Adverse drug reactions from postmarketing experience

System Organ Class	Frequency Category
Blood and Lymphatic System Disorders	
Aplasia Pure Red Cell ^{1,2}	Unknown
Skin and Subcutaneous Tissue Disorders	
Stevens-Johnson syndrome / toxic epidermal necrolysis ²	Unknown

¹ see section 2.4

² Incidence rate and frequency category cannot be estimated based on available data.

Laboratory Abnormalities

Laboratory abnormalities reported during post marketing reflect the experience gained from clinical trials (see sections 2.4.1 Warnings and Precautions, General and 2.6.1 Clinical Trials).

2.7 Overdose

The therapeutic range of Recormon is wide and individual response to therapy must be considered when Recormon treatment is initiated. Overdose can result in manifestations of an exaggerated pharmacodynamic effect, e.g. excessive erythrocytosis which may be associated with life-threatening complications of the cardiovascular system. In case of excessive haemoglobin levels, Recormon should be temporarily withheld (see section 2.2 Dosage and Administration). If clinically indicated, phlebotomy may be performed.

2.8 Interactions with other Medicinal Products and other Forms of Interaction

No dedicated clinical interaction studies have been performed.

Clinical experience has not given evidence for potential interaction of Recormon with other medicinal products (for more information see also section 3.3.5 Nonclinical Safety).

In animal experiments epoetin did not increase the myelotoxicity of cytostatic medicinal products like etoposide, cisplatin, cyclophosphamide, and fluorouracil.

3. PHARMACOLOGICAL PROPERTIES AND EFFECTS

The biological efficacy of epoetin beta has been demonstrated after intravenous and subcutaneous administration in various animal models *in vivo* (normal and uraemic rats, polycythemic mice, dogs). After administration of epoetin beta, the number of erythrocytes, the Hb values and reticulocyte counts increase as well as the ⁵⁹Fe-incorporation rate.

An increased ³H-thymidine incorporation in the erythroid nucleated spleen cells has been found in vitro (mouse spleen cell culture) after incubation with epoetin beta. Investigations in cell cultures of human bone marrow cells showed that epoetin beta stimulates erythropoiesis specifically and does not affect leucopoiesis. Cytotoxic actions of epoetin beta on bone marrow or on human skin cells were not detected.

After single dose administration of epoetin beta no effects on behaviour or locomotor activity of mice and circulatory or respiratory function of dogs were observed.

3.1 Pharmacodynamic Properties

Epoetin beta is identical in its amino acid and carbohydrate composition to erythropoietin that has been isolated from the urine of anaemic patients. Erythropoietin is a glycoprotein that stimulates the formation of erythrocytes from its committed progenitors. It acts as a mitosis stimulating factor and differentiation hormone.

3.1.1 Mechanism of Action

Erythropoietin is a glycoprotein that, as a growth factor, primarily stimulates the formation of erythrocytes from its committed progenitors. It acts as a mitosis stimulating factor and differentiation hormone.

3.1.2 Clinical / Efficacy Studies

This section describes recently completed randomised, controlled studies with epoetin beta in patients with renal anemia or cancer patients receiving chemotherapy/radiotherapy.

Patients with anaemia due to chronic kidney disease

An open randomised study using epoetin beta was conducted in 605 pre-dialysis patients (CREATE) with mild to moderate anemia (Hb level: 11-12.5 g/dl). The primary objective was to explore whether high Hb correction (13–15 g/dl) would reduce cardiovascular (CV) morbidity as compared with standard anaemia treatment (target Hb 10.5–11.5 g/dl). There was no benefit observed with high Hb correction compared to standard anaemia correction. On the contrary, there were fewer events observed in the standard treatment group (47 versus 58 events, HR 0.78, p=0.20). A difference in time to initiation of dialysis was observed favouring the standard anaemia correction group (111 and 127 events, median time to dialysis 41 months and 36 months, log rank test p=0.034, respectively), although no difference in median creatinine clearance over time between the two study groups was observed. Quality of life (assessed by SF-36 Health Survey Questionnaire) was significantly improved (p=0.003) in the high target Hb group at 1 year.

In another open randomised study in 172 patients with early diabetic nephrology, (ACORD) the effect of high Hb correction (target Hb 13-15 g/dl) and standard Hb correction (target Hb 10.5-11.5 g/dl) on cardiac structure and function was investigated.

At the end of the study, there was no significant difference between the two groups with respect to the primary parameter, the baseline adjusted left ventricular mass index (p=0.88). There was no statistically significant difference between the treatment groups in change from baseline in calculated creatinine clearance, time to doubling of serum creatinine, or an analysis of rapid progressors. The General Health score of the quality of life assessment (using the SF-36 Health Survey Questionnaire) was significantly improved (p=0.04) in the high target Hb group.

Cancer patients with symptomatic anaemia receiving chemotherapy

In a placebo-controlled study using epoetin beta in 351 patients with head and neck cancer (ENHANCE), study drug was administered to maintain the haemoglobin levels of 14 g/dl in women and 15 g/dl in men. Locoregional progression free survival was significantly shorter in patients receiving epoetin beta (HR=1.62, p=0.0008). The results and interpretation of this study were confounded by imbalances between the treatment groups, especially with regard to tumor localization, smoking status and the heterogeneity of the study population.

A controlled, open-label, randomised study using epoetin beta in 463 patients with metastatic breast cancer receiving chemotherapy (BRAVE), which was designed to show a significant improvement in survival, did not show any statistically significant difference between the control and epoetin beta arms with regards to overall survival (p=0.52) or time to tumour progression (p=0.45). A greater number of patients in the control arm (64/232; 27.6%) had transfusion and severe anemic events compared with the epoetin beta arm (40/231; 17.3%) (p=0.009), reflecting the efficacy of epoetin beta treatment with respect to preventing transfusions by effective increase in hemoglobin.

A higher percentage of epoetin beta patients experienced thromboembolic events (TEEs) during the study compared with the control arm (13% versus 6%) and a shorter time to TEE for the epoetin beta treatment arm compared with control (p=0.008) was seen. However, the percentage of patients that experienced a serious TEE (3% control versus 4% epoetin beta) or TEE leading to death (2% in each arm) was comparable.

A controlled, open label, randomised study using epoetin beta in 74 patients with cervical cancer receiving radiochemotherapy (MARCH) did not show a correlation between haemoglobin increase and the reduction in treatment failures (response to radiochemotherapy). Therefore, it was decided not to proceed this study to its second stage.

A meta-analysis including all controlled clinical studies in anaemic cancer patients treated with epoetin beta was performed (12 studies with a total of 2301 patients). The results from this present meta-analysis confirm the known efficacy of epoetin beta with respect to increases in haemoglobin levels and a reduced risk of blood transfusion.

In the overall population including also patients with Hb initiation levels up to 13 g/dl, no statistically significant increase in risk of death in the epoetin beta group compared with the control group (HR: 1.13, 95% CI 0.87 to 1.46, p=0.34) was observed. In patients with baseline haemoglobin ≤ 11 g/dl, the HR for overall survival was 1.09 (95 % CI 0.80 to 1.47, p=0.58). For time to disease progression the HR was 0.85 (95% CI: 0.72 to 1.01, p=0.07) in the overall patient population. When the analysis was restricted to patients with baseline haemoglobin ≤ 11 g/dl the HR was 0.80 (95% CI 0.65 to 0.99, p=0.04).

This meta analysis also confirmed the increased rate of thromboembolic events (TEE) reported (see section 2.6.1 Undesirable Effects) with a TEE rate of 7% in the epoetin beta group compared with 4% in the control group.

3.1.3 Immunogenicity

(see section 2.4.1 Warnings and Precautions, General).

3.2 Pharmacokinetic Properties

Pharmacokinetic investigations in healthy volunteers and uraemic patients show that the half-life of intravenously administered epoetin beta is between 4 and 12 hours and that the distribution volume corresponds to one to two times the plasma volume. Analogous results have been found in animal experiments in uraemic and normal rats.

3.2.1 Absorption

After subcutaneous administration of epoetin beta to uraemic patients, the protracted absorption results in a serum concentration plateau, whereby the maximum concentration is reached after an average of 12 - 28 hours.

Bioavailability of epoetin beta after subcutaneous administration is between 23% and 42% as compared with intravenous administration.

3.2.2 Distribution

Pharmacokinetic investigations in healthy volunteers and uraemic patients show that the distribution volume corresponds to one to two times the plasma volume.

3.2.3 Metabolism

Not applicable.

3.2.4 Elimination

Pharmacokinetic investigations in healthy volunteers and uraemic patients show that the half-life of intravenously administered epoetin beta is between 4 and 12 hours.

After subcutaneous administration of epoetin beta to uraemic patients, the terminal half-life is higher than after intravenous administration, with an average of 13 - 28 hours.

3.2.5 Pharmacokinetics in Special Populations

No formal study of the effect of hepatic impairment on the pharmacokinetics of epoetin beta was conducted.

3.3 Nonclinical Safety

3.3.1 Carcinogenicity

A carcinogenicity study with homologous erythropoietin in mice did not reveal any signs of proliferative or tumorigenic potential.

3.3.2 Genotoxicity

Not applicable.

3.3.3 Impairment of Fertility

Not applicable.

3.3.4 Reproductive Toxicity

Not applicable.

3.3.5 Other

Nonclinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, and toxicity of reproduction.

4. PHARMACEUTICAL PARTICULARS

4.1 Storage

The medicine should not be used after the expiry date shown on the pre-filled syringe, and the pack. Storage: Store at 2-8°C (in a refrigerator).

Keep the pre-filled syringe in the outer carton, in order to protect from light.

For the purpose of ambulatory use, the patient may remove the pre-filled syringe from the refrigerator and store it at room temperature (not above 25°C) for one single period of up to 3 days.

Recormon should not be used after the expiry date (EXP) shown on the pack.

4.2 Special Instructions for Use, Handling and Disposal

Incompatibilities

In the absence of compatibility studies, this medicinal product should not be mixed with other medicinal products.

First wash your hands!

- Remove one syringe from the pack and check that the solution is clear, colourless and practically free from visible particles. Remove the cap from the syringe.
- Remove one needle from the pack, fix it on the syringe and remove the protective cap from the needle.
- Expel air from the syringe and needle by holding the syringe vertically and gently pressing the plunger upwards. Keep pressing the plunger until the amount of Recormon is in the syringe is as prescribed.
- Clean the skin at the site of injection using an alcohol wipe. Form a skin fold by pinching the skin between thumb and forefinger. Hold the syringe barrel near to the needle, and insert the needle into the skin fold with a quick, firm action. Inject the Recormon solution. Withdraw the needle quickly and apply pressure over the injection site with a dry, sterile pad.

This medicinal product is for single use only.

Disposal

The following points should be strictly adhered to regarding the use and disposal of syringes and other medicinal sharps:

- Needles and syringes should never be reused.
- Place all used needles and syringes into a sharps container (puncture-proof disposable container).
- Keep this container out of the reach of children.
- Placing used sharps containers in the household waste should be avoided.

Dispose of the full container according to local requirements or as instructed by your healthcare provider.

The release of pharmaceuticals in the environment should be minimized. Medicines should not be disposed of via wastewater, and disposal through household waste should be avoided. Use established 'collection systems' if available in your location.

4.3 Packs

Recormon 2000 IU, 4000 IU, 5000 IU

Syringes with solution for injection

Medicine: keep out of reach of children

MYRecormonPFS20221202CDS7.0

Revision Date: December 2022



Made for

F. Hoffmann-La Roche Ltd, Basel, Switzerland
by Vetter Pharma Fertigung GmbH & Co KG, Ravensburg, Germany.

Made for

F. Hoffmann-La Roche Ltd, Basel, Switzerland
by Roche Diagnostics GmbH, Mannheim, Germany

Instructions for Use Recormon pre-filled syringe

The following instructions explain how to give an injection of Recormon. Be sure that you read, understand and follow the Instructions for Use as well as the package leaflet before injecting Recormon. Your healthcare provider will show you how to prepare and inject Recormon properly before you use it for the first time.

Do not inject yourself unless you have received training. Consult your healthcare provider if you require further information.

Always follow all directions in these Instructions for Use as they may differ from your experiences. These instructions will minimize risk such as accidental needle stick and prevent incorrect use.

Recormon can be administered through 2 manners, your doctor will decide which way is right for you:

- Intravenous administration (into the vein or vein port), only to be performed by healthcare professionals.
- Subcutaneous administration (under the skin).

Before you begin

- Do not** take the needle cap off until you are ready to inject Recormon.
- Do not** try to take the syringe apart at any time.
- Do not** reuse the same syringe.
- Do not** use if the syringe has been dropped or damaged.
- Do not** leave the syringe unattended.
- Keep the syringe and needle and the puncture-resistant or sharps disposal container out of reach of children.
- Contact your healthcare professional if you have any questions.


Storage Instructions

- Keep your unused syringe(s) in the original carton and store in a refrigerator at 2°C to 8°C.
- Keep your syringe and needle out of direct sunlight.
- Do not** freeze.
- Do not** use if the syringe has been frozen.
- Always keep the syringe and needle dry.

Supplies needed to give your injection

Included in the box:

- Recormon pre-filled syringe(s).




- Injection needle(s) (27G ½, used for priming, setting the dose and injecting the medicine.
Note: Each Recormon carton contains 6 syringes/6 needles.
 - Package leaflet with Instructions for Use.
- Not included in the box:**
- 1 alcohol swab.
 - 1 dry sterile pad.
 - 1 puncture-resistant container or sharps container for safe disposal of rubber cap, needle cap and used syringe.

Preparing for injection


- Find a well-lighted, clean, flat, working surface.
 - Take the carton with the syringe(s) and needle(s) out of the refrigerator.
- Check the carton has not been damaged and check the expiration date on the carton has not passed.
 - Do not** use if the expiration date has passed, if the syringe is dropped or damaged, or if the carton appears to be tampered with. In this case, proceed to *step 17* and contact your healthcare provider.
- Take one syringe out of the carton and one needle. Be careful when taking out the syringe. Make sure you always hold the syringe on the glass barrel.
 - Do not** flip the carton upside down to remove the syringe.
 - Do not** handle the syringe by holding the plunger or needle cap.

Remark: If you have a multipack, put the carton with the remaining syringe(s) and needle(s) back into the refrigerator
- Inspect the syringe and needle closely
 - Check the syringe and needle for any damage. **Do not** use the syringe if you have dropped the syringe or if any part of the syringe appears to be damaged.
 - Check the expiration date on the syringe and the needle. **Do not** use the syringe or the needle if the expiration date has passed.
 - Check the liquid in the syringe. The liquid should be clear and colorless. **Do not** use the syringe if the liquid is cloudy, discolored, or has particles.
- Place the syringe on a clean, flat surface.
 - Set aside the syringe for 30 minutes so it can warm up on its own to room temperature. Leave the needle cap on while it warms up.
 - Do not** speed up the warming process in any way, and **do not** put the syringe in a microwave or in warm water.

Remark: If the syringe does not reach room temperature, this could cause the injection to feel uncomfortable and make it hard to push the plunger.




- Attach the needle to the syringe.
 - Remove the needle from its blister.
 - Remove the outer packaging from the needle. **Do not** remove the needle cap.
 - Pull the rubber cap off the end of the syringe.



- Dispose of the rubber cap in a puncture-resistant or sharps container immediately.
- Do not** touch the tip of the syringe.
- Do not** push or pull the plunger.
- Hold the syringe by the barrel and push the needle onto the syringe.
- Gently twist until it is fully attached.

- Place the syringe on a clean, flat surface until ready to use.
- Wash your hands with soap and water.
- Choose an injection site:
 - The recommended injection sites are the top of your thigh or the lower part of your abdomen below the belly button.



- Do not** inject within 5 cm (2 inches) area directly around your belly button.
- Choose a different injection site for each new injection.
- Do not** inject into moles, scars, bruises or areas where the skin is tender, red, hard or not intact.
- Do not** inject into vein or into a muscle.

- Wipe the injection site with an alcohol swab and let it air dry for 10 seconds.




- Do not** fan or blow on the cleaned area.
- Do not** touch the injection site again before giving the injection.

Administering the subcutaneous injection

- Hold the syringe and needle tightly at the hub and carefully pull the injection needle cap away from the syringe. Use the syringe within 5 minutes of removing the cap; otherwise, the needle may clog.
 - Do not** hold the plunger while you remove the needle cap.
 - Do not** touch the needle after removing the needle cap.
 - Do not** re-cap the needle.

- Do not** straighten needle if needle is bent or damaged. Throw away the needle cap in a sharps container immediately.
- Hold the syringe with the needle pointing up. Remove the larger air bubbles by gently tapping the syringe barrel with your fingers until the air bubbles rise to the top of the syringe. Then, slowly push the plunger up to push the air bubbles out of the syringe.



- Adjust to your prescribed dose by slowly pushing the plunger.
- Pinch the selected injection site and fully insert the needle at a 45° to 90° angle with a quick, firm action.
 - Do not** touch the plunger while inserting the needle into the skin.
 - Do not** insert the needle through clothing. Once the needle is inserted, release the pinch and hold the syringe tightly in place.
- Slowly inject your prescribed dose by gently pushing the plunger all the way down.
 - Remove the needle and syringe from the injection site at the same angle as inserted.

After the injection

- There may be a little bleeding at the injection site. You can press dry sterile pad over the injection site. **Do not** rub the injection site.
 - If needed, you may cover the injection site with a small bandage.
 - In case of skin contact with medicine, wash the area that touched the medicine with water.
- Put your used syringe in a sharps disposal container right away after use.
 - Do not** try to remove the used injection needle from the used syringe.
 - Do not** recap the injection needle with the cap.
 - Do not** throw away (dispose of) the syringe in your household trash.

Important: Always keep the sharps disposal container out of reach of children.

Instructions for Use for intravenous injection, intended for healthcare professional only

The following Instructions for Use explain how to give an intravenous injection of Recormon. Be sure that you read, understand and follow the Instructions for Use as well as the package leaflet before injecting Recormon.

Administering the intravenous injection (healthcare professional only)

Preparing for injection: follow steps 1 to 8 of subcutaneous injection (above)

- Select a vein. Change vein with each injection to prevent soreness in one spot.
 - Do not** inject into a red or swollen area.
 - Do not** inject into a muscle.

Clean the skin above the vein with an alcohol swab and let dry.

 - Do not** fan or blow on the cleaned area.
 - Do not** touch the injection site again before giving the injection.
 - Prepare the syringe and needle: follow steps 11 to 14 of subcutaneous injection (above).
 - Insert the needle into the vein.
 - Do not** hold or push on the plunger while inserting the needle.
 - Slowly inject the prescribed dose by gently pushing the plunger all the way down. Remove the needle and syringe from the injection site at the same angle as inserted.

After the intravenous injection, follow steps 16 to 17 of subcutaneous injection (above).
- ###### Administering the intravenous injection via injection port (healthcare professional only)
- Preparing for injection: follow steps 1 to 8 of subcutaneous injection (above)
- Clean the skin above the injection port with an alcohol swab and let dry.

Clean the injection port as instructed by the provider.

 - Do not** fan or blow on the cleaned area.
 - Do not** touch injection site again before giving the injection.
 - Prepare the syringe and needle: follow steps 11 to 14 of subcutaneous injection (above).
 - Insert the needle into the injection port (follow the instruction of the injection port provider).
 - Do not** hold or push on the plunger while inserting the needle.
 - Slowly inject the prescribed dose by gently pushing the plunger all the way down. Remove the needle and syringe from the injection port at the same angle as inserted.
- After the intravenous injection via injection port,** follow steps 16 to 17 subcutaneous injection (above).