

Colimix<sup>®</sup>  
Syrup

KBXXXXXXX  
001

Colimix<sup>®</sup>  
Syrup

Relieves infant colic,  
wind and abdominal  
discomfort



For children and adults

60ml



Colimix<sup>®</sup>  
Syrup

Dosage:

- Infants over 6 months: 5ml before each feed. Do not exceed 4 doses in 24 hours.
- Children 4 to 12 years: 5 to 10ml 4 times daily.
- Adults: 10ml 3 to 4 times daily and at bedtime. Or as advised by a physician.

Caution: It is dangerous to exceed the stated dose.

Dicyclomine is not recommended for use in infants under 6 months of age

Each 5ml contains:  
Dicyclomine HCl 5mg  
Simethicone 50mg  
Methyl Hydroxybenzoate 0.18% w/v and  
Propyl Hydroxybenzoate 0.02% w/v as preservatives.  
Ethanol 96% 0.09ml/5ml

Storage: Store below 30°C. Protect from light.  
Refer to enclosed leaflet for further information.  
SHAKE WELL BEFORE USE



9 556492 1003070



Xepa<sup>SP</sup>

Manufacturer and Product Registration Holder  
(in Malaysia)  
**Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.**  
1-5, Cheng Industrial Estate,  
75250 Melaka, Malaysia.

Colimix<sup>®</sup>  
Syrup

Melegakan kesakitan  
spasma usus pada bayi,  
kembung dan  
ketidakelesaan perut



Untuk kanak-kanak  
dan dewasa

60ml



Colimix<sup>®</sup>  
Syrup

Dos:

- Bayi berumur lebih dari 6 bulan: 5ml sebelum setiap penyusuan. Jangan melebihi 4 dosis dalam tempoh 24 jam.
- Kanak-kanak 4 - 12 tahun: 5 - 10ml 4 kali sehari.
- Dewasa: 10ml 3 - 4 kali sehari dan sebelum waktu tidur.

Atau dinasihatkan oleh para doktor.

Amaran: Adalah merbahaya jika melebihi dos yang dinyatakan.

Dicyclomine tidak harus diberi kepada kanak-kanak berumur kurang dari 6 bulan

Simpan di bawah suhu 30°C. Lindungi dari cahaya.  
GONCANG SEBELUM GUNA

服量:

- 6個月以上嬰兒：餵哺前服用5毫升。每日不可超過4次服量。
- 小童：4至12歲，每次5至10毫升，每日4次。
- 成人：每次10毫升，每日3至4次及睡前服用。或遵照醫生指示。

注意：服食過量有危險。  
置放在不高於30°C陰涼處。遠離光線。

不適用於6個月以下的嬰兒。

服用前請搖均勻。

**CONTROLLED MEDICINE**

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存於兒童的活動範圍以外

FOR ORAL USE ONLY

MAL 19871663AZ

SIN 3333P

HK-29775

MM Reg: No. R2808AA855

BRU 10040454NP

Scan for leaflet  
or package insert



B  
MFG  
EXP



Syrup  
**Colimix**<sup>®</sup>

KBXXXXXXX  
001

**Colimix**<sup>®</sup>  
Syrup

Relieves infant colic,  
wind and abdominal  
discomfort



For children and adults

90ml



**Colimix**<sup>®</sup>  
Syrup

**Dosage:**

- Infants over 6 months: 5ml before each feed. Do not exceed 4 doses in 24 hours.
- Children 4 to 12 years: 5 to 10ml 4 times daily.
- Adults: 10ml 3 to 4 times daily and at bedtime. Or as advised by a physician.

Caution: It is dangerous to exceed the stated dose.

Dicyclomine is not recommended for use in infants under 6 months of age

**Each 5ml contains:**

Dicyclomine HCl 5mg  
 Simethicone 50mg  
 Methyl Hydroxybenzoate 0.18% w/v and  
 Propyl Hydroxybenzoate 0.02% w/v as  
 preservatives.  
 Ethanol 96% 0.09ml/5ml

Storage: Store below 30°C. Protect from light.

Refer to enclosed leaflet for further information.

SHAKE WELL BEFORE USE



9 556492 100225 7



Manufacturer and Product Registration Holder (in Malaysia)  
**Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.**  
 1-5 Cheng Industrial Estate  
 75250 Melaka, Malaysia

**Colimix**<sup>®</sup>  
Syrup

Melegakan kesakitan  
spasma usus pada bayi,  
kembung dan  
ketidakelesaan perut



Untuk kanak-kanak  
dan dewasa

90ml



**Colimix**<sup>®</sup>  
Syrup

**Dos:**

- Bayi berumur lebih dari 6 bulan: 5ml sebelum setiap penyusuan. Jangan melebihi 4 dosis dalam tempoh 24 jam.
- Kanak-kanak 4-12 tahun: 5 - 10ml 4 kali sehari.
- Dewasa: 10ml 3 - 4 kali sehari dan sebelum waktu tidur.

Atau dinasihatkan oleh para doktor.

Amaran: Adalah merbahaya jika melebihi dos yang dinyatakan.

Dicyclomine tidak harus diberi kepada kanak-kanak berumur kurang dari 6 bulan

Simpan di bawah suhu 30°C. Lindungi dari cahaya.

**GONCANG SEBELUM GUNA**

**服量:**

- 6个月以上婴儿: 喂奶前服用5毫升。每日不可超过4次服量。
- 小童: 4至12岁, 每次5至10毫升, 每日4次。
- 成人: 每次10毫升, 每日3至4次及在睡前服用。或遵照医生指示。

注意: 服食过量有危险。置放在不高于30°C阴凉处。远离光线。

不適用於6個月以下的嬰兒。

服用前請搖均勻。

**CONTROLLED MEDICINE**

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存於兒童的活動範圍以外

FOR ORAL USE ONLY

MAL 19871663AZ

SIN 3333P

HK-29775

MM Reg: No.R2808AA855

BRU 10040454NP

Scan for leaflet  
or package insert



B  
MFG  
EXP

