

Each 5 ml contains:

Dicyclomine HCl	5 mg
Simethicone	50 mg
Parabens 0.25% w/v as preservative	

Storage:

Store below 30°C. Protect from light.

Refer to accompanying package insert for further information.

SHAKE WELL BEFORE USE

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

储存于儿童的活动范围以外。

FOR ORAL USE ONLY



Xepa^{SP}

Manufactured and Marketed by

Xepa-Soul Pattinson (M) Sdn Bhd

1-5, Cheng Industrial Estate,
75250 Melaka, Malaysia.

Colimix[®]

Syrup



60ml

Dosage:

- Infants over 6 months:
5ml before each feed. Do not exceed 4 doses in 24 hours.
 - Children 4 to 12 years:
5 to 10ml 4 times daily.
 - Adults: 10ml 3 to 4 times daily and at bedtime.
- Or as advised by a physician.

Caution: It is dangerous to exceed the stated dose.

注意：服食过量有危险。

Dicyclomine is not recommended for use in infants under 6 months of age

MAL 19871663AZ

SIN 3333P HK-29775

MM Reg : No. R1808AA855

BRU 10040454P

CODE CO0101

CO0101

BATCH

MFG EXP

003

Each 5ml contains:
Dicyclomine HCl 5mg
Simethicone 50mg
Parabens 0.25%w/v as preservative

Storage:

Store below 30°C. Protect from light.

Refer to accompanying package insert
for further information.

SHAKE WELL BEFORE USE

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

儲存于兒童的活動範圍以外

FOR ORAL USE ONLY



Manufactured and Marketed by
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate
75250 Melaka, Malaysia

Colimix® Syrup



90ml

Dosage:

- Infants over 6 months: 5ml before each feed. Do not exceed 4 doses in 24 hours.
 - Children 4 to 12 years: 5 to 10ml 4 times daily
 - Adults: 10ml 3 to 4 times daily and at bedtime.
- Or as advised by a physician.

Caution: It is dangerous to exceed the stated dose.

注意：服食过量有危险。

Dicyclomine is not recommended for use in infants under 6 months of age

MAL 19871663AZ
SIN 3333P HK-29775
MM Reg : No. R1808AA855
BRU 10040454P
CODE CO0104

CO0104

BATCH
MFG EXP

003