

Important Information. Please read carefully.

NOVOMIN[®]

SYRUP

COMPOSITION

Each 5ml contains Dimenhydrinate 15mg, Sodium Benzoate 0.13%w/v as preservative and Ethanol 96% 0.05ml/5ml.

PHARMACODYNAMICS/ PHARMACOKINETICS

Dimenhydrinate has antihistaminic, antiemetic, anticholinergic and local anaesthetic properties. Dimenhydrinate is rapidly absorbed from the gastrointestinal tract, metabolised in the liver and excreted mainly as metabolites in the urine.

INDICATIONS

Novomin has anti-emetic properties and is used for the prevention and treatment of nausea and vomiting, such as in travel or motion sickness.

DOSAGE AND ADMINISTRATION

To be administered orally. Unless otherwise prescribed by the physician:

For prevention of travel sickness, the first dose should be taken 30 minutes before the journey. Dosage can be repeated every 8 hours if necessary.

Children 2 to 6 years	:	2.5ml to 5ml
Children 7 to 12 years	:	10ml to 15ml

CONTRAINDICATIONS

It should not be used in persons with known allergy to this drug.

PRECAUTIONS

This medicine may cause drowsiness, If affected, do not drive motor vehicles or operate machinery. Avoid alcoholic drinks.

SIDE EFFECTS

Drowsiness may be experienced by some patients, especially on higher dosage.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Convulsion may occur with massive overdose. No specific antidote is available. The stomach should be emptied by aspiration and lavage. Convulsion may be controlled with diazepam.

SHELF-LIFE

The expiry date is indicated on the packaging.

PRESENTATION

Yellow, orange flavoured syrup in pack of 60ml.

STORAGE

Store below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 23-Oct-2025

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn. Bhd.

1-5, Cheng Industrial Estate,
75250 Melaka, Malaysia.

KLFXXXXXX

XXX