

NOVOMIN

Tablet: Each tablet contains Dimenhydrinate 50mg.

Syrup: Each 5ml contains Dimenhydrinate 15mg.

What is in this leaflet

1. What Novomin is used for
2. How Novomin works
3. Before you use Novomin
4. How to use Novomin
5. While you are using it
6. Side effects
7. Storage and Disposal of Novomin
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What Novomin is used for

It helps in the prevention and treatment of nausea, vomiting and motion sickness.

How Novomin works

Dimenhydrinate has the antihistamine property which prevents the release of histamine due to allergic reaction.

Histamine is a chemical substance that released in the body as part of an allergic reaction.

Together with its antiemetic, anticholinergic and local anaesthetic properties, it blocks the signal of nausea, vomiting and motion sickness in the brain.

Before you use Novomin

- When you must not use it

Do not take this medicine if:

- you are allergic to Dimenhydrinate or any other ingredients of this medicine (listed in Product Description)

- Before you start to use it

Check with your doctor or pharmacist before taking Novomin if you:

- are pregnant or plan to pregnant.
- are breast-feeding or plan to start breast-feeding.
- are going to drive motor vehicles or operate machinery.
- have taken alcoholic drinks or other drugs that may produce drowsiness.

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

How to use Novomin

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

Dosage can be repeated every 8 hours if necessary.

Novomin Tablet

- *Adults:* 1 to 2 tablets
- *Children over 12 years:* 1 tablet
- *Children 7 to 12 years:* ½ to 1 tablet.
- *Children 2 to 6 years:* Recommended for Novomin Syrup.

Novomin Syrup

- *Adults and Children over 12 years:* Recommended for Novomin Tablet
- *Children 7 to 12 years:* 10ml to 15ml
- *Children 2 to 6 years:* 2.5ml to 5ml.

- When to use it

To be taken before or after meals.

For prevention of motion sickness, the first dose should be taken 30 minutes before the journey.

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking Novomin for as long as your doctor recommends.

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do

not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Overdose may result in sudden and irregular movement of muscles.

While you are using Novomin

- Things you must do

- Always wash your hand before handle the medicine.
- Take the medicine at the same time each day.
- For syrup, sip and swallow the medicine slowly, preferably taken without dilution.
- Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

- Do not chew the tablets. Swallow it with water.
- Do not stop taking the medicine unless advised by your doctor.
- Do not take any new medicines without consulting your doctor.
- Do not give Novomin to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If this medicine makes you feel sick, dizzy or tired, or this medicine gives you a headache do not drive or use machines and contact your doctor immediately.

Side effects

Drowsiness may be experienced by some patients, especially on higher dose.

Like all medicines, Novomin can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Novomin

- Storage

Keep out of the reach of the children.
Store below 30°C.
Do not use this product after the expiry date.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Novomin Tablet

White colour, round, convex tablet with single breakline on one side, 7.9mm in diameter.
Available in blister strips of 10's in packing of 10 tablets per box.

Novomin Syrup

Yellow, orange flavoured syrup.
Available in pack of 60ml.

- Ingredients

Novomin Tablet

- Active Ingredient
Dimenhydrinate

- Inactive ingredients
Maize Starch, Lactose, Colloidal Silicon Dioxide, Magnesium Stearate, Microcrystalline Cellulose.

Novomin Syrup

- Active Ingredient
Dimenhydrinate

- Inactive ingredients
Saccharin Sodium, Orange Flavour, Syrup, Purified Water, Ethanol 96%, Glycerin BP, Sodium Benzoate, Quinoline Yellow Colour Powder, Sunset Yellow Colour Powder.

- MAL numbers:

Novomin Tablet: MAL19870329AZ
Novomin Syrup: MAL19870318AZ

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision

31/10/2025

Serial Number

NPRA(R2/2)260414/00042