

Each Tablet Contains Dimenhydrinate 50mg

NOVOMIN

Each Tablet Contains Dimenhydrinate 50mg

10 TABLETS



FOR TRAVEL SICKNESS

NOVOMIN

NOVOMIN

Each Tablet Contains Dimenhydrinate 50mg

FOR TRAVEL SICKNESS

Take the first dose 30 minutes before starting your journey

INDICATIONS

- For prevention and treatment of nausea and vomiting such as in travel or motion sickness.
- Adults - 1 to 2 tablets. Children over 12 years - 1 tablet, Children 7 to 12 years - half to 1 tablet.

DOSE

SHORT JOURNEY

- A single dose is sufficient.

LONG JOURNEY

- You may repeat the dose every 8 hours.

CAUTION

- This medicine may cause drowsiness. If affected, do not drive or operate machinery.

STORAGE

- Store below 30°C.

Refer to accompanying package insert for further information.

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK

MAL 19870329A
SIN 3403P
HK 35601

CODE NO0401
Medico Trading
CAM 813 08 T

Manufactured by
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.

須在啟程卅分鐘之前服用

功效：NOVOMIN能消除嘔吐，適合用於預防及治療噁心及嘔吐，例如旅途所引起的嘔吐（暈車）。

服法：成年人：一至二粒；十二歲以上的兒童：一粒；七至十二歲兒童：半至一粒。

短途：服用一次。

長途：若必要時，每八小時重複服用。

注意：此藥可使人昏昏欲睡，服後如有此情形，不得駕駛或動用機械。

若須進一步詳情，請詢問您的藥劑師或醫生。

HOLOGRAM



9 556492 001908